



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:44 pm GMT

PDB ID : 4MUI  
Title : Crystal structure of pantothenate synthetase in complex with 2-(5-methoxy-2-(4-methoxyphenylsulfonylcarbamoyl)-1H-indol-1-yl)acetic acid  
Authors : Silvestre, H.L.; Blundell, T.L.  
Deposited on : 2013-09-22  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	
1	B	300	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EOH	A	401	-	-	X	X
2	EOH	A	402	-	-	-	X
2	EOH	A	403	-	-	-	X
2	EOH	A	404	-	-	-	X
3	2DQ	B	403	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pantothenate synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	3	0
			2108	1337	378	387	6			
1	B	272	Total	C	N	O	S	0	2	0
			2016	1277	366	368	5			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ALA	THR	ENGINEERED MUTATION	UNP P0A5R0
A	77	GLY	GLU	ENGINEERED MUTATION	UNP P0A5R0
B	2	ALA	THR	ENGINEERED MUTATION	UNP P0A5R0
B	77	GLY	GLU	ENGINEERED MUTATION	UNP P0A5R0

- Molecule 2 is ETHANOL (three-letter code: EOH) (formula: C<sub>2</sub>H<sub>6</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 3	C 2	O 1	0	0
2	A	1	Total 3	C 2	O 1	0	0
2	A	1	Total 3	C 2	O 1	0	0
2	A	1	Total 3	C 2	O 1	0	0
2	B	1	Total 3	C 2	O 1	0	0

- Molecule 3 is (5-METHOXY-2-[(4-METHOXYPHENYL)SULFONYL]CARBAMOYL}-1 H-INDOL-1-YL)ACETIC ACID (three-letter code: 2DQ) (formula: C<sub>19</sub>H<sub>18</sub>N<sub>2</sub>O<sub>7</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 29	C 19	N 2	O 7	S 1	0	0
3	B	1	Total 29	C 19	N 2	O 7	S 1	0	0
3	B	1	Total 29	C 19	N 2	O 7	S 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $\text{C}_2\text{H}_6\text{O}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	148	Total	O	0	0
			148	148		
5	B	138	Total	O	0	0
			138	138		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Pantothenate synthetase



#### • Molecule 1: Pantothenate synthetase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.97Å 70.69Å 81.69Å 90.00° 99.91° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 28.57 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.10) 100.0 (28.57-2.10)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.53 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.186 , 0.251 0.186 , 0.249	Depositor DCC
$R_{free}$ test set	1631 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.4	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 57.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4524	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EOH, 2DQ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.93	3/2146 (0.1%)	0.87	2/2931 (0.1%)
1	B	0.94	2/2053 (0.1%)	0.86	2/2803 (0.1%)
All	All	0.93	5/4199 (0.1%)	0.87	4/5734 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	GLU	CG-CD	-6.06	1.42	1.51
1	A	184	VAL	CB-CG1	5.30	1.64	1.52
1	A	51	VAL	CB-CG1	5.26	1.63	1.52
1	B	143	VAL	CB-CG2	5.25	1.63	1.52
1	B	182	VAL	CB-CG1	5.04	1.63	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	278	ARG	NE-CZ-NH1	-6.17	117.21	120.30
1	B	280	LEU	CA-CB-CG	6.09	129.32	115.30
1	A	56	ARG	NE-CZ-NH2	5.46	123.03	120.30
1	A	274	LEU	CA-CB-CG	-5.23	103.27	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	289	THR	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2108	0	2153	37	0
1	B	2016	0	2081	37	0
2	A	12	0	24	2	0
2	B	3	0	6	0	0
3	A	29	0	17	2	0
3	B	58	0	34	5	0
4	A	4	0	6	1	0
4	B	8	0	12	2	0
5	A	148	0	0	5	0
5	B	138	0	0	3	0
All	All	4524	0	4333	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:ARG:HG2	1:B:278:ARG:HH11	1.32	0.92
1:A:115:ARG:O	2:A:401:EOH:H22	1.70	0.92
1:B:278:ARG:HG2	1:B:278:ARG:NH1	1.90	0.83
1:A:114:LEU:HD21	5:A:646:HOH:O	1.83	0.78
1:B:44:HIS:HD2	1:B:46:GLY:H	1.36	0.73
1:B:44:HIS:H	1:B:47:HIS:HD1	1.37	0.72
1:B:221:ALA:HA	1:B:285:ILE:HD13	1.73	0.70
1:B:278:ARG:HH11	1:B:278:ARG:CG	2.05	0.69
1:A:263:ASN:H	1:A:263:ASN:HD22	1.39	0.69
4:B:401:EDO:H22	3:B:402:2DQ:H15	1.75	0.68
1:B:263:ASN:H	1:B:263:ASN:ND2	1.95	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:HIS:HD2	1:A:46:GLY:H	1.44	0.65
1:A:286:GLU:HG2	1:A:290:PHE:CD2	2.33	0.63
1:B:187:VAL:HG12	1:B:195:MET:HG3	1.79	0.63
1:A:151:ARG:HH22	1:B:151:ARG:CD	2.12	0.62
1:A:157:PHE:CZ	1:A:168:ILE:HD13	2.36	0.61
1:A:255:ILE:HG13	5:A:648:HOH:O	2.01	0.60
1:A:151:ARG:HH22	1:B:151:ARG:HD2	1.65	0.60
1:B:52:ARG:O	1:B:56:ARG:HG3	2.01	0.59
1:B:3:ILE:HD13	1:B:26:ALA:HB2	1.84	0.58
1:A:3:ILE:HG22	1:A:22:ASP:HB3	1.84	0.58
1:B:217:LEU:HD23	1:B:268:LEU:HD23	1.87	0.57
1:A:44:HIS:H	1:A:47:HIS:HD1	1.51	0.57
1:B:160:LYS:HE3	5:B:595:HOH:O	2.05	0.56
1:A:117:THR:OG1	2:A:401:EOH:H21	2.05	0.56
1:A:44:HIS:HE1	3:A:405:2DQ:O	1.88	0.56
1:A:47:HIS:NE2	3:A:405:2DQ:NAQ	2.50	0.56
1:B:263:ASN:N	1:B:263:ASN:ND2	2.54	0.55
1:B:254:ASP:HB3	1:B:261:PRO:HD3	1.87	0.55
1:B:278:ARG:NH1	1:B:278:ARG:CG	2.61	0.54
1:A:197:SER:O	1:A:200:ARG:HG2	2.09	0.52
1:A:286:GLU:HG2	1:A:290:PHE:CG	2.43	0.52
1:A:24:SER:OG	1:A:28:ARG:NH2	2.42	0.52
1:A:174:ASP:HB3	4:B:404:EDO:H21	1.92	0.51
1:B:46:GLY:O	3:B:402:2DQ:H7	2.11	0.50
3:B:403:2DQ:H3	3:B:403:2DQ:OAD	2.12	0.50
1:A:160[A]:LYS:HE3	5:A:528:HOH:O	2.11	0.49
1:A:114:LEU:HD22	1:A:117:THR:HG22	1.94	0.49
1:A:286:GLU:HB3	1:A:290:PHE:HB3	1.94	0.49
1:A:27:LEU:HD13	1:A:61:VAL:HG11	1.94	0.48
3:B:403:2DQ:CA	3:B:403:2DQ:OAD	2.61	0.48
1:B:263:ASN:H	1:B:263:ASN:HD22	1.58	0.48
1:A:69:ASN:HD22	1:A:72:GLN:HE21	1.61	0.47
1:A:131:PRO:HG2	1:A:132:ARG:NH2	2.29	0.47
1:B:124:ALA:O	1:B:129:GLY:HA3	2.15	0.47
1:B:217:LEU:HD23	1:B:268:LEU:CD2	2.44	0.47
1:A:262:LEU:HD21	1:A:287:ILE:HG13	1.97	0.46
1:A:151:ARG:NH2	1:B:151:ARG:HD2	2.29	0.46
1:B:191:ASP:OD2	1:B:211[B]:VAL:CG2	2.64	0.46
1:A:202:LEU:HD22	1:A:206:GLN:HB3	1.98	0.45
1:B:132:ARG:HB3	1:B:135:HIS:HB2	1.99	0.45
1:A:120:PRO:HB2	1:A:124:ALA:HB2	1.99	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:HIS:HE1	3:B:402:2DQ:OXT	2.00	0.44
1:A:268:LEU:HB2	1:A:285:ILE:HG13	2.00	0.43
1:B:191:ASP:OD2	1:B:211[B]:VAL:HG22	2.18	0.43
1:A:290:PHE:HD1	1:A:290:PHE:HA	1.74	0.43
1:A:45:GLU:HG3	1:A:187:VAL:HG11	2.01	0.43
1:A:43:LEU:HA	1:A:47:HIS:HD1	1.84	0.43
1:B:287:ILE:HA	1:B:287:ILE:HD13	1.87	0.43
1:A:221:ALA:HA	1:A:285:ILE:HD13	2.01	0.42
1:B:44:HIS:CE1	1:B:47:HIS:CE1	3.07	0.42
1:A:174:ASP:OD2	1:B:115:ARG:HB3	2.19	0.42
1:B:269:LEU:HD12	1:B:269:LEU:N	2.35	0.42
4:A:406:EDO:H12	5:A:639:HOH:O	2.19	0.41
1:B:148:GLN:OE1	5:B:622:HOH:O	2.22	0.41
1:B:208:ALA:O	1:B:211[A]:VAL:HG12	2.20	0.41
1:A:148:GLN:NE2	5:A:552:HOH:O	2.49	0.41
1:B:37:VAL:HB	1:B:64:VAL:HG22	2.02	0.41
1:A:232:LEU:O	1:A:236:ARG:HG3	2.20	0.41
1:B:142:VAL:CG1	1:B:143:VAL:N	2.84	0.41
1:B:151:ARG:NH2	5:B:604:HOH:O	2.52	0.41
1:A:255:ILE:HG23	1:A:266:GLY:HA2	2.03	0.41
1:B:33:ARG:HH11	1:B:33:ARG:HA	1.85	0.40
1:B:44:HIS:CD2	1:B:46:GLY:H	2.26	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/300 (93%)	271 (97%)	8 (3%)	1 (0%)	38	35
1	B	270/300 (90%)	264 (98%)	6 (2%)	0	100	100
All	All	550/600 (92%)	535 (97%)	14 (2%)	1 (0%)	51	52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/223 (96%)	206 (96%)	9 (4%)	34	33
1	B	205/223 (92%)	198 (97%)	7 (3%)	42	43
All	All	420/446 (94%)	404 (96%)	16 (4%)	38	38

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	HIS
1	A	30	THR
1	A	114	LEU
1	A	236	ARG
1	A	254	ASP
1	A	260	MET
1	A	263	ASN
1	A	289	THR
1	A	290	PHE
1	B	33	ARG
1	B	87	ASP
1	B	88	ASP
1	B	142	VAL
1	B	160	LYS
1	B	263	ASN
1	B	278	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	72	GLN
1	A	222	HIS
1	A	263	ASN
1	B	44	HIS
1	B	263	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EOH	A	401	-	2,2,2	0.51	0	1,1,1	0.17	0
2	EOH	A	402	-	2,2,2	0.55	0	1,1,1	0.38	0
2	EOH	A	403	-	2,2,2	0.55	0	1,1,1	0.23	0
2	EOH	A	404	-	2,2,2	0.51	0	1,1,1	0.31	0
3	2DQ	A	405	-	26,31,31	2.61	9 (34%)	35,45,45	2.55	6 (17%)
4	EDO	A	406	-	3,3,3	0.46	0	2,2,2	0.48	0
4	EDO	B	401	-	3,3,3	0.47	0	2,2,2	0.47	0
3	2DQ	B	402	-	26,31,31	3.01	9 (34%)	35,45,45	2.81	10 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	2DQ	B	403	-	26,31,31	3.41	12 (46%)	35,45,45	2.28	7 (20%)
4	EDO	B	404	-	3,3,3	0.39	0	2,2,2	0.59	0
2	EOH	B	405	-	2,2,2	0.38	0	1,1,1	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EOH	A	401	-	-	0/0/0/0	0/0/0/0
2	EOH	A	402	-	-	0/0/0/0	0/0/0/0
2	EOH	A	403	-	-	0/0/0/0	0/0/0/0
2	EOH	A	404	-	-	0/0/0/0	0/0/0/0
3	2DQ	A	405	-	-	0/18/23/23	0/3/3/3
4	EDO	A	406	-	-	0/1/1/1	0/0/0/0
4	EDO	B	401	-	-	0/1/1/1	0/0/0/0
3	2DQ	B	402	-	-	0/18/23/23	0/3/3/3
3	2DQ	B	403	-	-	0/18/23/23	0/3/3/3
4	EDO	B	404	-	-	0/1/1/1	0/0/0/0
2	EOH	B	405	-	-	0/0/0/0	0/0/0/0

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	2DQ	CAU-NAQ	-5.27	1.32	1.39
3	B	403	2DQ	CAO-CAY	-4.65	1.33	1.38
3	A	405	2DQ	CAO-CAY	-4.54	1.34	1.38
3	A	405	2DQ	CAU-NAQ	-4.32	1.33	1.39
3	B	403	2DQ	CAU-NAQ	-3.61	1.34	1.39
3	B	402	2DQ	CAM-CBA	-3.50	1.33	1.41
3	B	403	2DQ	CAM-CBA	-3.36	1.34	1.41
3	A	405	2DQ	CAZ-CBA	-3.13	1.34	1.41
3	B	402	2DQ	CAN-CAZ	-3.07	1.34	1.42
3	B	403	2DQ	CAN-CAZ	-3.05	1.34	1.42
3	B	403	2DQ	CAZ-CBA	-2.98	1.34	1.41
3	A	405	2DQ	CAM-CBA	-2.75	1.35	1.41
3	A	405	2DQ	CAN-CAZ	-2.55	1.36	1.42
3	B	403	2DQ	CAY-CAU	-2.40	1.35	1.50
3	B	402	2DQ	CAO-CAY	-2.13	1.36	1.38
3	B	402	2DQ	CAZ-CBA	-2.00	1.37	1.41
3	B	403	2DQ	CA-N	2.09	1.52	1.48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	2DQ	CAN-CAW	2.40	1.41	1.37
3	B	402	2DQ	CAX-SBC	2.44	1.80	1.76
3	B	403	2DQ	CAX-SBC	2.45	1.80	1.76
3	B	402	2DQ	CA-N	2.61	1.53	1.48
3	A	405	2DQ	SBC-NAQ	2.93	1.70	1.64
3	A	405	2DQ	CAN-CAW	3.83	1.44	1.37
3	A	405	2DQ	OAE-SBC	4.28	1.48	1.43
3	B	403	2DQ	SBC-NAQ	6.30	1.77	1.64
3	A	405	2DQ	OAF-SBC	7.26	1.51	1.43
3	B	402	2DQ	OAF-SBC	8.57	1.52	1.43
3	B	402	2DQ	OAE-SBC	8.63	1.52	1.43
3	B	403	2DQ	OAE-SBC	9.06	1.53	1.43
3	B	403	2DQ	OAF-SBC	9.27	1.53	1.43

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	405	2DQ	OAE-SBC-OAF	-12.55	103.49	119.55
3	B	402	2DQ	OAE-SBC-OAF	-10.49	106.13	119.55
3	B	403	2DQ	OAE-SBC-OAF	-8.31	108.92	119.55
3	B	403	2DQ	CAU-NAQ-SBC	-7.92	113.36	123.44
3	B	402	2DQ	CAU-NAQ-SBC	-4.11	118.21	123.44
3	B	402	2DQ	OAF-SBC-CAX	-3.00	104.25	107.95
3	B	402	2DQ	CAA-OAR-CAV	-2.72	111.55	117.50
3	B	402	2DQ	CAL-CAW-CAN	-2.59	117.31	120.82
3	B	402	2DQ	OAD-CAU-NAQ	-2.58	117.87	121.11
3	B	403	2DQ	C-CA-N	-2.42	110.57	114.00
3	B	403	2DQ	CAM-CAL-CAW	-2.06	117.39	120.15
3	A	405	2DQ	CAB-OAS-CAW	-2.05	113.02	117.50
3	B	403	2DQ	CAO-CAZ-CBA	2.09	108.10	106.27
3	B	402	2DQ	CAB-OAS-CAW	2.09	122.08	117.50
3	A	405	2DQ	CAY-CAO-CAZ	2.29	109.41	106.55
3	B	403	2DQ	OAE-SBC-NAQ	2.40	113.64	106.75
3	A	405	2DQ	CAX-SBC-NAQ	2.63	109.47	105.91
3	B	402	2DQ	CAK-CAX-SBC	3.00	123.09	119.77
3	B	403	2DQ	OAF-SBC-CAX	3.19	111.90	107.95
3	A	405	2DQ	OAE-SBC-CAX	3.91	112.79	107.95
3	A	405	2DQ	OAF-SBC-NAQ	3.95	118.09	106.75
3	B	402	2DQ	OAF-SBC-NAQ	4.86	120.69	106.75
3	B	402	2DQ	OAE-SBC-CAX	6.92	116.51	107.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	EOH	2	0
3	A	405	2DQ	2	0
4	A	406	EDO	1	0
4	B	401	EDO	1	0
3	B	402	2DQ	3	0
3	B	403	2DQ	2	0
4	B	404	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/300 (94%)	0.27	17 (6%)	23 28	11, 25, 46, 57	4 (1%)
1	B	272/300 (90%)	0.35	25 (9%)	10 12	12, 23, 43, 59	1 (0%)
All	All	555/600 (92%)	0.31	42 (7%)	15 19	11, 24, 45, 59	5 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	262	LEU	8.0
1	A	261	PRO	6.9
1	A	258	GLY	6.8
1	A	81	ALA	5.8
1	B	86	PRO	4.9
1	B	70	PRO	4.8
1	A	3	ILE	4.6
1	A	290	PHE	4.5
1	B	263	ASN	4.4
1	A	95	ALA	4.3
1	B	201	TYR	4.0
1	B	91	ALA	3.9
1	B	87	ASP	3.8
1	A	260	MET	3.7
1	B	130	GLY	3.5
1	B	133	PRO	3.4
1	B	131	PRO	3.2
1	B	30	THR	3.2
1	B	261	PRO	3.0
1	B	200	ARG	3.0
1	A	5	ALA	3.0
1	A	4	PRO	2.9
1	A	90	LEU	2.9
1	B	288	GLY	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	69	ASN	2.8
1	B	190	ALA	2.7
1	B	88	ASP	2.7
1	A	289	THR	2.7
1	B	255	ILE	2.6
1	A	259	PRO	2.5
1	A	79	LEU	2.5
1	B	132	ARG	2.5
1	A	71	MET	2.4
1	B	2	ALA	2.4
1	B	29	LEU	2.3
1	A	263	ASN	2.1
1	B	68	VAL	2.1
1	B	95	ALA	2.1
1	A	10	GLU	2.0
1	B	10	GLU	2.0
1	B	41	GLY	2.0
1	A	276	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	EOH	A	403	3/3	0.87	0.23	15.05	33,33,34,34	0
2	EOH	A	402	3/3	0.84	0.40	7.01	48,48,48,49	0
3	2DQ	B	403	29/29	0.80	0.30	2.84	48,50,55,56	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EOH	A	404	3/3	0.78	0.22	2.66	45,45,46,47	0
2	EOH	A	401	3/3	0.88	0.15	2.58	23,23,32,33	0
2	EOH	B	405	3/3	0.88	0.16	0.97	31,31,32,33	0
4	EDO	A	406	4/4	0.98	0.11	0.81	23,26,26,27	0
3	2DQ	B	402	29/29	0.94	0.15	-0.01	23,27,30,33	0
4	EDO	B	401	4/4	0.85	0.18	-0.11	40,43,45,47	0
3	2DQ	A	405	29/29	0.94	0.12	-0.15	25,28,34,34	0
4	EDO	B	404	4/4	0.98	0.08	-0.94	20,23,25,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.