



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 7, 2018 – 11:09 AM EST

PDB ID : 5MU6  
Title : Human N-myristoyltransferase (NMT1) with Myristoyl-CoA and IMP-1088 inhibitor bound  
Authors : Perez-Dorado, I.; Bell, A.S.; Tate, E.W.  
Deposited on : 2017-01-12  
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736

**i**

## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



<b>Metric</b>	<b>Whole archive (#Entries)</b>	<b>Similar resolution (#Entries, resolution range(Å))</b>
R <sub>free</sub>	100719	7505 (1.90-1.86)
Clashscore	112137	8369 (1.90-1.86)
Ramachandran outliers	110173	8279 (1.90-1.86)
Sidechain outliers	110143	8280 (1.90-1.86)
RSRZ outliers	101464	7571 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	391	<div> <div></div> <div>92%</div> <div>6%</div> <div>..</div> </div>
1	B	391	<div> <div></div> <div>92%</div> <div></div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	B	504	-	-	-	X

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	1	0
			3139	2033	533	557	16			
1	B	377	Total	C	N	O	S	0	6	0
			3128	2030	528	553	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	106	GLY	-	expression tag	UNP P30419
A	107	PRO	-	expression tag	UNP P30419
A	108	HIS	-	expression tag	UNP P30419
B	106	GLY	-	expression tag	UNP P30419
B	107	PRO	-	expression tag	UNP P30419
B	108	HIS	-	expression tag	UNP P30419

- Molecule 2 is TETRADECANOYL-COA (three-letter code: MYA) (formula:  $C_{35}H_{62}N_7O_{17}P_3S$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	
			63	35	7	17	3	1	0
2	B	1	Total	C	N	O	P	S	
			63	35	7	17	3	1	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O		
			6	3	3	0	0
4	A	1	Total	C	O		
			6	3	3	0	0
4	B	1	Total	C	O		
			6	3	3	0	0
4	B	1	Total	C	O		
			6	3	3	0	0

- Molecule 5 is 1-[5-[3,4-bis(fluoranyl)-2-[2-(1,3,5-trimethylpyrazol-4-yl)ethoxy]phenyl]-1-methyl-indazol-3-yl]- {N}, {N}-dimethyl-methanamine (three-letter code: KFK) (formula:

C<sub>25</sub>H<sub>29</sub>F<sub>2</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	0
			33	25	2	5	1		
5	B	1	Total	C	F	N	O	0	0
			33	25	2	5	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	228	Total	O	0	0
			228	228		
6	B	228	Total	O	0	0
			228	228		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1

Chain A: 



- Molecule 1: Glycylpeptide N-tetradecanoyltransferase 1

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.73Å 180.73Å 58.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.18 – 1.88 72.18 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.8 (72.18-1.88) 99.9 (72.18-1.88)	Depositor EDS
$R_{merge}$	0.25	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 1.88Å)	Xtrriage
Refinement program	REFMAC 5.8.0073, PHENIX 1.8.4_1496	Depositor
R, $R_{free}$	0.204 , 0.247 0.210 , 0.249	Depositor DCC
$R_{free}$ test set	3450 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.1	Xtrriage
Anisotropy	0.992	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6941	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MYA, GOL, MG, KFK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3231	0.53	1/4389 (0.0%)
1	B	0.41	0/3228	0.54	0/4381
All	All	0.41	0/6459	0.54	1/8770 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	408	MET	CG-SD-CE	5.43	108.89	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3139	0	3138	15	0
1	B	3128	0	3150	12	0
2	A	63	0	58	0	0
2	B	63	0	58	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	16	0	0
4	B	12	0	16	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	33	0	0	0	0
5	B	33	0	0	0	0
6	A	228	0	0	1	0
6	B	228	0	0	3	0
All	All	6941	0	6436	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:325:LYS:HG3	1:B:328:ARG:HH12	1.47	0.79
1:B:393:GLU:OE2	6:B:601:HOH:O	2.03	0.76
1:B:240:LYS:NZ	6:B:602:HOH:O	2.15	0.75
1:A:410:HIS:HD2	1:A:412:THR:H	1.43	0.67
1:A:127:LYS:N	1:A:130:GLU:OE1	2.28	0.64
1:A:166:ARG:NH1	6:A:604:HOH:O	2.31	0.60
1:A:130:GLU:O	1:A:289:LYS:NZ	2.33	0.59
1:B:145[B]:ILE:HD12	1:B:272:HIS:HB3	1.86	0.58
1:B:325:LYS:HG3	1:B:328:ARG:NH1	2.18	0.55
1:A:143:ASP:O	1:B:350:HIS:ND1	2.37	0.55
1:B:448:ASP:OD1	6:B:603:HOH:O	2.19	0.52
1:B:145[A]:ILE:HD11	1:B:273:LEU:HG	1.92	0.52
1:A:410:HIS:CD2	1:A:413:HIS:H	2.29	0.51
1:A:458:ASN:HA	1:A:461:PHE:CE2	2.46	0.51
1:A:115:ARG:NH2	1:A:117:TYR:OH	2.44	0.51
1:A:410:HIS:CD2	1:A:412:THR:H	2.27	0.50
1:B:458:ASN:HA	1:B:461:PHE:CE2	2.52	0.45
1:A:135:HIS:HA	1:A:482:LYS:O	2.19	0.43
1:A:315:SER:HB3	1:A:318:MET:HB2	2.01	0.43
1:A:189:ARG:NH1	1:A:408:MET:SD	2.91	0.43
1:A:410:HIS:HA	1:A:411:PRO:HD3	1.77	0.42
1:B:170:LYS:HD3	1:B:170:LYS:HA	1.82	0.42
1:B:376:TYR:HA	1:B:377:PRO:HD3	1.92	0.41
1:B:379:GLU:O	1:B:379:GLU:HG3	2.21	0.40
1:A:440:VAL:O	1:A:444:MET:HG3	2.21	0.40
1:A:144:ASN:OD1	1:A:144:ASN:N	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/391 (97%)	371 (97%)	10 (3%)	0	100	100
1	B	379/391 (97%)	368 (97%)	11 (3%)	0	100	100
All	All	760/782 (97%)	739 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	349/355 (98%)	346 (99%)	3 (1%)	82	81
1	B	349/355 (98%)	345 (99%)	4 (1%)	78	75
All	All	698/710 (98%)	691 (99%)	7 (1%)	82	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	ARG
1	A	170	LYS
1	A	455	LEU
1	B	145[A]	ILE
1	B	145[B]	ILE
1	B	304	ARG
1	B	455	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	HIS
1	A	317	ASN
1	A	410	HIS
1	A	428	GLN
1	A	475	GLN
1	B	135	HIS
1	B	351	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	MYA	A	501	3	55,65,65	0.77	1 (1%)	63,91,91	1.66	6 (9%)
4	GOL	A	503	-	5,5,5	0.42	0	5,5,5	0.51	0
4	GOL	A	504	-	5,5,5	0.40	0	5,5,5	0.45	0
5	KFK	A	505	-	33,36,36	2.20	12 (36%)	40,53,53	1.80	9 (22%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MYA	B	501	-	55,65,65	0.80	1 (1%)	63,91,91	1.69	7 (11%)
4	GOL	B	503	-	5,5,5	0.37	0	5,5,5	0.31	0
4	GOL	B	504	-	5,5,5	0.38	0	5,5,5	0.61	0
5	KFK	B	505	-	33,36,36	2.04	9 (27%)	40,53,53	1.79	9 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MYA	A	501	3	-	0/59/80/80	0/3/3/3
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
4	GOL	A	504	-	-	0/4/4/4	0/0/0/0
5	KFK	A	505	-	-	0/13/14/14	0/4/4/4
2	MYA	B	501	-	-	0/59/80/80	0/3/3/3
4	GOL	B	503	-	-	0/4/4/4	0/0/0/0
4	GOL	B	504	-	-	0/4/4/4	0/0/0/0
5	KFK	B	505	-	-	0/13/14/14	0/4/4/4

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	505	KFK	CAV-CAW	-4.14	1.27	1.36
2	B	501	MYA	C2-S1	-4.01	1.76	1.81
5	A	505	KFK	CAS-CAR	-3.98	1.29	1.38
5	A	505	KFK	CAS-CAT	-3.83	1.35	1.42
5	A	505	KFK	CAV-CAW	-3.82	1.28	1.36
5	B	505	KFK	CAS-CAR	-3.73	1.29	1.38
2	A	501	MYA	C2-S1	-3.67	1.76	1.81
5	A	505	KFK	CAV-CAU	-3.54	1.33	1.41
5	B	505	KFK	CAS-CAT	-3.36	1.35	1.42
5	B	505	KFK	CAC-NAB	-3.31	1.32	1.37
5	B	505	KFK	CAV-CAU	-3.31	1.34	1.41
5	A	505	KFK	CAC-NAB	-3.19	1.32	1.37
5	A	505	KFK	CAZ-NBA	2.07	1.36	1.33
5	A	505	KFK	CAQ-CAR	2.42	1.53	1.49
5	B	505	KFK	NAA-NAB	2.60	1.40	1.37
5	A	505	KFK	NAA-NAB	2.70	1.40	1.37
5	B	505	KFK	NBA-NBB	2.79	1.40	1.37
5	A	505	KFK	CBD-CAZ	2.86	1.52	1.51
5	A	505	KFK	NBA-NBB	3.79	1.41	1.37

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	505	KFK	CAD-CAE	4.26	1.47	1.38
5	A	505	KFK	CAC-CAD	4.35	1.48	1.39
5	B	505	KFK	CAC-CAD	4.54	1.48	1.39
5	B	505	KFK	CAD-CAE	4.59	1.48	1.38

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	MYA	N3A-C2A-N1A	-8.99	121.03	128.86
2	A	501	MYA	N3A-C2A-N1A	-7.98	121.91	128.86
5	B	505	KFK	CAV-CAU-CAT	-2.77	117.17	120.94
2	A	501	MYA	C4A-C5A-N7A	-2.76	106.75	109.41
5	A	505	KFK	CAV-CAU-CAT	-2.71	117.26	120.94
5	B	505	KFK	CAR-CAS-CAT	-2.67	117.02	122.30
5	B	505	KFK	CAP-CAQ-CAR	-2.32	114.34	118.68
2	A	501	MYA	C4M-C3M-C2M	-2.21	107.61	113.97
5	A	505	KFK	CAR-CAS-CAT	-2.12	118.12	122.30
2	B	501	MYA	C4A-C5A-N7A	-2.03	107.45	109.41
2	B	501	MYA	C7-C6-C5	-2.01	108.98	112.22
2	B	501	MYA	C6-C7-N8	-2.01	107.72	111.87
5	A	505	KFK	CAR-CAQ-CAI	2.03	125.70	122.10
5	A	505	KFK	CAS-CAT-CAU	2.32	122.43	119.65
2	B	501	MYA	C7-N8-C9	2.37	127.00	122.59
2	A	501	MYA	C7-N8-C9	2.44	127.14	122.59
5	A	505	KFK	CAZ-CBD-NBE	2.47	117.00	113.48
5	B	505	KFK	CAS-CAT-CAU	2.59	122.75	119.65
5	B	505	KFK	CAL-CAE-NAA	2.64	125.48	119.78
5	B	505	KFK	CAR-CAQ-CAI	2.98	127.39	122.10
5	A	505	KFK	CAL-CAE-NAA	3.08	126.43	119.78
5	B	505	KFK	CAZ-NBA-NBB	3.36	107.38	104.48
5	A	505	KFK	CAZ-NBA-NBB	3.37	107.39	104.48
5	A	505	KFK	CAV-CAU-NBB	3.45	135.72	132.40
5	B	505	KFK	CAV-CAU-NBB	3.72	135.97	132.40
2	A	501	MYA	C2-S1-C2M	3.83	106.21	100.36
5	B	505	KFK	CAE-NAA-NAB	4.62	108.33	104.35
2	B	501	MYA	C2-S1-C2M	4.66	107.49	100.36
5	A	505	KFK	CAE-NAA-NAB	5.22	108.84	104.35
2	B	501	MYA	O2M-C2M-C3M	5.30	118.99	109.14
2	A	501	MYA	O2M-C2M-C3M	5.98	120.26	109.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	382/391 (97%)	-0.21	5 (1%) 77 79	7, 12, 30, 58	0
1	B	377/391 (96%)	-0.16	3 (0%) 86 87	7, 12, 29, 57	0
All	All	759/782 (97%)	-0.18	8 (1%) 80 82	7, 12, 30, 58	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	316	ARG	3.4
1	A	315	SER	2.8
1	B	317	ASN	2.8
1	B	115	ARG	2.7
1	A	317	ASN	2.6
1	A	130	GLU	2.4
1	A	115	ARG	2.3
1	B	336	ALA	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	504	6/6	0.88	0.15	6.69	17,18,20,20	0
3	MG	B	502	1/1	0.94	0.17	1.35	21,21,21,21	0
3	MG	A	502	1/1	0.93	0.14	1.24	22,22,22,22	0
5	KFK	A	505	33/33	0.94	0.10	1.20	4,8,11,13	0
4	GOL	A	504	6/6	0.91	0.10	1.10	13,16,21,23	0
5	KFK	B	505	33/33	0.95	0.10	0.27	6,9,12,13	0
4	GOL	A	503	6/6	0.92	0.11	0.23	9,17,21,22	0
4	GOL	B	503	6/6	0.94	0.10	0.13	9,10,14,15	0
2	MYA	A	501	63/63	0.96	0.09	-0.47	5,10,15,20	0
2	MYA	B	501	63/63	0.96	0.09	-0.54	3,11,16,19	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.