



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:30 am GMT

PDB ID : 3MV5  
Title : Crystal structure of Akt-1-inhibitor complexes  
Authors : Pandit, J.  
Deposited on : 2010-05-03  
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4719 (2.50-2.46)
Clashscore	112137	5483 (2.50-2.46)
Ramachandran outliers	110173	5388 (2.50-2.46)
Sidechain outliers	110143	5390 (2.50-2.46)
RSRZ outliers	101464	4754 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	
2	C	10	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2749 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called v-akt murine thymoma viral oncogene homolog 1 (AKT1).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	316	2597	1668	437	476	1	15	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	GLY	-	EXPRESSION TAG	UNP B2RAM5
A	140	ALA	-	EXPRESSION TAG	UNP B2RAM5
A	141	MET	-	EXPRESSION TAG	UNP B2RAM5
A	142	ASP	-	EXPRESSION TAG	UNP B2RAM5
A	143	PRO	-	EXPRESSION TAG	UNP B2RAM5
A	473	ASP	SER	ENGINEERED	UNP B2RAM5


- Molecule 2 is a protein called GSK3-beta peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	10	79	47	16	16	0	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mn	0	0
			1	1		

- Molecule 4 is (3R)-1-(5-METHYL-7H-PYRROLO[2,3-D]PYRIMIDIN-4-YL)PYRROLIDIN-3-AMINE (three-letter code: XFE) (formula: C<sub>11</sub>H<sub>15</sub>N<sub>5</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C N	0	0
			16	11 5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	55	Total	O	0	0
			55	55		
5	C	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: v-akt murine thymoma viral oncogene homolog 1 (AKT1)



- Molecule 2: GSK3-beta peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.71Å 55.42Å 93.29Å 90.00° 104.00° 90.00°	Depositor
Resolution (Å)	19.92 – 2.47 19.92 – 2.47	Depositor EDS
% Data completeness (in resolution range)	96.8 (19.92-2.47) 97.0 (19.92-2.47)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 2.47Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.200 , 0.282 0.200 , 0.280	Depositor DCC
$R_{free}$ test set	758 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.2	Xtrriage
Anisotropy	0.138	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2749	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MN, XFE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.15	7/2647 (0.3%)	1.07	10/3558 (0.3%)
2	C	1.12	0/80	0.90	0/105
All	All	1.15	7/2727 (0.3%)	1.06	10/3663 (0.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	350	TYR	CD2-CE2	8.96	1.52	1.39
1	A	350	TYR	CE2-CZ	6.59	1.47	1.38
1	A	243	ARG	CG-CD	6.08	1.67	1.51
1	A	229	TYR	CD2-CE2	5.57	1.47	1.39
1	A	397	GLU	CG-CD	5.55	1.60	1.51
1	A	359	GLU	CD-OE1	5.33	1.31	1.25
1	A	258	VAL	CB-CG1	5.26	1.64	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	ARG	NE-CZ-NH2	-7.32	116.64	120.30
1	A	350	TYR	CB-CG-CD1	-7.29	116.62	121.00
1	A	350	TYR	CB-CG-CD2	6.71	125.03	121.00
1	A	325	ASP	CB-CG-OD1	6.66	124.29	118.30
1	A	367	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	251	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	387	ASP	CB-CG-OD1	5.28	123.05	118.30
1	A	377	LYS	CD-CE-NZ	5.21	123.67	111.70
1	A	205	SER	CB-CA-C	-5.16	100.30	110.10
1	A	406	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2597	0	2560	44	0
2	C	79	0	77	5	0
3	A	1	0	0	0	0
4	A	16	0	15	0	0
5	A	55	0	0	5	0
5	C	1	0	0	0	0
All	All	2749	0	2652	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:PHE:HD2	2:C:10:GLU:HG2	1.13	1.05
1:A:390:GLN:HG3	5:A:40:HOH:O	1.65	0.95
1:A:309:PHE:CD2	2:C:10:GLU:HG2	2.06	0.88
1:A:404:GLN:HB2	5:A:7:HOH:O	1.72	0.87
1:A:309:PHE:HD2	2:C:10:GLU:CG	1.90	0.84
1:A:206:ARG:HH11	1:A:206:ARG:CG	1.95	0.80
1:A:159:GLY:HA3	5:A:53:HOH:O	1.92	0.69
1:A:283:ASP:HB2	5:A:22:HOH:O	1.96	0.65
1:A:206:ARG:HH11	1:A:206:ARG:HG2	1.62	0.65
1:A:168:LYS:HD2	1:A:173:GLY:O	1.99	0.63
1:A:353:ASP:OD2	1:A:356:LYS:HG2	2.01	0.60
1:A:406:ARG:HH11	1:A:406:ARG:HG2	1.68	0.58
1:A:335:LEU:O	1:A:339:MET:HG2	2.07	0.55
1:A:206:ARG:NH1	1:A:206:ARG:CG	2.64	0.54
1:A:206:ARG:NH1	1:A:206:ARG:HG2	2.23	0.53
1:A:271:VAL:HG22	1:A:328:ARG:HB3	1.89	0.52
1:A:148:ASN:O	1:A:170:LYS:HE3	2.10	0.52
1:A:155:LEU:HB3	1:A:442:PHE:O	2.10	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:LYS:HD3	1:A:175:TYR:CZ	2.45	0.51
1:A:372:LEU:HB3	1:A:377:LYS:HE3	1.92	0.51
1:A:309:PHE:CD2	2:C:10:GLU:CG	2.79	0.51
1:A:207:HIS:ND1	1:A:208:PRO:HD2	2.26	0.50
1:A:231:ASN:H	1:A:231:ASN:ND2	2.10	0.50
1:A:206:ARG:HH11	1:A:206:ARG:HG3	1.72	0.49
1:A:406:ARG:CG	1:A:406:ARG:HH11	2.27	0.47
2:C:2:ARG:NH1	2:C:5:THR:HG23	2.30	0.47
1:A:403:MET:HA	1:A:413:TRP:CZ2	2.51	0.46
1:A:207:HIS:CG	1:A:208:PRO:HD2	2.51	0.46
1:A:204:ASN:O	1:A:268:LYS:HE2	2.15	0.45
1:A:328:ARG:HG2	5:A:14:HOH:O	2.16	0.45
1:A:207:HIS:CE1	1:A:208:PRO:HD2	2.52	0.44
1:A:353:ASP:OD2	1:A:356:LYS:CG	2.64	0.44
1:A:317:ALA:O	1:A:318:PRO:C	2.55	0.43
1:A:474:TYR:CG	1:A:475:SER:N	2.86	0.43
1:A:216:SER:HB3	1:A:472:PHE:CE2	2.54	0.43
1:A:150:PHE:HA	1:A:168:LYS:O	2.19	0.42
1:A:281:MET:SD	1:A:291:THR:HG21	2.60	0.42
1:A:366:ILE:HG21	1:A:366:ILE:HD13	1.83	0.42
1:A:175:TYR:CE1	1:A:433:THR:HG22	2.55	0.42
1:A:218:GLN:HG2	1:A:473:ASP:OD2	2.20	0.41
1:A:414:GLN:HG2	1:A:418:GLU:OE2	2.21	0.41
1:A:176:TYR:OH	1:A:214:LYS:HD3	2.21	0.41
1:A:350:TYR:HD1	1:A:350:TYR:HA	1.74	0.41
1:A:297:LYS:HG2	1:A:306:MET:SD	2.60	0.40
1:A:379:LEU:HD22	1:A:407:PHE:CD2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/342 (91%)	298 (96%)	13 (4%)	1 (0%)	44	64
2	C	8/10 (80%)	8 (100%)	0	0	100	100
All	All	320/352 (91%)	306 (96%)	13 (4%)	1 (0%)	44	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	292	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/299 (93%)	252 (91%)	25 (9%)	11	20
2	C	8/8 (100%)	8 (100%)	0	100	100
All	All	285/307 (93%)	260 (91%)	25 (9%)	12	21

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	144	ARG
1	A	149	GLU
1	A	151	GLU
1	A	160	THR
1	A	168	LYS
1	A	189	LYS
1	A	206	ARG
1	A	216	SER
1	A	231	ASN
1	A	243	ARG
1	A	244	VAL
1	A	267	GLU
1	A	268	LYS
1	A	284	LYS
1	A	285	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	372	LEU
1	A	404	GLN
1	A	406	ARG
1	A	411	ILE
1	A	422	SER
1	A	432	GLU
1	A	447	ILE
1	A	471	GLN
1	A	475	SER
1	A	477	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	204	ASN
1	A	231	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	A	308	1	9,10,11	0.83	0	10,14,16	1.37	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	308	1	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	308	TPO	O3P-P-O2P	2.32	116.99	107.61
1	A	308	TPO	O3P-P-O1P	2.55	120.47	110.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	XFE	A	999	-	16,18,18	1.15	1 (6%)	14,26,26	3.55	10 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	XFE	A	999	-	-	0/4/13/13	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	999	XFE	C8-N9	3.56	1.38	1.32

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	999	XFE	N9-C8-N7	-8.69	121.29	128.86
4	A	999	XFE	C14-C10-C11	-4.34	98.93	106.55
4	A	999	XFE	C15-C14-C13	-3.89	119.49	127.92
4	A	999	XFE	N7-C6-N4	-2.31	111.74	116.85
4	A	999	XFE	C10-C6-N7	-2.08	118.74	122.84
4	A	999	XFE	C3-N4-C6	-2.02	117.70	123.52
4	A	999	XFE	C13-C14-C10	2.58	110.86	107.56
4	A	999	XFE	C8-N9-C11	3.37	121.24	113.33
4	A	999	XFE	C8-N7-C6	3.83	121.22	111.82
4	A	999	XFE	C10-C6-N4	4.14	129.51	121.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/342 (92%)	-0.12	5 (1%) 72 73	11, 30, 53, 65	0
2	C	10/10 (100%)	0.71	1 (10%) 8 7	34, 38, 45, 52	0
All	All	325/352 (92%)	-0.09	6 (1%) 69 70	11, 30, 52, 65	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	477	SER	5.0
1	A	220	HIS	3.6
1	A	160	THR	3.1
2	C	10	GLU	2.6
1	A	446	MET	2.5
1	A	466	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	308	11/12	0.96	0.13	-	33,35,44,46	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	XFE	A	999	16/16	0.97	0.13	-0.52	21,23,25,27	0
3	MN	A	1	1/1	0.97	0.15	-	63,63,63,63	0

## 6.5 Other polymers

There are no such residues in this entry.