



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 08:19 AM EDT

PDB ID : 3MZV  
Title : Crystal structure of a decaprenyl diphosphate synthase from *Rhodobacter capsulatus*  
Authors : Quartararo, C.E.; Patskovsky, Y.; Bonanno, J.B.; Rutter, M.; Bain, K.T.; Chang, S.; Toro, R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : unknown  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	<div> <div>11%</div> <div> <div></div> <div>76%</div> <div>12%</div> <div>9%</div> </div> </div>
1	B	341	<div> <div>11%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>5%</div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Decaprenyl diphosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	1	0
			2337	1475	418	434	10			
1	B	325	Total	C	N	O	S	0	2	0
			2475	1552	437	475	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	334	GLU	-	EXPRESSION TAG	UNP Q07D00
A	335	GLY	-	EXPRESSION TAG	UNP Q07D00
A	336	HIS	-	EXPRESSION TAG	UNP Q07D00
A	337	HIS	-	EXPRESSION TAG	UNP Q07D00
A	338	HIS	-	EXPRESSION TAG	UNP Q07D00
A	339	HIS	-	EXPRESSION TAG	UNP Q07D00
A	340	HIS	-	EXPRESSION TAG	UNP Q07D00
A	341	HIS	-	EXPRESSION TAG	UNP Q07D00
B	334	GLU	-	EXPRESSION TAG	UNP Q07D00
B	335	GLY	-	EXPRESSION TAG	UNP Q07D00
B	336	HIS	-	EXPRESSION TAG	UNP Q07D00
B	337	HIS	-	EXPRESSION TAG	UNP Q07D00
B	338	HIS	-	EXPRESSION TAG	UNP Q07D00
B	339	HIS	-	EXPRESSION TAG	UNP Q07D00
B	340	HIS	-	EXPRESSION TAG	UNP Q07D00
B	341	HIS	-	EXPRESSION TAG	UNP Q07D00


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	126	Total	O	0	0
			126	126		
2	B	179	Total	O	0	0
			179	179		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Decaprenyl diphosphate synthase



#### • Molecule 1: Decaprenyl diphosphate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.46Å 90.25Å 80.17Å 90.00° 125.76° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.94 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-1.90) 99.4 (19.94-1.89)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.89 (at 1.89Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.193 , 0.221 0.207 , 0.237	Depositor DCC
$R_{free}$ test set	2839 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 53.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5117	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	1/2378 (0.0%)	0.78	2/3228 (0.1%)
1	B	0.81	1/2521 (0.0%)	0.82	5/3420 (0.1%)
All	All	0.79	2/4899 (0.0%)	0.80	7/6648 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	287	LYS	CE-NZ	5.92	1.63	1.49
1	B	260	GLU	CG-CD	5.24	1.59	1.51

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	127	ARG	NE-CZ-NH2	11.47	126.03	120.30
1	A	132	MET	CG-SD-CE	9.97	116.16	100.20
1	B	33	MET	CG-SD-CE	8.29	113.47	100.20
1	A	132	MET	CA-CB-CG	7.98	126.87	113.30
1	B	127	ARG	NE-CZ-NH1	-7.61	116.49	120.30
1	B	93	ASP	CB-CG-OD1	6.72	124.35	118.30
1	B	59	MET	CG-SD-CE	-5.83	90.87	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2337	0	2320	35	0
1	B	2475	0	2444	38	0
2	A	126	0	0	1	0
2	B	179	0	0	3	0
All	All	5117	0	4764	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:33:MET:HE2	1:B:45:THR:HG21	1.36	1.06
1:B:129:PHE:HA	1:B:132[B]:MET:HG3	1.38	1.06
1:B:33:MET:HE2	1:B:45:THR:CG2	1.89	1.01
1:B:132[B]:MET:HE1	1:B:144:LEU:HD12	1.43	0.97
1:B:33:MET:CE	1:B:45:THR:CG2	2.49	0.90
1:B:132[B]:MET:CE	1:B:144:LEU:HD12	2.04	0.88
1:B:26:ASN:OD1	1:B:55:ARG:NH2	2.09	0.86
1:B:33:MET:CE	1:B:45:THR:HG22	2.07	0.85
1:A:278:LEU:O	1:A:278:LEU:HD12	1.78	0.84
1:A:177:ARG:HA	1:A:181:ALA:HB3	1.63	0.80
1:A:249:VAL:HG22	1:A:285:MET:HE3	1.69	0.74
1:A:249:VAL:HG22	1:A:285:MET:CE	2.18	0.72
1:B:129:PHE:CE2	1:B:132[B]:MET:SD	2.82	0.72
1:B:234:LYS:HG3	1:B:235:ASN:ND2	2.05	0.72
1:B:33:MET:HE1	1:B:45:THR:HG22	1.75	0.69
1:B:129:PHE:CD2	1:B:132[B]:MET:SD	2.86	0.68
1:B:78:LEU:CD2	1:B:132[A]:MET:CE	2.72	0.68
1:B:11:HIS:NE2	1:B:54:LYS:O	2.15	0.68
1:B:78:LEU:HD23	1:B:132[A]:MET:SD	2.34	0.68
1:A:309:LEU:HD23	1:A:312:LEU:HD12	1.80	0.64
1:A:299:ILE:HD13	1:A:329:VAL:HG11	1.79	0.63
1:B:78:LEU:CD2	1:B:132[A]:MET:HE1	2.28	0.63
1:A:278:LEU:C	1:A:278:LEU:HD12	2.19	0.63
1:A:48:LEU:HD23	1:A:87:THR:HG23	1.81	0.62
1:B:67:LEU:CD1	1:B:315:HIS:HB3	2.30	0.61
1:A:41:ILE:HG21	1:A:123:TYR:CD2	2.36	0.60
1:B:48:LEU:HD23	1:B:87:THR:HG23	1.82	0.60
1:A:245:LEU:HD21	1:A:268:ILE:HG21	1.84	0.59
1:A:249:VAL:CG2	1:A:285:MET:CE	2.82	0.58
1:B:78:LEU:HD21	1:B:132[A]:MET:CE	2.36	0.56

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:LEU:HD23	1:B:132[A]:MET:CE	2.37	0.54
1:B:41:ILE:HG21	1:B:123:TYR:CD2	2.43	0.54
1:B:33:MET:HE1	1:B:45:THR:CG2	2.34	0.53
1:A:326:ASP:OD1	1:A:333:ARG:NH1	2.41	0.53
1:B:261:ARG:HD3	2:B:562:HOH:O	2.08	0.53
1:A:245:LEU:HD21	1:A:268:ILE:CG2	2.39	0.52
1:A:249:VAL:CG2	1:A:285:MET:HE1	2.40	0.52
1:A:260:GLU:OE2	1:A:287:LYS:NZ	2.37	0.50
1:A:245:LEU:HD23	2:A:748:HOH:O	2.12	0.50
1:A:28:LEU:HD12	1:A:76:VAL:HG12	1.94	0.50
1:A:109:ASN:HB3	1:A:117:SER:OG	2.11	0.50
1:A:132:MET:HE3	1:A:141:MET:HB3	1.94	0.50
1:A:95:VAL:HG21	1:B:118:VAL:HG21	1.94	0.49
1:A:40:ARG:O	1:A:44:VAL:HG23	2.13	0.49
1:B:177:ARG:HA	1:B:181:ALA:HB3	1.94	0.48
1:A:281:ALA:O	1:A:285:MET:HG3	2.14	0.47
1:B:40:ARG:HD2	1:B:112:TRP:CD1	2.49	0.47
1:A:168:THR:OG1	1:A:169:GLU:N	2.47	0.47
1:A:115:LYS:HA	1:B:96:VAL:HG22	1.98	0.46
1:B:289:GLY:HA2	2:B:392:HOH:O	2.14	0.46
1:A:85:ILE:O	1:A:89:THR:HG23	2.16	0.45
1:B:132[B]:MET:HE3	1:B:144:LEU:HD12	1.95	0.45
1:A:315:HIS:CE1	1:A:317:LEU:HD12	2.52	0.45
1:B:132[A]:MET:HE2	1:B:132[A]:MET:HB3	1.74	0.45
1:A:67:LEU:CD1	1:A:315:HIS:HB3	2.47	0.44
1:A:9:LYS:HB2	1:A:12:ASP:OD2	2.18	0.44
1:A:109:ASN:O	1:A:113:ASP:HA	2.17	0.44
1:B:174:ARG:NH1	2:B:704:HOH:O	2.50	0.44
1:B:45:THR:HG22	1:B:49:ILE:HD12	2.00	0.44
1:B:38:ALA:HB3	1:B:41:ILE:HD12	2.00	0.43
1:A:171:ILE:HD13	1:A:171:ILE:N	2.33	0.43
1:A:112:TRP:HB2	1:A:116:SER:OG	2.18	0.43
1:A:299:ILE:HD13	1:A:329:VAL:CG1	2.45	0.43
1:B:78:LEU:HD21	1:B:132[A]:MET:HE3	2.00	0.42
1:B:228:LYS:HG3	1:B:292:GLU:OE1	2.20	0.42
1:B:315:HIS:ND1	1:B:316:PRO:HD2	2.35	0.41
1:A:247:MET:N	1:A:248:PRO:HD2	2.35	0.41
1:B:67:LEU:HD13	1:B:315:HIS:HB3	2.00	0.41
1:A:247:MET:SD	1:A:251:LYS:HE2	2.61	0.41
1:A:249:VAL:HA	1:A:285:MET:HE2	2.04	0.40
1:B:78:LEU:HD21	1:B:132[A]:MET:HE1	2.02	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/341 (89%)	291 (96%)	13 (4%)	0	100	100
1	B	325/341 (95%)	321 (99%)	4 (1%)	0	100	100
All	All	629/682 (92%)	612 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/268 (86%)	223 (96%)	8 (4%)	41	30
1	B	249/268 (93%)	243 (98%)	6 (2%)	54	47
All	All	480/536 (90%)	466 (97%)	14 (3%)	48	39

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	HIS
1	A	55	ARG
1	A	114	ASN
1	A	168	THR
1	A	171	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	247	MET
1	A	278	LEU
1	A	333	ARG
1	B	13	ARG
1	B	36	GLU
1	B	101	GLN
1	B	111	LEU
1	B	224	ASP
1	B	306	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	HIS
1	A	114	ASN
1	B	101	GLN
1	B	217	GLN
1	B	235	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/341 (90%)	0.67	36 (11%) 5 5	27, 47, 80, 88	0
1	B	325/341 (95%)	0.53	38 (11%) 5 5	29, 40, 71, 83	0
All	All	634/682 (92%)	0.60	74 (11%) 5 5	27, 42, 76, 88	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	GLY	9.2
1	A	104	GLY	8.8
1	A	103	ARG	8.1
1	B	232	ILE	7.2
1	A	111	LEU	7.1
1	A	236	THR	6.2
1	B	51	ALA	5.5
1	A	112	TRP	5.4
1	A	51	ALA	5.3
1	B	52	GLY	5.3
1	B	105	ARG	5.2
1	B	53	GLY	5.1
1	B	235	ASN	4.8
1	B	234	LYS	4.7
1	B	103	ARG	4.6
1	B	97	ASP	4.5
1	B	106	PRO	4.4
1	B	233	GLY	4.4
1	A	105	ARG	4.2
1	B	100	ARG	4.1
1	A	47	HIS	3.9
1	A	52	GLY	3.7
1	A	53	GLY	3.7
1	A	106	PRO	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	50	GLU	3.5
1	B	98	GLU	3.4
1	A	259	ALA	3.4
1	A	110	LEU	3.4
1	B	12	ASP	3.2
1	A	102	ARG	3.1
1	B	281	ALA	3.1
1	A	333	ARG	3.1
1	A	37	HIS	3.1
1	A	50	GLU	3.1
1	B	101	GLN	3.1
1	B	96	VAL	3.1
1	A	281	ALA	3.1
1	A	275	ASP	2.9
1	B	37	HIS	2.9
1	B	54	LYS	2.9
1	A	223	LEU	2.9
1	A	165	LEU	2.8
1	B	47	HIS	2.8
1	A	278	LEU	2.7
1	B	231	GLU	2.7
1	B	11	HIS	2.7
1	A	331	ARG	2.7
1	B	229	SER	2.6
1	B	48	LEU	2.6
1	A	264	TRP	2.6
1	B	107	THR	2.6
1	B	118	VAL	2.6
1	B	117	SER	2.5
1	A	160	THR	2.5
1	B	99	SER	2.5
1	B	190	VAL	2.4
1	B	321	LEU	2.4
1	A	109	ASN	2.4
1	A	243	ARG	2.4
1	A	272	ASP	2.4
1	B	102	ARG	2.4
1	A	256	ALA	2.3
1	B	36	GLU	2.3
1	B	49	ILE	2.3
1	A	285	MET	2.3
1	A	49	ILE	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	268	ILE	2.3
1	B	112	TRP	2.2
1	B	333	ARG	2.2
1	A	222	LEU	2.2
1	A	267	VAL	2.2
1	A	284	LEU	2.1
1	B	113	ASP	2.0
1	A	249	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.