



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2017 – 12:55 AM EDT

PDB ID : 3N0U  
Title : Crystal structure of Tm1821, the 8-oxoguanine DNA glycosylase of *Thermotoga maritima*  
Authors : Cooper, D.R.; Roy, A.; Derewenda, Z.S.; Integrated Center for Structure and Function Innovation (ISFI)  
Deposited on : unknown  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2279 (1.50-1.50)
Clashscore	112137	2503 (1.50-1.50)
Ramachandran outliers	110173	2445 (1.50-1.50)
Sidechain outliers	110143	2443 (1.50-1.50)
RSRZ outliers	101464	2305 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	<div> <div>4%</div> <div>89%</div> <div>6%</div> <div>5%</div> </div>
1	B	219	<div> <div>4%</div> <div>85%</div> <div>10%</div> <div>5%</div> </div>
1	C	219	<div> <div>5%</div> <div>87%</div> <div>7%</div> <div>5%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11303 atoms, of which 5395 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable N-glycosylase/DNA lyase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	209	Total	C	H	N	O	S	0	5	0
			3581	1132	1823	313	308	5			
1	B	209	Total	C	H	N	O	S	0	4	0
			3537	1121	1794	308	309	5			
1	C	208	Total	C	H	N	O	S	0	2	0
			3505	1111	1778	306	305	5			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	expression tag	UNP Q9X2E1
A	-10	GLY	-	expression tag	UNP Q9X2E1
A	-9	SER	-	expression tag	UNP Q9X2E1
A	-8	ASP	-	expression tag	UNP Q9X2E1
A	-7	LYS	-	expression tag	UNP Q9X2E1
A	-6	ILE	-	expression tag	UNP Q9X2E1
A	-5	HIS	-	expression tag	UNP Q9X2E1
A	-4	HIS	-	expression tag	UNP Q9X2E1
A	-3	HIS	-	expression tag	UNP Q9X2E1
A	-2	HIS	-	expression tag	UNP Q9X2E1
A	-1	HIS	-	expression tag	UNP Q9X2E1
A	0	HIS	-	expression tag	UNP Q9X2E1
B	-11	MET	-	expression tag	UNP Q9X2E1
B	-10	GLY	-	expression tag	UNP Q9X2E1
B	-9	SER	-	expression tag	UNP Q9X2E1
B	-8	ASP	-	expression tag	UNP Q9X2E1
B	-7	LYS	-	expression tag	UNP Q9X2E1
B	-6	ILE	-	expression tag	UNP Q9X2E1
B	-5	HIS	-	expression tag	UNP Q9X2E1
B	-4	HIS	-	expression tag	UNP Q9X2E1
B	-3	HIS	-	expression tag	UNP Q9X2E1
B	-2	HIS	-	expression tag	UNP Q9X2E1
B	-1	HIS	-	expression tag	UNP Q9X2E1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	HIS	-	expression tag	UNP Q9X2E1
C	-11	MET	-	expression tag	UNP Q9X2E1
C	-10	GLY	-	expression tag	UNP Q9X2E1
C	-9	SER	-	expression tag	UNP Q9X2E1
C	-8	ASP	-	expression tag	UNP Q9X2E1
C	-7	LYS	-	expression tag	UNP Q9X2E1
C	-6	ILE	-	expression tag	UNP Q9X2E1
C	-5	HIS	-	expression tag	UNP Q9X2E1
C	-4	HIS	-	expression tag	UNP Q9X2E1
C	-3	HIS	-	expression tag	UNP Q9X2E1
C	-2	HIS	-	expression tag	UNP Q9X2E1
C	-1	HIS	-	expression tag	UNP Q9X2E1
C	0	HIS	-	expression tag	UNP Q9X2E1

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Na 1 1	0	0
2	A	1	Total Na 1 1	0	0
2	C	1	Total Na 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	232	Total O 232 232	0	0
3	B	221	Total O 221 221	0	0
3	C	224	Total O 224 224	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Probable N-glycosylase/DNA lyase



- Molecule 1: Probable N-glycosylase/DNA lyase



- Molecule 1: Probable N-glycosylase/DNA lyase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.87Å 93.64Å 135.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.85 – 1.50 25.85 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (25.85-1.50) 99.0 (25.85-1.50)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.6.1 _357	Depositor
R, $R_{free}$	0.150 , 0.190 0.145 , 0.184	Depositor DCC
$R_{free}$ test set	5585 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.3	Xtriage
Anisotropy	0.126	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.48 , 53.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11303	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 57.07 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.4955e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/1810	0.55	0/2419
1	B	0.47	0/1789	0.57	0/2392
1	C	0.44	0/1767	0.57	1/2362 (0.0%)
All	All	0.45	0/5366	0.56	1/7173 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	12	ARG	NE-CZ-NH1	5.23	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1758	1823	1816	9	0
1	B	1743	1794	1787	20	0
1	C	1727	1778	1773	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	232	0	0	3	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	221	0	0	4	0
3	C	224	0	0	4	3
All	All	5908	5395	5376	42	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90[B]:ARG:NH1	3:A:664:HOH:O	2.13	0.78
1:B:58:ARG:HA	1:B:61:LYS:CE	2.16	0.76
1:B:58:ARG:HA	1:B:61:LYS:HE3	1.73	0.71
1:B:6:LYS:HE2	1:B:10:ARG:HH12	1.58	0.69
1:B:36[B]:GLU:OE2	3:B:497:HOH:O	2.14	0.65
1:A:90[B]:ARG:NH2	3:A:664:HOH:O	2.33	0.62
1:B:58:ARG:HA	1:B:61:LYS:HE2	1.83	0.59
1:C:28:ARG:NH2	3:C:640:HOH:O	2.37	0.58
1:A:201:VAL:HG12	1:A:202:LYS:HG3	1.85	0.57
1:A:164:PRO:HB3	1:A:171[A]:ARG:CZ	2.35	0.56
1:A:90[B]:ARG:CZ	3:A:664:HOH:O	2.50	0.56
1:B:63:ILE:HG12	1:B:78:LYS:HG2	1.87	0.56
1:C:2:GLU:OE2	1:C:6:LYS:HD2	2.06	0.55
1:A:27:LYS:O	1:A:31[A]:GLU:HG3	2.07	0.55
1:B:6:LYS:HG2	1:B:10:ARG:NH1	2.22	0.54
1:B:90[B]:ARG:NH1	3:B:668:HOH:O	2.28	0.54
1:C:129:LYS:HD2	1:C:147:ASP:HB3	1.91	0.53
1:C:54:GLU:HG3	1:C:55:GLY:N	2.25	0.52
1:C:28:ARG:NH1	3:C:366:HOH:O	2.42	0.51
1:B:201:VAL:HG12	1:B:202:LYS:HG3	1.93	0.51
1:B:6:LYS:CE	1:B:10:ARG:HH12	2.25	0.49
1:C:12:ARG:HD3	3:C:254:HOH:O	2.13	0.49
1:B:129:LYS:HD2	1:B:147:ASP:HB3	1.93	0.49
1:B:57:ILE:O	1:B:61:LYS:HG3	2.13	0.48
1:B:36[B]:GLU:OE1	3:B:494:HOH:O	2.20	0.48
1:B:153:LEU:HD23	1:B:153:LEU:C	2.36	0.46
1:B:27:LYS:O	1:B:31:GLU:HG3	2.16	0.46
1:A:129:LYS:HD2	1:A:147:ASP:HB3	1.98	0.46
1:A:164:PRO:HG2	1:A:167:TRP:CH2	2.52	0.44
1:B:57:ILE:HG22	1:B:61:LYS:HE2	2.01	0.43
1:C:201:VAL:HG12	1:C:202:LYS:HG3	2.00	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36[B]:GLU:OE2	3:B:490:HOH:O	2.22	0.42
1:B:92:GLU:O	1:B:96:GLU:HG3	2.19	0.42
1:C:159:LEU:HD12	1:C:159:LEU:N	2.33	0.42
1:C:24:GLU:HG3	3:C:523:HOH:O	2.20	0.42
1:C:23:PHE:CZ	1:C:141:GLU:HG2	2.55	0.41
1:C:129:LYS:HE2	1:C:130:GLU:OE2	2.20	0.41
1:B:89:LYS:HE3	1:B:93:PHE:CE1	2.55	0.41
1:C:201:VAL:HG12	1:C:202:LYS:CG	2.51	0.41
1:B:82:VAL:HG22	1:B:82:VAL:O	2.22	0.40
1:C:84:HIS:CE1	1:C:90[B]:ARG:HG3	2.57	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:663:HOH:O	3:C:675:HOH:O[4_455]	1.87	0.33
3:A:264:HOH:O	3:C:420:HOH:O[1_655]	2.16	0.04
3:C:605:HOH:O	3:C:628:HOH:O[4_455]	2.19	0.01
3:A:483:HOH:O	3:A:511:HOH:O[3_655]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/219 (97%)	211 (99%)	2 (1%)	0	100	100
1	B	211/219 (96%)	208 (99%)	3 (1%)	0	100	100
1	C	208/219 (95%)	207 (100%)	1 (0%)	0	100	100
All	All	632/657 (96%)	626 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/190 (97%)	182 (98%)	3 (2%)	68	39
1	B	183/190 (96%)	182 (100%)	1 (0%)	91	80
1	C	181/190 (95%)	179 (99%)	2 (1%)	78	56
All	All	549/570 (96%)	543 (99%)	6 (1%)	78	56

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	PHE
1	A	89	LYS
1	A	170	LYS
1	B	156	ARG
1	C	23	PHE
1	C	171	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	209/219 (95%)	0.15	9 (4%) 36 41	16, 24, 48, 81	0
1	B	209/219 (95%)	0.10	9 (4%) 36 41	15, 25, 48, 71	0
1	C	208/219 (94%)	0.16	12 (5%) 24 27	16, 27, 54, 76	0
All	All	626/657 (95%)	0.14	30 (4%) 31 35	15, 25, 51, 81	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-1	HIS	7.3
1	B	-1	HIS	6.1
1	B	156	ARG	3.8
1	A	156	ARG	3.3
1	C	134	PHE	3.2
1	C	145	ILE	3.0
1	A	161	GLN	2.9
1	B	145	ILE	2.9
1	A	193	PHE	2.8
1	C	156	ARG	2.7
1	B	134	PHE	2.7
1	A	145	ILE	2.6
1	B	193	PHE	2.5
1	C	193	PHE	2.5
1	C	0	HIS	2.5
1	C	152	ARG	2.4
1	B	163	ILE	2.4
1	C	109	LYS	2.3
1	A	162	GLU	2.3
1	A	152	ARG	2.3
1	A	170	LYS	2.3
1	A	46	VAL	2.3
1	B	81	GLU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	2	GLU	2.1
1	B	162	GLU	2.1
1	B	131	ALA	2.1
1	C	106	ASN	2.1
1	C	131	ALA	2.1
1	C	28	ARG	2.1
1	C	46	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	A	208	1/1	0.99	0.06	-1.24	22,22,22,22	0
2	NA	B	208	1/1	0.99	0.05	-1.78	20,20,20,20	0
2	NA	C	208	1/1	0.99	0.06	-1.85	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.