



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 24, 2018 – 03:05 AM EST

PDB ID : 1N46
Title : CRYSTAL STRUCTURE OF HUMAN TR BETA LIGAND-BINDING DOMAIN COMPLEXED WITH A POTENT SUBTYPE-SELECTIVE THYROMIMETIC
Authors : Dow, R.L.; Schneider, S.R.; Paight, E.S.; Hank, R.F.; Chiang, P.; Cornelius, P.; Lee, E.; Newsome, W.P.; Swick, A.G.; Spitzer, J.; Hargrove, D.M.; Patterson, T.A.; Pandit, J.; Chrunk, B.A.; LeMotte, P.K.; Danley, D.E.; Rosner, M.H.; Ammirati, M.J.; Simons, S.P.; Schulte, G.K.; Tate, B.F.; DaSilva-Jardine, P.
Deposited on : 2002-10-30
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>2%</div> <div>67%</div> <div>24%</div> <div>• • •</div> </div>
1	B	258	<div> <div>5%</div> <div>63%</div> <div>23%</div> <div>6%</div> <div>7%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thyroid hormone receptor Beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1968	1264	326	362	16			
1	B	239	Total	C	N	O	S	0	0	0
			1904	1223	315	350	16			

- Molecule 2 is [4-(4-HYDROXY-3-ISOPROPYL-PHENOXY)-3,5-DIMETHYL-PHENYL]-6-AZAURACIL (three-letter code: PFA) (formula: C₂₀H₂₁N₃O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	20	3	4		
2	B	1	Total	C	N	O	0	0
			27	20	3	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	85	Total 85	O 85	0	0
3	B	29	Total 29	O 29	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thyroid hormone receptor Beta-1



• Molecule 1: Thyroid hormone receptor Beta-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.92Å 105.31Å 55.99Å 90.00° 98.19° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.66 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.7 (30.00-2.20) 96.7 (29.66-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.192 , 0.256 0.197 , 0.259	Depositor DCC
R_{free} test set	1226 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	19.2	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4040	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PFA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	2/2012 (0.1%)	1.24	20/2722 (0.7%)
1	B	1.03	2/1945 (0.1%)	1.18	18/2628 (0.7%)
All	All	1.05	4/3957 (0.1%)	1.21	38/5350 (0.7%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	313	MET	SD-CE	-6.75	1.40	1.77
1	A	313	MET	SD-CE	-6.74	1.40	1.77
1	A	217	GLU	CG-CD	5.22	1.59	1.51
1	B	430	MET	CG-SD	-5.09	1.68	1.81

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	383	ARG	NE-CZ-NH2	-12.86	113.87	120.30
1	A	338	ARG	NE-CZ-NH2	-12.20	114.20	120.30
1	A	320	ARG	NE-CZ-NH2	-9.91	115.35	120.30
1	A	282	ARG	NE-CZ-NH1	9.88	125.24	120.30
1	A	282	ARG	NE-CZ-NH2	-9.02	115.79	120.30
1	B	429	ARG	NE-CZ-NH1	8.15	124.38	120.30
1	B	383	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	A	351	ASP	CB-CG-OD2	7.38	124.94	118.30
1	A	320	ARG	NE-CZ-NH1	7.29	123.94	120.30
1	B	410	ARG	NE-CZ-NH2	-7.28	116.66	120.30
1	A	355	ASP	CB-CG-OD2	7.07	124.67	118.30
1	A	304	LEU	CB-CG-CD1	6.92	122.76	111.00
1	A	265	ASP	CB-CG-OD2	6.53	124.17	118.30
1	A	338	ARG	CA-CB-CG	6.50	127.71	113.40
1	B	429	ARG	NE-CZ-NH2	-6.33	117.14	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	429	ARG	NE-CZ-NH1	6.27	123.44	120.30
1	A	397	ASP	CB-CG-OD2	6.22	123.90	118.30
1	B	423	MET	CG-SD-CE	-6.20	90.29	100.20
1	A	300	ASP	CB-CG-OD1	6.16	123.85	118.30
1	A	461	ASP	CB-CG-OD2	6.06	123.76	118.30
1	B	305	LEU	CA-CB-CG	-6.03	101.43	115.30
1	B	397	ASP	CB-CG-OD2	5.98	123.68	118.30
1	B	430	MET	CA-CB-CG	-5.94	103.20	113.30
1	B	355	ASP	CB-CG-OD2	5.93	123.63	118.30
1	A	322	ASP	CB-CG-OD2	5.77	123.50	118.30
1	A	427	ASP	CB-CG-OD1	5.74	123.46	118.30
1	B	300	ASP	CB-CG-OD2	5.64	123.38	118.30
1	B	366	ASP	CB-CG-OD1	5.64	123.37	118.30
1	A	430	MET	CB-CA-C	-5.61	99.18	110.40
1	B	430	MET	CG-SD-CE	-5.57	91.28	100.20
1	B	351	ASP	CB-CG-OD2	5.52	123.27	118.30
1	A	367	ASP	CB-CG-OD2	5.51	123.26	118.30
1	A	249	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	391	ARG	NE-CZ-NH1	-5.40	117.60	120.30
1	B	320	ARG	NE-CZ-NH2	-5.37	117.61	120.30
1	B	338	ARG	NE-CZ-NH2	-5.25	117.67	120.30
1	B	249	ASP	CB-CG-OD2	5.21	122.98	118.30
1	B	438	ARG	NE-CZ-NH1	5.07	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1968	0	1972	39	0
1	B	1904	0	1902	44	0
2	A	27	0	21	4	0
2	B	27	0	21	5	0
3	A	85	0	0	3	0
3	B	29	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4040	0	3916	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:310:MET:HE3	2:B:463:PFA:H29C	1.54	0.89
1:A:310:MET:HE3	2:A:462:PFA:H29C	1.58	0.85
1:B:429:ARG:HB3	3:B:13:HOH:O	1.78	0.82
1:A:310:MET:HA	1:A:313:MET:HE3	1.63	0.79
1:B:276:ILE:HD12	2:B:463:PFA:HC72	1.64	0.78
1:B:291:PRO:O	1:B:295:GLU:HG2	1.86	0.76
1:A:390:GLU:HG2	3:A:90:HOH:O	1.87	0.75
1:A:310:MET:CE	2:A:462:PFA:H29C	2.19	0.71
1:B:320:ARG:HH22	2:B:463:PFA:HN6	1.39	0.71
1:B:310:MET:CE	1:B:313:MET:HE3	2.24	0.67
1:B:310:MET:HE2	1:B:313:MET:HE3	1.79	0.65
1:A:320:ARG:HH22	2:A:462:PFA:HN6	1.44	0.65
1:A:456:LEU:O	1:A:460:GLU:HB2	1.96	0.65
1:B:253:ALA:HB2	1:B:343:ASN:HD21	1.65	0.60
1:B:456:LEU:O	1:B:460:GLU:HB3	2.00	0.60
1:A:281:THR:HG22	1:A:454:LEU:CD2	2.32	0.59
1:A:430:MET:CE	1:B:223:LYS:HA	2.33	0.59
1:A:277:THR:O	1:A:281:THR:HG23	2.04	0.58
1:A:433:ALA:HB1	1:B:409:TYR:OH	2.05	0.57
1:B:310:MET:CE	2:B:463:PFA:H29C	2.33	0.56
1:B:304:LEU:HD13	1:B:386:LEU:HD21	1.88	0.56
1:A:429:ARG:HB3	3:A:45:HOH:O	2.06	0.56
1:A:430:MET:HE3	1:B:223:LYS:HA	1.88	0.55
1:A:248:GLU:OE1	1:A:248:GLU:HA	2.07	0.54
1:A:248:GLU:OE1	1:A:252:GLN:OE1	2.24	0.54
1:B:264:VAL:HG11	1:B:446:CYS:SG	2.48	0.54
1:B:325:SER:HB2	1:B:327:THR:HG23	1.88	0.54
1:A:260:GLU:HG2	1:A:343:ASN:HD21	1.74	0.53
1:B:213:GLU:CD	1:B:409:TYR:HB2	2.29	0.53
1:B:310:MET:HA	1:B:313:MET:HE3	1.91	0.52
1:B:414:VAL:HG22	1:B:417:PHE:HD1	1.75	0.52
1:A:418:TRP:HB3	1:A:419:PRO:CD	2.40	0.52
1:B:213:GLU:OE2	1:B:409:TYR:HB2	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:313:MET:O	2:B:463:PFA:HC3	2.10	0.50
1:B:407:ILE:CG1	1:B:417:PHE:HE2	2.24	0.50
1:B:421:LEU:O	1:B:424:LYS:HB2	2.11	0.50
1:A:281:THR:HG22	1:A:454:LEU:HD21	1.92	0.49
1:A:426:THR:HG23	3:A:95:HOH:O	2.12	0.49
1:A:272:PHE:HE1	1:A:336:VAL:HG11	1.78	0.49
1:B:219:TRP:O	1:B:223:LYS:HD3	2.12	0.48
1:B:310:MET:HE1	1:B:313:MET:HE3	1.94	0.48
1:A:414:VAL:HG22	1:A:417:PHE:HD1	1.78	0.48
1:B:280:ILE:O	1:B:284:VAL:HG23	2.14	0.48
1:A:234:ALA:C	1:A:235:GLN:HG2	2.34	0.47
1:B:379:MET:O	1:B:396:GLN:HB2	2.13	0.47
1:A:423:MET:CE	1:B:227:GLU:HG2	2.45	0.47
1:A:447:PRO:HG2	1:A:450:LEU:HD11	1.96	0.47
1:A:356:LEU:HG	1:A:360:LEU:HD22	1.97	0.47
1:B:277:THR:HB	1:B:278:PRO:HD3	1.97	0.46
1:A:272:PHE:HB3	2:A:462:PFA:HC93	1.98	0.46
1:B:407:ILE:HG12	1:B:417:PHE:HE2	1.81	0.46
1:A:270:SER:HB3	1:A:450:LEU:HD22	1.98	0.46
1:B:429:ARG:HD2	3:B:8:HOH:O	2.15	0.46
1:B:365:LEU:HA	1:B:369:GLU:OE1	2.15	0.45
1:B:292:MET:HG2	1:B:395:TYR:CE1	2.52	0.45
1:B:239:TRP:O	1:B:240:LYS:C	2.55	0.45
1:A:422:LEU:HA	1:A:422:LEU:HD23	1.78	0.44
1:A:279:ALA:HB2	1:A:331:ASN:HD21	1.83	0.44
1:A:304:LEU:HG	1:A:378:LEU:O	2.18	0.44
1:A:377:LEU:HD23	1:A:425:VAL:HG13	1.99	0.44
1:B:226:THR:O	1:B:230:VAL:HG22	2.18	0.44
1:B:304:LEU:HG	1:B:378:LEU:O	2.18	0.44
1:B:409:TYR:CG	1:B:409:TYR:O	2.70	0.43
1:A:309:CYS:SG	1:A:313:MET:HE2	2.58	0.43
1:B:264:VAL:HG12	1:B:264:VAL:O	2.18	0.43
1:B:304:LEU:HD12	1:B:304:LEU:HA	1.81	0.43
1:A:388:CYS:O	1:A:392:ILE:HD12	2.19	0.42
1:B:313:MET:HE1	1:B:459:PHE:HZ	1.83	0.42
1:A:416:HIS:HB3	1:A:419:PRO:HG2	2.01	0.42
1:B:310:MET:HE2	1:B:310:MET:HB2	1.87	0.42
1:A:277:THR:N	1:A:278:PRO:HD2	2.34	0.42
1:A:312:ILE:HG23	1:A:374:GLN:HG2	2.01	0.42
1:A:278:PRO:O	1:A:282:ARG:HG3	2.20	0.41
1:A:322:ASP:HA	1:A:323:PRO:HD3	1.84	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418:TRP:HB3	1:A:419:PRO:HD3	2.02	0.41
1:A:313:MET:HE1	1:A:459:PHE:HZ	1.85	0.40
1:B:409:TYR:C	1:B:409:TYR:CD2	2.94	0.40
1:B:330:LEU:O	1:B:331:ASN:HB2	2.21	0.40
1:B:422:LEU:O	1:B:425:VAL:HB	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/258 (95%)	236 (97%)	7 (3%)	1 (0%)	38	41
1	B	233/258 (90%)	222 (95%)	7 (3%)	4 (2%)	11	7
All	All	477/516 (92%)	458 (96%)	14 (3%)	5 (1%)	18	16

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	234	ALA
1	A	234	ALA
1	B	411	LYS
1	B	240	LYS
1	B	264	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/226 (96%)	196 (90%)	22 (10%)	9	8
1	B	211/226 (93%)	186 (88%)	25 (12%)	6	5
All	All	429/452 (95%)	382 (89%)	47 (11%)	7	6

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	220	GLU
1	A	221	LEU
1	A	235	GLN
1	A	240	LYS
1	A	242	LYS
1	A	243	ARG
1	A	248	GLU
1	A	274	LYS
1	A	276	ILE
1	A	281	THR
1	A	304	LEU
1	A	324	GLU
1	A	333	GLU
1	A	360	LEU
1	A	394	LYS
1	A	403	PHE
1	A	409	TYR
1	A	411	LYS
1	A	430	MET
1	A	437	SER
1	A	448	THR
1	A	450	LEU
1	B	220	GLU
1	B	223	LYS
1	B	241	GLN
1	B	243	ARG
1	B	252	GLN
1	B	274	LYS
1	B	281	THR
1	B	288	LYS
1	B	295	GLU
1	B	299	GLU
1	B	304	LEU
1	B	305	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	324	GLU
1	B	338	ARG
1	B	359	SER
1	B	360	LEU
1	B	361	SER
1	B	407	ILE
1	B	409	TYR
1	B	422	LEU
1	B	424	LYS
1	B	426	THR
1	B	430	MET
1	B	460	GLU
1	B	461	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	252	GLN
1	A	331	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PFA	A	462	-	26,29,29	2.10	7 (26%)	33,42,42	3.06	12 (36%)
2	PFA	B	463	-	26,29,29	2.23	10 (38%)	33,42,42	2.45	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PFA	A	462	-	-	0/12/12/12	0/3/3/3
2	PFA	B	463	-	-	0/12/12/12	0/3/3/3

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	463	PFA	C17-N3	-3.03	1.25	1.32
2	B	463	PFA	C4-N5	-2.41	1.39	1.44
2	A	462	PFA	C7-C6	-2.28	1.46	1.51
2	A	462	PFA	C5-C6	2.01	1.42	1.39
2	A	462	PFA	C10-N6	2.17	1.37	1.33
2	B	463	PFA	C24-C25	2.18	1.55	1.52
2	B	463	PFA	C1-C6	2.33	1.44	1.40
2	B	463	PFA	C5-C4	2.37	1.41	1.38
2	B	463	PFA	C1-C2	2.96	1.45	1.40
2	A	462	PFA	C31-C21	3.02	1.44	1.38
2	A	462	PFA	C22-C21	3.19	1.44	1.38
2	B	463	PFA	C31-C21	3.30	1.45	1.38
2	B	463	PFA	C3-C2	3.39	1.44	1.39
2	B	463	PFA	C3-C4	4.44	1.44	1.38
2	A	462	PFA	C17-C10	4.72	1.46	1.39
2	B	463	PFA	C17-C10	5.46	1.47	1.39
2	A	462	PFA	C5-C4	5.49	1.45	1.38

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	462	PFA	C17-C10-N6	-7.82	117.96	123.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	463	PFA	C17-C10-N6	-5.82	119.28	123.11
2	B	463	PFA	C5-C4-N5	-5.53	112.40	119.13
2	A	462	PFA	C5-C4-N5	-5.05	112.97	119.13
2	B	463	PFA	C4-C5-C6	-3.92	118.18	120.12
2	A	462	PFA	C4-C5-C6	-3.74	118.27	120.12
2	A	462	PFA	C9-C25-C24	-3.24	106.06	111.77
2	A	462	PFA	C10-C17-N3	-2.93	118.69	121.78
2	B	463	PFA	C9-C25-C8	-2.65	103.74	110.34
2	A	462	PFA	C29-C31-C21	-2.64	116.44	119.74
2	A	462	PFA	C8-C25-C24	-2.37	107.60	111.77
2	A	462	PFA	O20-C1-C6	2.36	123.64	118.29
2	A	462	PFA	C22-C24-C26	2.38	118.87	116.89
2	B	463	PFA	C5-C6-C1	2.62	121.79	117.84
2	A	462	PFA	C4-C3-C2	4.56	122.38	120.12
2	B	463	PFA	C3-C4-N5	5.18	125.45	119.13
2	A	462	PFA	C3-C4-N5	6.64	127.22	119.13
2	B	463	PFA	C10-N6-C14	7.61	120.67	114.13
2	A	462	PFA	C10-N6-C14	8.81	121.70	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	462	PFA	4	0
2	B	463	PFA	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/258 (96%)	-0.30	5 (2%) 65 63	13, 23, 47, 109	0
1	B	239/258 (92%)	0.04	14 (5%) 23 22	18, 29, 56, 108	0
All	All	487/516 (94%)	-0.13	19 (3%) 40 38	13, 26, 52, 109	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	461	ASP	4.6
1	B	253	ALA	4.3
1	B	239	TRP	3.9
1	B	235	GLN	3.8
1	A	240	LYS	3.7
1	B	449	GLU	3.6
1	A	241	GLN	3.2
1	B	242	LYS	3.1
1	B	413	HIS	2.9
1	B	250	ILE	2.7
1	B	248	GLU	2.6
1	B	252	GLN	2.6
1	A	234	ALA	2.6
1	A	239	TRP	2.6
1	B	240	LYS	2.4
1	B	409	TYR	2.1
1	B	211	LYS	2.1
1	B	244	LYS	2.1
1	A	461	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PFA	B	463	27/27	0.95	0.13	0.07	17,25,29,29	0
2	PFA	A	462	27/27	0.96	0.11	-0.49	10,16,18,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.