



Full wwPDB NMR Structure Validation Report ⓘ

Feb 13, 2017 – 01:41 am GMT

PDB ID : 2N54
Title : Solution structure of a disulfide stabilized XCL1 dimer
Authors : Tyler, R.C.; Tuinstra, R.L.; Peterson, F.F.; Volkman, B.F.
Deposited on : 2015-07-07

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 80%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	93	
1	B	93	

2 Ensemble composition and analysis

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:51, B:210-B:251 (84)	0.50	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 8, 12, 13, 15, 17, 18, 19
2	3, 4, 5, 6, 9
3	10, 11, 14, 16
4	2, 7, 20

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2140 atoms, of which 1100 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Lymphotactin.

Mol	Chain	Residues	Atoms						Trace
1	A	66	Total	C	H	N	O	S	0
			1071	322	551	98	95	5	
1	B	66	Total	C	H	N	O	S	0
			1069	322	549	98	95	5	

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	CYS	ALA	engineered mutation	UNP P47992
A	49	CYS	ALA	engineered mutation	UNP P47992
B	236	CYS	ALA	engineered mutation	UNP P47992
B	249	CYS	ALA	engineered mutation	UNP P47992

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Lymphotactin



- Molecule 1: Lymphotactin




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin




4.2.2 Score per residue for model 2

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin




4.2.3 Score per residue for model 3

- Molecule 1: Lymphotactin



- Molecule 1: Lymphotactin






4.2.4 Score per residue for model 4


- Molecule 1: Lymphotactin

Chain A: 33% 11% 26% 29%



- Molecule 1: Lymphotactin


Chain B: 32% 12% 26% 29%



4.2.5 Score per residue for model 5


- Molecule 1: Lymphotactin

Chain A: 29% 14% 26% 29%




- Molecule 1: Lymphotactin

Chain B: 33% 12% 26% 29%




4.2.6 Score per residue for model 6

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotoctin




4.2.7 Score per residue for model 7

- Molecule 1: Lymphotoctin




- Molecule 1: Lymphotoctin



4.2.8 Score per residue for model 8


- Molecule 1: Lymphotoctin





• Molecule 1: Lymphotactin


Chain B: 27% 17% 26% 29%



4.2.9 Score per residue for model 9


• Molecule 1: Lymphotactin

Chain A: 33% 11% 26% 29%



• Molecule 1: Lymphotactin


Chain B: 32% 12% 26% 29%




4.2.10 Score per residue for model 10

• Molecule 1: Lymphotactin

Chain A: 26% 19% 26% 29%




- Molecule 1: Lymphotactin




4.2.11 Score per residue for model 11

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin




4.2.12 Score per residue for model 12

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin






4.2.13 Score per residue for model 13

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin




4.2.14 Score per residue for model 14

- Molecule 1: Lymphotactin




- Molecule 1: Lymphotactin




4.2.15 Score per residue for model 15

● Molecule 1: Lymphotactin




● Molecule 1: Lymphotactin




4.2.16 Score per residue for model 16

● Molecule 1: Lymphotactin




● Molecule 1: Lymphotactin




4.2.17 Score per residue for model 17

● Molecule 1: Lymphotactin







- Molecule 1: Lymphotoctin





4.2.18 Score per residue for model 18

- Molecule 1: Lymphotoctin





- Molecule 1: Lymphotoctin




4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: Lymphotoctin




● Molecule 1: Lymphotactin




4.2.20 Score per residue for model 20

● Molecule 1: Lymphotactin



● Molecule 1: Lymphotactin



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
TALOS	geometry optimization	
CYANA	structure solution	
CYANA	refinement	
X-PLOR NIH	geometry optimization	
X-PLOR NIH	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	2n54_cs.str
Number of chemical shift lists	1
Total number of shifts	1432
Number of shifts mapped to atoms	1432
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	80%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	B	0.0±0.0	0.1±0.2
All	All	0	1

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	B	235	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	325	351	351	11±2
1	B	325	351	351	11±2
All	All	13000	14040	14040	359

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:230:THR:HA	1:B:234:LEU:O	0.70	1.86	3	15

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:25:LYS:HE2	1:A:25:LYS:N	0.68	2.03	5	6
1:A:30:THR:HA	1:A:34:LEU:O	0.67	1.89	7	16
1:A:26:THR:HA	1:A:38:ILE:O	0.67	1.89	11	20
1:B:226:THR:HA	1:B:238:ILE:O	0.67	1.89	8	20
1:A:29:ILE:HD13	1:B:240:ILE:HD11	0.66	1.68	14	19
1:A:40:ILE:HD11	1:B:229:ILE:HD13	0.62	1.69	2	19
1:A:34:LEU:HD11	1:A:49:CYS:SG	0.62	2.34	10	1
1:A:25:LYS:HE2	1:A:25:LYS:H	0.61	1.55	5	2
1:A:25:LYS:NZ	1:A:40:ILE:HB	0.61	2.10	5	1
1:A:28:THR:HA	1:A:36:CYS:O	0.59	1.96	16	11
1:B:234:LEU:HD11	1:B:236:CYS:SG	0.59	2.38	10	5
1:A:27:TYR:O	1:A:37:VAL:HA	0.59	1.98	18	3
1:A:34:LEU:HD11	1:A:36:CYS:SG	0.58	2.38	17	4
1:A:14:LEU:HG	1:B:214:LEU:CD2	0.58	2.28	7	2
1:B:228:THR:HA	1:B:236:CYS:O	0.58	1.99	13	15
1:A:31:GLU:O	1:A:34:LEU:HB3	0.58	1.99	2	5
1:A:29:ILE:HB	1:A:36:CYS:HB3	0.57	1.76	11	10
1:A:34:LEU:HD21	1:A:49:CYS:SG	0.56	2.39	18	3
1:B:224:ILE:HA	1:B:240:ILE:O	0.56	2.01	5	8
1:B:231:GLU:O	1:B:234:LEU:HB3	0.56	2.01	1	4
1:B:217:GLN:HA	1:B:217:GLN:HE21	0.56	1.61	9	1
1:A:45:LEU:HD11	1:B:249:CYS:SG	0.55	2.41	18	1
1:B:229:ILE:HB	1:B:236:CYS:HB3	0.55	1.78	7	7
1:A:14:LEU:HG	1:B:214:LEU:HD21	0.55	1.77	7	3
1:A:14:LEU:HB3	1:B:212:VAL:CG2	0.52	2.33	12	1
1:B:225:LYS:N	1:B:225:LYS:HE2	0.51	2.20	7	1
1:A:39:PHE:HB2	1:A:46:LYS:HB3	0.51	1.82	9	13
1:A:19:LEU:HD23	1:A:19:LEU:H	0.50	1.67	8	1
1:A:49:CYS:SG	1:B:245:LEU:HD11	0.49	2.47	13	2
1:A:29:ILE:CG2	1:A:31:GLU:HG2	0.49	2.37	11	4
1:A:29:ILE:O	1:A:35:ARG:HA	0.49	2.07	1	2
1:A:17:GLN:HE21	1:A:17:GLN:HA	0.49	1.67	9	1
1:A:25:LYS:HE3	1:A:40:ILE:HB	0.48	1.85	3	6
1:A:39:PHE:CD1	1:A:46:LYS:HD2	0.48	2.43	4	1
1:A:25:LYS:H	1:A:25:LYS:HE2	0.48	1.68	13	1
1:A:12:VAL:CG2	1:B:214:LEU:HB3	0.48	2.38	8	1
1:A:15:THR:HG23	1:A:43:ARG:O	0.48	2.08	19	1
1:A:38:ILE:HD13	1:B:238:ILE:CD1	0.48	2.39	20	2
1:A:34:LEU:HD21	1:A:49:CYS:HB3	0.48	1.85	2	1
1:B:239:PHE:HB2	1:B:246:LYS:HB3	0.48	1.85	3	11
1:B:235:ARG:HB2	1:B:250:ASP:HB2	0.47	1.85	8	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:25:LYS:HD3	1:A:40:ILE:HD12	0.47	1.86	2	1
1:A:23:ARG:HA	1:A:23:ARG:NE	0.47	2.25	5	1
1:A:29:ILE:HD11	1:A:38:ILE:HD11	0.47	1.86	6	3
1:B:234:LEU:HD21	1:B:249:CYS:SG	0.47	2.50	20	2
1:A:24:ILE:HA	1:A:40:ILE:O	0.47	2.10	4	2
1:B:219:LEU:O	1:B:242:LYS:HA	0.47	2.10	15	1
1:A:29:ILE:HG21	1:B:240:ILE:CD1	0.47	2.40	17	1
1:A:12:VAL:HG13	1:B:214:LEU:HD12	0.47	1.85	11	1
1:B:225:LYS:HE3	1:B:240:ILE:HB	0.47	1.87	3	4
1:A:19:LEU:O	1:A:42:LYS:HA	0.46	2.10	16	2
1:B:217:GLN:HG3	1:B:218:ARG:H	0.46	1.70	6	1
1:B:229:ILE:HG22	1:B:231:GLU:OE2	0.46	2.10	18	1
1:A:24:ILE:HD12	1:A:41:THR:HA	0.46	1.88	6	5
1:B:229:ILE:HD11	1:B:238:ILE:HD11	0.46	1.86	18	5
1:A:49:CYS:SG	1:B:245:LEU:HD21	0.46	2.49	4	1
1:A:25:LYS:CE	1:A:40:ILE:HB	0.46	2.39	5	1
1:B:229:ILE:O	1:B:235:ARG:HA	0.46	2.10	12	4
1:A:18:ARG:O	1:A:18:ARG:HD3	0.46	2.10	17	1
1:B:229:ILE:CG2	1:B:231:GLU:HG2	0.46	2.40	12	2
1:B:242:LYS:H	1:B:242:LYS:HD3	0.46	1.71	1	1
1:B:224:ILE:HD12	1:B:241:THR:HA	0.46	1.86	2	6
1:A:31:GLU:CD	1:B:225:LYS:HZ3	0.46	2.14	3	1
1:B:242:LYS:HD3	1:B:242:LYS:H	0.45	1.70	13	1
1:A:36:CYS:HA	1:A:48:CYS:O	0.45	2.11	16	3
1:B:225:LYS:HE2	1:B:225:LYS:N	0.45	2.27	4	3
1:A:14:LEU:HD21	1:B:214:LEU:HG	0.45	1.89	12	3
1:B:236:CYS:HA	1:B:248:CYS:O	0.44	2.13	2	2
1:B:238:ILE:HA	1:B:246:LYS:O	0.44	2.13	2	2
1:A:25:LYS:H	1:A:25:LYS:CE	0.44	2.25	13	1
1:B:224:ILE:CD1	1:B:241:THR:HA	0.44	2.43	11	11
1:A:40:ILE:CD1	1:B:229:ILE:HG21	0.43	2.43	10	2
1:B:243:ARG:O	1:B:243:ARG:HG2	0.43	2.13	13	1
1:B:219:LEU:HD12	1:B:219:LEU:O	0.43	2.14	6	1
1:A:22:SER:HB2	1:B:231:GLU:OE1	0.43	2.13	2	1
1:A:24:ILE:CD1	1:A:41:THR:HA	0.43	2.43	10	2
1:A:38:ILE:CD1	1:B:238:ILE:HD13	0.43	2.44	10	2
1:A:11:CYS:HA	1:A:48:CYS:HA	0.43	1.90	8	1
1:A:18:ARG:HD3	1:A:18:ARG:H	0.43	1.73	14	1
1:A:16:THR:HG22	1:A:44:GLY:HA2	0.43	1.91	11	1
1:A:14:LEU:CD2	1:B:214:LEU:HG	0.42	2.44	12	1
1:B:225:LYS:CE	1:B:225:LYS:H	0.42	2.27	13	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:38:ILE:HD13	1:B:238:ILE:HD13	0.42	1.90	20	2
1:A:14:LEU:CD1	1:B:212:VAL:HB	0.42	2.44	2	2
1:A:43:ARG:O	1:A:43:ARG:HG2	0.42	2.15	6	1
1:B:225:LYS:HD3	1:B:240:ILE:HD12	0.42	1.90	2	1
1:A:34:LEU:HA	1:A:51:PRO:HA	0.42	1.92	7	1
1:A:40:ILE:CD1	1:B:229:ILE:HD13	0.42	2.44	5	2
1:A:14:LEU:HD12	1:B:212:VAL:HB	0.42	1.91	17	1
1:B:224:ILE:HG23	1:B:241:THR:HG22	0.41	1.92	20	1
1:A:45:LEU:HD21	1:B:249:CYS:SG	0.41	2.55	7	1
1:A:49:CYS:SG	1:B:219:LEU:HD21	0.41	2.55	15	1
1:A:25:LYS:CE	1:A:25:LYS:H	0.41	2.28	3	1
1:A:37:VAL:O	1:A:47:VAL:HA	0.41	2.16	5	1
1:A:36:CYS:HA	1:A:49:CYS:HA	0.41	1.93	11	1
1:B:211:CYS:HA	1:B:248:CYS:HA	0.41	1.92	10	1
1:A:29:ILE:CD1	1:B:240:ILE:HD11	0.41	2.46	18	1
1:B:236:CYS:HA	1:B:249:CYS:HA	0.40	1.93	12	2
1:B:225:LYS:HD2	1:B:240:ILE:HD12	0.40	1.93	14	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	42/93 (45%)	38±1 (90±3%)	4±2 (9±4%)	0±1 (1±2%)	21	68
1	B	42/93 (45%)	38±1 (90±3%)	4±1 (9±3%)	0±1 (1±1%)	23	70
All	All	1680/3720 (45%)	1511 (90%)	152 (9%)	17 (1%)	23	70

All 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	217	GLN	4
1	A	17	GLN	3
1	A	18	ARG	3
1	B	218	ARG	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	23	ARG	2
1	B	232	GLY	1
1	A	43	ARG	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	40/86 (47%)	38±1 (95±2%)	2±1 (5±2%)	33	79
1	B	40/86 (47%)	38±1 (95±2%)	2±1 (5±2%)	33	79
All	All	1600/3440 (47%)	1522 (95%)	78 (5%)	33	79

All 15 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	42	LYS	20
1	B	242	LYS	20
1	A	25	LYS	7
1	B	231	GLU	6
1	B	225	LYS	6
1	A	31	GLU	5
1	A	23	ARG	2
1	A	17	GLN	2
1	B	236	CYS	2
1	B	217	GLN	2
1	A	18	ARG	2
1	B	234	LEU	1
1	B	223	ARG	1
1	A	46	LYS	1
1	B	218	ARG	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 80% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: 2n54_cs.str

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1432
Number of shifts mapped to atoms	1432
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	10

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	130	-0.51 ± 0.13	Should be applied
$^{13}\text{C}_\beta$	122	-0.42 ± 0.11	None needed (< 0.5 ppm)
$^{13}\text{C}'$	92	0.31 ± 0.36	None needed (< 0.5 ppm)
^{15}N	122	-0.20 ± 0.28	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 80%, i.e. 834 atoms were assigned a chemical shift out of a possible 1048. 8 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	376/412 (91%)	160/164 (98%)	138/168 (82%)	78/80 (98%)
Sidechain	434/602 (72%)	258/352 (73%)	176/218 (81%)	0/32 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	24/34 (71%)	16/18 (89%)	8/16 (50%)	0/0 (—%)
Overall	834/1048 (80%)	434/534 (81%)	322/402 (80%)	78/112 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 76%, i.e. 1278 atoms were assigned a chemical shift out of a possible 1678. 10 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	588/652 (90%)	244/260 (94%)	222/264 (84%)	122/128 (95%)
Sidechain	642/968 (66%)	382/566 (67%)	260/340 (76%)	0/62 (0%)
Aromatic	48/58 (83%)	28/30 (93%)	18/26 (69%)	2/2 (100%)
Overall	1278/1678 (76%)	654/856 (76%)	500/630 (79%)	124/192 (65%)

7.1.4 Statistically unusual chemical shifts ⓘ


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	5	VAL	CG2	33.12	29.20 – 13.40	7.5
1	B	205	VAL	CG2	33.12	29.20 – 13.40	7.5
1	B	266	LYS	CD	36.05	34.86 – 23.06	6.0
1	A	66	LYS	CD	36.05	34.86 – 23.06	6.0
1	B	202	GLY	CA	52.88	51.81 – 38.91	5.8
1	A	2	GLY	CA	52.88	51.81 – 38.91	5.8
1	B	221	VAL	HB	3.76	3.59 – 0.39	5.5
1	A	21	VAL	HB	3.76	3.59 – 0.39	5.5
1	A	44	GLY	C	163.80	183.33 – 164.53	-5.4
1	B	244	GLY	C	163.80	183.33 – 164.53	-5.4

7.1.5 Random Coil Index (RCI) plots ⓘ

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

