



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 10:54 AM EDT

PDB ID : 3N5W
Title : Structure of neuronal nitric oxide synthase heme domain in complex with 6,6'-((2,2'-(pyridine-3,5-diyl)bis(ethane-2,1-diyl))bis(4-methylpyridin-2-amine)
Authors : Li, H.; Delker, S.L.; Poulos, T.L.
Deposited on : unknown
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2694 (1.76-1.72)
Clashscore	112137	2854 (1.76-1.72)
Ramachandran outliers	110173	2824 (1.76-1.72)
Sidechain outliers	110143	2824 (1.76-1.72)
RSRZ outliers	101464	2705 (1.76-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>15%</div> <div> <div></div> <div>91%</div> <div>5%</div> <div>.</div> </div> </div>
1	B	422	<div> <div>8%</div> <div> <div></div> <div>93%</div> <div>.</div> <div>.</div> </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7476 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	1	0
			3322	2129	566	606	21			
1	B	411	Total	C	N	O	S	0	2	0
			3357	2150	574	612	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 6,6'-(pyridine-3,5-diyl)diethane-2,1-diyl)bis(4-methylpyridin-2-amine) (three-letter code: XFJ) (formula: $C_{21}H_{25}N_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			26	21	5		
3	A	1	Total	C	N	0	0
			26	21	5		
3	B	1	Total	C	N	0	0
			26	21	5		
3	B	1	Total	C	N	0	0
			26	21	5		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Cl 1 1	0	0
5	A	1	Total Cl 1 1	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total Zn 3 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	263	Total O 263 263	0	0
7	B	331	Total O 331 331	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase



- Molecule 1: Nitric oxide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.34Å 111.21Å 164.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.11 – 1.73 38.11 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.11-1.73) 99.2 (38.11-1.73)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.73Å)	Xtriage
Refinement program	CNS, REFMAC 5.5.0102	Depositor
R, R_{free}	0.182 , 0.205 0.193 , 0.219	Depositor DCC
R_{free} test set	5018 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7476	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, HEM, ZN, XFJ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3419	0.57	0/4639
1	B	0.60	0/3457	0.60	0/4687
All	All	0.58	0/6876	0.58	0/9326

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3322	0	3230	16	0
1	B	3357	0	3273	10	0
2	A	43	0	30	2	0
2	B	43	0	30	2	0
3	A	52	0	50	1	0
3	B	52	0	50	1	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	263	0	0	1	0
7	B	331	0	0	2	0
All	All	7476	0	6669	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:ARG:HB3	1:A:318:LEU:HD21	1.61	0.81
1:A:371:ARG:HH21	1:A:371:ARG:CG	2.00	0.75
1:A:609:GLU:HG3	7:A:1047:HOH:O	1.97	0.63
1:A:628:GLN:HG2	1:B:631:VAL:HG11	1.81	0.61
2:B:750:HEM:HBB2	2:B:750:HEM:HHC	1.83	0.61
1:A:371:ARG:CG	1:A:371:ARG:NH2	2.65	0.59
1:A:371:ARG:HH21	1:A:371:ARG:HG3	1.70	0.56
1:A:299:ARG:CB	1:A:318:LEU:HD21	2.35	0.55
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.91	0.53
1:A:569:ASN:HD22	1:A:569:ASN:H	1.57	0.50
2:A:750:HEM:C1C	3:A:800:XFJ:H07B	2.46	0.50
1:B:364:GLN:NE2	7:B:1503:HOH:O	2.45	0.50
1:B:686:SER:HA	1:B:691:PHE:CG	2.46	0.50
1:A:371:ARG:HG2	1:A:371:ARG:NH2	2.27	0.49
1:A:371:ARG:HG2	1:A:371:ARG:HH21	1.77	0.48
1:A:686:SER:HA	1:A:691:PHE:CG	2.50	0.47
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.97	0.47
1:B:325:GLY:O	1:B:332:MET:HG3	2.15	0.45
1:B:299:ARG:HB3	1:B:318:LEU:HD21	1.98	0.45
1:B:535:GLN:HG3	7:B:1440:HOH:O	2.16	0.45
2:A:750:HEM:HHC	2:A:750:HEM:HBB2	1.99	0.44
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.06	0.43
2:B:750:HEM:C1C	3:B:800:XFJ:H07B	2.54	0.43
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.56	0.41
1:B:614:MET:HE3	1:B:632:GLU:HG3	2.02	0.41
1:A:307:GLU:HG2	1:A:692:HIS:CG	2.56	0.41
1:B:475:TRP:CG	1:B:710:PRO:HG2	2.55	0.41
1:A:361:PHE:O	1:A:364:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/422 (96%)	393 (97%)	11 (3%)	0	100	100
1	B	409/422 (97%)	404 (99%)	5 (1%)	0	100	100
All	All	813/844 (96%)	797 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/377 (97%)	356 (98%)	8 (2%)	57	33
1	B	368/377 (98%)	364 (99%)	4 (1%)	78	64
All	All	732/754 (97%)	720 (98%)	12 (2%)	68	49

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	350	THR
1	A	352	ASP
1	A	371	ARG
1	A	507	GLN
1	A	547	ARG
1	A	569	ASN
1	A	620	LYS
1	A	645	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	299	ARG
1	B	350	THR
1	B	353	GLN
1	B	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	454	ASN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	642	GLN
1	A	697	ASN
1	B	364	GLN
1	B	385	ASN
1	B	425	GLN
1	B	454	ASN
1	B	507	GLN
1	B	605	ASN
1	B	697	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 5 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	750	1	28,50,50	2.18	10 (35%)	17,82,82	2.00	3 (17%)
3	XFJ	A	800	-	28,28,28	0.62	0	38,38,38	2.14	10 (26%)
3	XFJ	A	805	6	28,28,28	0.64	0	38,38,38	1.82	11 (28%)
4	ACT	A	860	-	1,3,3	1.04	0	0,3,3	0.00	-
2	HEM	B	750	1	28,50,50	2.33	8 (28%)	17,82,82	1.66	3 (17%)
3	XFJ	B	800	-	28,28,28	0.75	0	38,38,38	2.03	11 (28%)
3	XFJ	B	805	6	28,28,28	0.59	0	38,38,38	1.66	9 (23%)
4	ACT	B	860	-	1,3,3	1.39	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/6/54/54	0/0/8/8
3	XFJ	A	800	-	-	0/10/10/10	0/3/3/3
3	XFJ	A	805	6	-	0/10/10/10	0/3/3/3
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/6/54/54	0/0/8/8
3	XFJ	B	800	-	-	0/10/10/10	0/3/3/3
3	XFJ	B	805	6	-	0/10/10/10	0/3/3/3
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3C-C2C	-5.31	1.33	1.40
2	A	750	HEM	C3B-C2B	-4.71	1.34	1.40
2	B	750	HEM	C3B-C2B	-4.58	1.34	1.40
2	A	750	HEM	C3C-C2C	-3.33	1.36	1.40
2	A	750	HEM	C1D-ND	2.12	1.40	1.36
2	A	750	HEM	CMC-C2C	2.18	1.56	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C1C-NC	2.20	1.39	1.36
2	B	750	HEM	CAA-C2A	2.27	1.55	1.52
2	A	750	HEM	CMD-C2D	2.53	1.56	1.51
2	A	750	HEM	CAA-C2A	2.55	1.56	1.52
2	A	750	HEM	C4D-ND	3.17	1.40	1.36
2	B	750	HEM	C4D-ND	3.53	1.40	1.36
2	A	750	HEM	C3B-CAB	3.69	1.55	1.47
2	A	750	HEM	C3C-CAC	3.90	1.55	1.47
2	B	750	HEM	C3C-CAC	4.03	1.55	1.47
2	B	750	HEM	C3B-CAB	4.05	1.55	1.47
2	B	750	HEM	C3D-C2D	4.14	1.50	1.37
2	A	750	HEM	C3D-C2D	4.65	1.51	1.37

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CBD-CAD-C3D	-5.36	102.24	112.47
2	B	750	HEM	CBD-CAD-C3D	-4.70	103.50	112.47
3	A	800	XFJ	C05-C06-N01	-3.73	118.87	122.91
2	A	750	HEM	CMA-C3A-C4A	-3.45	123.17	128.46
3	B	800	XFJ	C05-C06-N01	-3.41	119.21	122.91
3	B	805	XFJ	C08-C06-C05	-2.78	117.44	121.19
3	B	800	XFJ	C25-C26-N21	-2.73	119.95	122.91
3	A	805	XFJ	C25-C26-N21	-2.72	119.96	122.91
3	A	800	XFJ	C15-C14-C13	-2.72	117.83	121.26
3	A	800	XFJ	C24-C25-C26	-2.68	118.58	120.26
3	A	805	XFJ	C08-C06-C05	-2.57	117.73	121.19
3	B	800	XFJ	C15-C14-C13	-2.53	118.07	121.26
3	A	800	XFJ	C25-C26-N21	-2.48	120.23	122.91
3	B	800	XFJ	C09-C08-C06	-2.30	107.91	112.71
3	B	800	XFJ	C09-C13-C12	-2.29	118.08	121.82
2	B	750	HEM	CMA-C3A-C4A	-2.28	124.96	128.46
3	A	805	XFJ	C04-C05-C06	-2.17	118.90	120.26
3	A	805	XFJ	C09-C13-C12	-2.07	118.44	121.82
3	B	805	XFJ	C15-C14-C13	-2.05	118.66	121.26
3	B	800	XFJ	C12-N11-C16	2.14	120.48	117.45
3	A	805	XFJ	C14-C13-C12	2.16	118.80	116.71
3	B	800	XFJ	C14-C13-C12	2.20	118.83	116.71
3	A	805	XFJ	C12-N11-C16	2.23	120.60	117.45
2	B	750	HEM	CMB-C2B-C3B	2.29	129.15	124.89
3	A	805	XFJ	C14-C15-C16	2.32	118.95	116.71
3	A	805	XFJ	C18-C26-N21	2.32	119.13	115.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	805	XFJ	N02-C02-N01	2.34	120.59	116.64
3	A	800	XFJ	C08-C06-N01	2.37	119.21	115.78
3	B	805	XFJ	C02-N01-C06	2.45	119.90	118.17
3	B	800	XFJ	C08-C06-N01	2.61	119.56	115.78
3	B	805	XFJ	C14-C15-C16	2.66	119.28	116.71
3	A	800	XFJ	C18-C26-N21	2.67	119.64	115.78
2	A	750	HEM	CMC-C2C-C3C	2.71	129.93	124.89
3	B	805	XFJ	C18-C26-N21	2.73	119.73	115.78
3	B	805	XFJ	C14-C13-C12	2.95	119.56	116.71
3	A	800	XFJ	C14-C15-C16	2.97	119.58	116.71
3	A	800	XFJ	C14-C13-C12	3.05	119.65	116.71
3	B	800	XFJ	C14-C15-C16	3.46	120.05	116.71
3	B	805	XFJ	C08-C06-N01	3.48	120.81	115.78
3	A	805	XFJ	C08-C06-N01	3.64	121.04	115.78
3	B	805	XFJ	C22-N21-C26	3.74	120.82	118.17
3	A	805	XFJ	C22-N21-C26	4.24	121.17	118.17
3	A	805	XFJ	C02-N01-C06	4.26	121.18	118.17
3	B	800	XFJ	C22-N21-C26	5.03	121.73	118.17
3	A	800	XFJ	C02-N01-C06	6.07	122.46	118.17
3	A	800	XFJ	C22-N21-C26	6.29	122.62	118.17
3	B	800	XFJ	C02-N01-C06	6.30	122.62	118.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	2	0
3	A	800	XFJ	1	0
2	B	750	HEM	2	0
3	B	800	XFJ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.86	64 (15%) 2 3	18, 35, 62, 78	0
1	B	411/422 (97%)	0.43	33 (8%) 13 17	17, 28, 49, 68	0
All	All	818/844 (96%)	0.64	97 (11%) 5 7	17, 31, 58, 78	0

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	488	PRO	7.6
1	B	300	PHE	7.0
1	A	300	PHE	6.5
1	B	348	VAL	6.4
1	B	350	THR	6.4
1	A	352	ASP	6.4
1	A	715	VAL	5.8
1	A	350	THR	5.7
1	B	715	VAL	5.7
1	A	712	ASN	5.4
1	B	619	ARG	5.4
1	A	351	LYS	5.3
1	A	713	THR	4.9
1	A	299	ARG	4.9
1	A	716	TRP	4.9
1	A	490	GLY	4.7
1	B	322	LEU	4.7
1	A	355	PHE	4.4
1	A	486	LYS	4.3
1	A	491	SER	4.2
1	A	321	THR	4.0
1	B	352	ASP	3.9
1	A	322	LEU	3.7
1	A	391	THR	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	567	VAL	3.6
1	B	718	GLY	3.5
1	A	385	ASN	3.4
1	A	619	ARG	3.4
1	A	390	SER	3.4
1	B	321	THR	3.4
1	A	338	PRO	3.4
1	A	489	ASP	3.3
1	A	392	SER	3.3
1	A	386	LYS	3.3
1	A	507	GLN	3.1
1	B	620	LYS	3.1
1	B	677	VAL	3.1
1	B	299	ARG	3.1
1	B	616	LEU	3.0
1	A	469	LYS	3.0
1	A	503	GLU	3.0
1	A	677	VAL	3.0
1	B	680	VAL	3.0
1	A	511	LYS	3.0
1	A	676	TRP	3.0
1	B	691	PHE	3.0
1	A	328	GLU	2.9
1	B	667	ARG	2.9
1	A	389	GLU	2.9
1	B	713	THR	2.8
1	B	338	PRO	2.8
1	A	667	ARG	2.8
1	A	679	ILE	2.7
1	A	680	VAL	2.7
1	B	567	VAL	2.7
1	A	593	ILE	2.7
1	A	382	GLU	2.7
1	A	388	ILE	2.7
1	A	552	ASP	2.7
1	B	679	ILE	2.6
1	A	479	LEU	2.6
1	B	615	ASP	2.6
1	A	711	TRP	2.5
1	A	353	GLN	2.5
1	A	714	HIS	2.5
1	A	480	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	678	TRP	2.5
1	B	595	VAL	2.4
1	A	710	PRO	2.4
1	A	393	THR	2.4
1	B	479	LEU	2.4
1	A	588	TYR	2.4
1	A	487	GLN	2.3
1	B	584	PHE	2.3
1	A	504	ILE	2.3
1	A	415	CYS	2.3
1	A	467	ASP	2.3
1	A	591	THR	2.3
1	A	704	PHE	2.3
1	B	617	ASP	2.3
1	B	591	THR	2.2
1	A	519	VAL	2.2
1	A	561	TRP	2.2
1	A	311	VAL	2.2
1	A	370	LYS	2.1
1	A	595	VAL	2.1
1	B	389	GLU	2.1
1	B	351	LYS	2.1
1	A	506	ILE	2.1
1	B	561	TRP	2.1
1	A	512	ALA	2.0
1	B	309	ASP	2.0
1	B	353	GLN	2.0
1	A	330	ILE	2.0
1	B	668	CYS	2.0
1	A	302	LYS	2.0
1	B	676	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	A	860	4/4	0.93	0.12	1.50	40,41,42,43	0
2	HEM	A	750	43/43	0.97	0.19	1.18	18,20,29,33	0
2	HEM	B	750	43/43	0.98	0.16	1.08	16,18,24,29	0
3	XFJ	A	800	26/26	0.94	0.19	0.85	21,25,28,31	0
3	XFJ	B	800	26/26	0.96	0.16	0.66	17,23,26,29	0
4	ACT	B	860	4/4	0.96	0.08	0.41	29,32,33,34	0
3	XFJ	A	805	26/26	0.88	0.17	0.15	26,29,43,44	0
5	CL	A	910	1/1	0.97	0.16	0.07	27,27,27,27	0
3	XFJ	B	805	26/26	0.90	0.15	0.02	25,29,37,39	0
5	CL	B	910	1/1	0.98	0.14	-0.19	25,25,25,25	0
6	ZN	A	900	1/1	0.99	0.08	-0.28	26,26,26,26	0
6	ZN	A	901	1/1	0.99	0.13	-	22,22,22,22	1
6	ZN	A	719	1/1	0.99	0.14	-	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.