



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 9, 2017 – 04:48 PM EST

PDB ID : 4N8H
Title : E61V mutant, RipA structure
Authors : Torres, R.; Goulding, C.W.
Deposited on : unknown
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 88%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 3%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div> <div style="width: 1%; background-color: red;"></div> <div style="width: 88%; background-color: green;"></div> <div style="width: 8%; background-color: yellow;"></div> <div style="width: 3%; background-color: grey;"></div> </div> </div>
1	B	476	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 88%; height: 10px; background-color: green;"></div> <div style="width: 8%; height: 10px; background-color: yellow;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 88%; background-color: green;"></div> <div style="width: 8%; background-color: yellow;"></div> <div style="width: 2%; background-color: grey;"></div> </div> </div>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 13860 atoms, of which 6744 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxybutyrate coenzyme A transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	440	Total	C	H	N	O	S	0	0	0
			6766	2134	3368	610	637	17			
1	B	440	Total	C	H	N	O	S	0	0	0
			6774	2134	3376	610	637	17			

There are 74 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-35	MET	-	INITIATING METHIONINE	UNP Q9ZC36
A	-34	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-33	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-32	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-31	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-30	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-29	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-28	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-27	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-26	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-25	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-24	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-23	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-22	LEU	-	EXPRESSION TAG	UNP Q9ZC36
A	-21	VAL	-	EXPRESSION TAG	UNP Q9ZC36
A	-20	PRO	-	EXPRESSION TAG	UNP Q9ZC36
A	-19	ARG	-	EXPRESSION TAG	UNP Q9ZC36
A	-18	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-17	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-16	HIS	-	EXPRESSION TAG	UNP Q9ZC36
A	-15	MET	-	EXPRESSION TAG	UNP Q9ZC36
A	-14	ALA	-	EXPRESSION TAG	UNP Q9ZC36
A	-13	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-12	MET	-	EXPRESSION TAG	UNP Q9ZC36
A	-11	THR	-	EXPRESSION TAG	UNP Q9ZC36

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-9	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-8	GLN	-	EXPRESSION TAG	UNP Q9ZC36
A	-7	GLN	-	EXPRESSION TAG	UNP Q9ZC36
A	-6	MET	-	EXPRESSION TAG	UNP Q9ZC36
A	-5	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-4	ARG	-	EXPRESSION TAG	UNP Q9ZC36
A	-3	GLY	-	EXPRESSION TAG	UNP Q9ZC36
A	-2	SER	-	EXPRESSION TAG	UNP Q9ZC36
A	-1	GLU	-	EXPRESSION TAG	UNP Q9ZC36
A	0	PHE	-	EXPRESSION TAG	UNP Q9ZC36
A	61	VAL	GLU	ENGINEERED MUTATION	UNP Q9ZC36
B	-35	MET	-	INITIATING METHIONINE	UNP Q9ZC36
B	-34	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-33	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-32	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-31	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-30	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-29	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-28	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-27	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-26	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-25	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-24	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-23	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-22	LEU	-	EXPRESSION TAG	UNP Q9ZC36
B	-21	VAL	-	EXPRESSION TAG	UNP Q9ZC36
B	-20	PRO	-	EXPRESSION TAG	UNP Q9ZC36
B	-19	ARG	-	EXPRESSION TAG	UNP Q9ZC36
B	-18	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-17	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-16	HIS	-	EXPRESSION TAG	UNP Q9ZC36
B	-15	MET	-	EXPRESSION TAG	UNP Q9ZC36
B	-14	ALA	-	EXPRESSION TAG	UNP Q9ZC36
B	-13	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-12	MET	-	EXPRESSION TAG	UNP Q9ZC36
B	-11	THR	-	EXPRESSION TAG	UNP Q9ZC36
B	-10	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-9	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-8	GLN	-	EXPRESSION TAG	UNP Q9ZC36
B	-7	GLN	-	EXPRESSION TAG	UNP Q9ZC36
B	-6	MET	-	EXPRESSION TAG	UNP Q9ZC36

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-4	ARG	-	EXPRESSION TAG	UNP Q9ZC36
B	-3	GLY	-	EXPRESSION TAG	UNP Q9ZC36
B	-2	SER	-	EXPRESSION TAG	UNP Q9ZC36
B	-1	GLU	-	EXPRESSION TAG	UNP Q9ZC36
B	0	PHE	-	EXPRESSION TAG	UNP Q9ZC36
B	61	VAL	GLU	ENGINEERED MUTATION	UNP Q9ZC36


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	177	Total O 177 177	0	0
2	B	143	Total O 143 143	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-hydroxybutyrate coenzyme A transferase



- Molecule 1: 4-hydroxybutyrate coenzyme A transferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	118.91Å 109.00Å 85.01Å 90.00° 120.07° 90.00°	Depositor
Resolution (Å)	39.70 – 2.40 39.70 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.7 (39.70-2.40) 98.5 (39.70-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.50 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.8.3_1479	Depositor
R, R_{free}	0.151 , 0.192 0.153 , 0.190	Depositor DCC
R_{free} test set	1985 reflections (5.48%)	DCC
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13860	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3463	0.53	2/4699 (0.0%)
1	B	0.59	1/3463 (0.0%)	0.53	1/4699 (0.0%)
All	All	0.59	1/6926 (0.0%)	0.53	3/9398 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	151	SER	CB-OG	-5.51	1.35	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	80	LYS	C-N-CD	5.92	140.83	128.40
1	B	80	LYS	C-N-CD	5.67	140.31	128.40
1	A	169	MET	C-N-CD	5.20	139.32	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3398	3368	3405	8	0
1	B	3398	3376	3405	10	0
2	A	177	0	0	0	0
2	B	143	0	0	2	0
All	All	7116	6744	6810	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:341:GLN:OE1	1:B:343:SER:HB2	2.05	0.57
1:A:108:TYR:OH	1:A:345:SER:OG	2.16	0.57
1:B:86:VAL:HG21	1:B:342:TYR:CD1	2.42	0.55
1:A:230:LEU:HB3	1:A:231:PRO:HD3	1.92	0.52
1:B:335:GLU:HB2	1:B:343:SER:OG	2.08	0.52
1:B:127:ASP:OD1	1:B:159:ARG:NH2	2.46	0.49
1:B:286:GLU:HG2	2:B:621:HOH:O	2.15	0.46
1:B:230:LEU:HB3	1:B:231:PRO:HD3	1.97	0.46
1:A:167:ARG:HG2	1:A:191:HIS:ND1	2.32	0.45
1:B:34:ALA:HB1	1:B:131:HIS:CE1	2.53	0.43
1:A:250:VAL:HG22	1:A:280:PHE:CE1	2.54	0.43
1:A:329:THR:HB	1:A:391:ARG:HD3	2.02	0.42
1:B:348:GLN:NE2	2:B:534:HOH:O	2.47	0.42
1:B:168:TYR:CD2	1:B:193:PRO:HA	2.54	0.42
1:A:35:GLU:HA	1:A:36:PRO:HD3	1.87	0.41
1:A:119:LEU:HD12	1:A:123:GLU:HG3	2.02	0.41
1:A:7:TYR:HA	1:A:181:ILE:HD11	2.03	0.41
1:B:206:TYR:CD2	1:B:233:LEU:HD12	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/476 (92%)	427 (98%)	11 (2%)	0	100	100
1	B	438/476 (92%)	426 (97%)	12 (3%)	0	100	100
All	All	876/952 (92%)	853 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	366/394 (93%)	362 (99%)	4 (1%)	78	90
1	B	366/394 (93%)	362 (99%)	4 (1%)	78	90
All	All	732/788 (93%)	724 (99%)	8 (1%)	78	90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	LYS
1	A	83	SER
1	A	232	ASN
1	A	279	THR
1	B	72	ARG
1	B	83	SER
1	B	279	THR
1	B	286	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	440/476 (92%)	-0.36	7 (1%) 72 70	14, 27, 59, 106	0
1	B	440/476 (92%)	-0.30	10 (2%) 61 58	15, 31, 66, 99	0
All	All	880/952 (92%)	-0.33	17 (1%) 67 64	14, 29, 61, 106	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	338	LEU	4.9
1	A	339	GLY	4.0
1	A	345	SER	3.8
1	B	343	SER	3.8
1	A	338	LEU	3.8
1	A	343	SER	3.1
1	B	340	HIS	3.0
1	B	373	LYS	2.8
1	A	388	THR	2.6
1	B	375	THR	2.3
1	B	95	ARG	2.2
1	B	376	VAL	2.2
1	B	436	LYS	2.2
1	B	4	ARG	2.2
1	A	373	LYS	2.2
1	A	372	ALA	2.1
1	B	440	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.