



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2018 – 01:34 PM EST

PDB ID : 5N8S
Title : Crystal Structure of Drosophila DHX36 helicase in complex with polyT
Authors : Chen, W.-F.; Rety, S.; Hai-Lei Guo, H.-L.; Wu, W.-Q.; Liu, N.-N.; Liu, Q.-W.;
Dai, Y.-X.; Xi, X.-G.
Deposited on : 2017-02-24
Resolution : 2.88 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2135 (2.90-2.86)
Clashscore	112137	2400 (2.90-2.86)
Ramachandran outliers	110173	2346 (2.90-2.86)
Sidechain outliers	110143	2349 (2.90-2.86)
RSRZ outliers	101464	2149 (2.90-2.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	944	<div> <div>4%</div> <div> <div></div> <div>47%</div> <div>39%</div> <div>• • 10%</div> </div> </div>
1	B	944	<div> <div>2%</div> <div> <div></div> <div>43%</div> <div>42%</div> <div>• • 10%</div> </div> </div>
2	C	16	<div> <div></div> <div> <div>44%</div> <div>13%</div> <div>6%</div> <div>38%</div> </div> </div>
2	D	16	<div> <div></div> <div> <div>38%</div> <div>25%</div> <div>38%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14069 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CG9323, isoform A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	850	Total	C	N	O	S	0	0	0
			6825	4310	1204	1266	45			
1	B	851	Total	C	N	O	S	0	0	0
			6834	4315	1205	1269	45			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	943	VAL	-	expression tag	UNP Q8SWT2
A	944	ASP	-	expression tag	UNP Q8SWT2
B	943	VAL	-	expression tag	UNP Q8SWT2
B	944	ASP	-	expression tag	UNP Q8SWT2

- Molecule 2 is a DNA chain called DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			200	100	20	70	10			
2	D	10	Total	C	N	O	P	0	0	0
			200	100	20	70	10			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

3 Residue-property plots



These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CG9323, isoform A







• Molecule 1: CG9323, isoform A



• Molecule 2: DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3')



- Molecule 2: DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	297.98Å 49.92Å 163.85Å 90.00° 115.01° 90.00°	Depositor
Resolution (Å)	81.10 – 2.88 83.80 – 2.88	Depositor EDS
% Data completeness (in resolution range)	99.4 (81.10-2.88) 99.4 (83.80-2.88)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.86Å)	Xtriage
Refinement program	PHENIX (1.10 _2155: ???)	Depositor
R, R_{free}	0.207 , 0.285 0.205 , 0.284	Depositor DCC
R_{free} test set	2474 reflections (4.93%)	DCC
Wilson B-factor (Å ²)	70.4	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 52.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14069	wwPDB-VP
Average B, all atoms (Å ²)	88.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	14/6944 (0.2%)	1.01	33/9367 (0.4%)
1	B	0.67	15/6953 (0.2%)	0.97	29/9379 (0.3%)
2	C	1.02	0/219	1.40	1/336 (0.3%)
2	D	1.04	0/219	1.42	0/336
All	All	0.69	29/14335 (0.2%)	1.01	63/19418 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8
1	B	0	6
All	All	0	14

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	140	GLU	CD-OE1	-11.02	1.13	1.25
1	B	691	ARG	NE-CZ	-10.27	1.19	1.33
1	A	140	GLU	CD-OE2	-9.82	1.14	1.25
1	B	691	ARG	CD-NE	-9.52	1.30	1.46
1	B	518	GLU	CD-OE1	-8.58	1.16	1.25

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	151	LYS	CD-CE-NZ	-16.01	74.88	111.70
1	A	146	ARG	NE-CZ-NH2	-15.51	112.55	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	396	ARG	NE-CZ-NH1	15.10	127.85	120.30
1	B	691	ARG	NE-CZ-NH1	-13.51	113.55	120.30
1	B	691	ARG	CD-NE-CZ	12.77	141.47	123.60

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	140	GLU	Peptide
1	A	141	GLU	Peptide
1	A	353	PHE	Peptide
1	A	542	GLY	Peptide
1	A	678	ARG	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6825	0	6925	491	1
1	B	6834	0	6930	501	0
2	C	200	0	121	4	0
2	D	200	0	121	5	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
All	All	14069	0	14097	993	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 993 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:141:GLU:CB	1:B:145:LYS:HZ1	1.18	1.55
1:A:67:ARG:NH2	1:A:70:GLU:HB2	1.32	1.42
1:A:138:GLN:C	1:A:144:LYS:HE3	1.50	1.32

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:394:LYS:NZ	1:A:397:LEU:HD13	1.45	1.28
1:B:141:GLU:HB3	1:B:145:LYS:NZ	1.45	1.27

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:GLU:OE2	1:A:555:ARG:NH2[2_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	842/944 (89%)	811 (96%)	27 (3%)	4 (0%)	32 66
1	B	843/944 (89%)	802 (95%)	38 (4%)	3 (0%)	38 70
All	All	1685/1888 (89%)	1613 (96%)	65 (4%)	7 (0%)	38 70

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	518	GLU
1	B	862	GLN
1	A	713	GLU
1	A	334	MET
1	A	835	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	758/842 (90%)	734 (97%)	24 (3%)	44	77
1	B	759/842 (90%)	724 (95%)	35 (5%)	31	64
All	All	1517/1684 (90%)	1458 (96%)	59 (4%)	37	70

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	115	ARG
1	B	265	SER
1	B	798	MET
1	B	137	ARG
1	B	151	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	814	ASN
1	B	138	GLN
1	B	471	GLN
1	B	57	ASN
1	B	136	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	A	1001	1	4,4,4	0.65	0	6,6,6	0.47	0
3	PO4	B	1001	-	4,4,4	0.74	0	6,6,6	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	1001	1	-	0/0/0/0	0/0/0/0
3	PO4	B	1001	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	850/944 (90%)	0.25	40 (4%) 32 28	43, 85, 149, 215	0
1	B	851/944 (90%)	0.09	18 (2%) 64 61	43, 78, 137, 214	0
2	C	10/16 (62%)	-0.24	0 100 100	66, 86, 131, 152	0
2	D	10/16 (62%)	-0.29	0 100 100	59, 74, 118, 140	0
All	All	1721/1920 (89%)	0.17	58 (3%) 46 40	43, 82, 142, 215	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	785	SER	5.9
1	A	136	GLN	4.9
1	A	52	ILE	4.8
1	B	355	LYS	4.3
1	B	709	TYR	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PO4	B	1001	5/5	0.94	0.22	0.94	102,111,117,121	0
3	PO4	A	1001	5/5	0.86	0.18	-0.09	118,144,149,150	0

6.5 Other polymers [i](#)

There are no such residues in this entry.