



wwPDB X-ray Structure Validation Summary Report

Feb 14, 2017 – 01:16 am GMT

PDB ID : 1NAN
Title : MCH CLASS I H-2KB MOLECULE COMPLEXED WITH PBM1 PEPTIDE
Authors : Reiser, J.-B.; Darnault, C.; Gregoire, C.; Mosser, T.; Mazza, G.; Kearney, A.;
van der Merwe, P.A.; Fontecilla-Camps, J.C.; Housset, D.; Malissen, B.
Deposited on : 2002-11-28
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	278	
1	L	278	
2	I	99	
2	P	99	
3	M	8	
3	Q	8	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, K-B alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	278	Total	C	N	O	S	0	0	0
			2262	1428	397	428	9			
1	L	278	Total	C	N	O	S	0	0	0
			2262	1428	397	428	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	P	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			
2	I	99	Total	C	N	O	S	0	0	0
			821	524	138	152	7			

- Molecule 3 is a protein called pBM1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	M	8	Total	C	N	O	0	0	0
			70	46	10	14			
3	Q	8	Total	C	N	O	0	0	0
			70	46	10	14			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	46	Total	O	0	0
			46	46		
4	I	24	Total	O	0	0
			24	24		
4	L	51	Total	O	0	0
			51	51		
4	M	6	Total	O	0	0
			6	6		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	P	28	Total 28	O 28	0	0
4	Q	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain



- Molecule 1: H-2 class I histocompatibility antigen, K-B alpha chain




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin





- Molecule 3: pBM1 peptide

Chain M: 63% 25% 13%



- Molecule 3: pBM1 peptide

Chain Q: 88% 13%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.98Å 88.61Å 89.17Å 90.00° 111.60° 90.00°	Depositor
Resolution (Å)	12.00 – 2.30 61.26 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) (12.00-2.30) 97.9 (61.26-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.14 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.224 , 0.279 0.229 , 0.285	Depositor DCC
R_{free} test set	4144 reflections (11.06%)	DCC
Wilson B-factor (Å ²)	35.5	Xtrriage
Anisotropy	0.823	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6462	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.98 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.3972e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.50	0/2326	0.77	5/3161 (0.2%)
1	L	0.52	0/2326	0.78	8/3161 (0.3%)
2	I	0.56	0/847	0.78	3/1148 (0.3%)
2	P	0.55	0/847	0.79	1/1148 (0.1%)
3	M	0.58	0/71	0.82	0/94
3	Q	0.58	0/71	0.83	0/94
All	All	0.52	0/6488	0.78	17/8806 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	137	ASP	CB-CG-OD2	7.23	124.81	118.30
1	L	212	ASP	CB-CG-OD2	6.93	124.53	118.30
1	L	119	ASP	CB-CG-OD2	6.67	124.30	118.30
1	H	129	ASP	CB-CG-OD2	6.10	123.79	118.30
1	H	137	ASP	CB-CG-OD2	5.99	123.69	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	195	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	2262	0	2150	55	0
1	L	2262	0	2150	64	0
2	I	821	0	796	13	0
2	P	821	0	796	10	0
3	M	70	0	65	5	0
3	Q	70	0	65	0	0
4	H	46	0	0	4	0
4	I	24	0	0	1	0
4	L	51	0	0	2	0
4	M	6	0	0	1	0
4	P	28	0	0	0	0
4	Q	1	0	0	0	0
All	All	6462	0	6022	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:1:ILE:H1	2:I:1:ILE:HD12	1.33	0.93
1:H:19:GLU:OE1	1:H:75:ARG:NH2	2.02	0.92
1:H:194:ARG:O	1:H:196:GLU:HA	1.70	0.90
1:H:21:ARG:HH11	1:H:21:ARG:HG3	1.37	0.89
2:I:1:ILE:N	2:I:1:ILE:CD1	2.38	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	276/278 (99%)	258 (94%)	9 (3%)	9 (3%)	4	2
1	L	276/278 (99%)	249 (90%)	16 (6%)	11 (4%)	3	1
2	I	97/99 (98%)	92 (95%)	5 (5%)	0	100	100
2	P	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
3	M	6/8 (75%)	6 (100%)	0	0	100	100
3	Q	6/8 (75%)	6 (100%)	0	0	100	100
All	All	758/770 (98%)	706 (93%)	32 (4%)	20 (3%)	6	4

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	177	ALA
1	H	178	THR
1	L	138	MET
1	L	176	ASN
1	L	225	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	236/236 (100%)	201 (85%)	35 (15%)	3	3
1	L	236/236 (100%)	200 (85%)	36 (15%)	3	3
2	I	94/94 (100%)	87 (93%)	7 (7%)	16	20
2	P	94/94 (100%)	86 (92%)	8 (8%)	12	15
3	M	8/8 (100%)	7 (88%)	1 (12%)	5	5
3	Q	8/8 (100%)	7 (88%)	1 (12%)	5	5
All	All	676/676 (100%)	588 (87%)	88 (13%)	5	5

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	44	ARG
1	L	110	LEU
2	I	50	GLU
1	L	45	TYR
1	L	73	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	149	GLN
1	L	226	GLN
2	P	38	GLN
1	L	114	GLN
2	I	8	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	278/278 (100%)	0.49	11 (3%) 39 46	27, 42, 64, 78	10 (3%)
1	L	278/278 (100%)	0.48	16 (5%) 24 30	25, 42, 64, 77	4 (1%)
2	I	99/99 (100%)	0.08	0 100 100	25, 37, 49, 56	4 (4%)
2	P	99/99 (100%)	0.10	0 100 100	26, 38, 49, 60	2 (2%)
3	M	8/8 (100%)	0.09	0 100 100	33, 36, 44, 44	0
3	Q	8/8 (100%)	0.21	0 100 100	30, 42, 56, 59	0
All	All	770/770 (100%)	0.37	27 (3%) 44 51	25, 40, 63, 78	20 (2%)

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	178	THR	12.4
1	H	177	ALA	11.1
1	H	278	PRO	6.0
1	H	180	LEU	5.6
1	L	277	PRO	5.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.