



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2017 – 05:08 PM EDT

PDB ID : 5NA5  
Title : Pseudomonas fluorescens kynurenine 3-monooxygenase (KMO) apo structure  
Authors : Rowland, P.  
Deposited on : 2017-02-27  
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029077  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029077


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3233 (1.96-1.92)
Clashscore	112137	3430 (1.96-1.92)
Ramachandran outliers	110173	3395 (1.96-1.92)
Sidechain outliers	110143	3395 (1.96-1.92)
RSRZ outliers	101464	3250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	<div> <div>3%</div> <div>94%</div> <div>5%</div> <div>...</div> </div>
1	B	461	<div> <div>6%</div> <div>93%</div> <div>5%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	503	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7725 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	3	0
			3482	2182	643	639	18			
1	B	451	Total	C	N	O	S	0	4	0
			3511	2201	650	642	18			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	252	SER	CYS	engineered mutation	UNP Q84HF5
A	461	SER	CYS	engineered mutation	UNP Q84HF5
B	252	SER	CYS	engineered mutation	UNP Q84HF5
B	461	SER	CYS	engineered mutation	UNP Q84HF5

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	338	Total	O	0	0
			338	338		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	274	Total 274	O 274	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kynurenine 3-monooxygenase



- Molecule 1: Kynurenine 3-monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.43Å 53.74Å 137.56Å 90.00° 103.64° 90.00°	Depositor
Resolution (Å)	133.68 – 1.94 133.68 – 1.94	Depositor EDS
% Data completeness (in resolution range)	96.0 (133.68-1.94) 96.4 (133.68-1.94)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 1.94Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, $R_{free}$	0.201 , 0.231 0.201 , 0.229	Depositor DCC
$R_{free}$ test set	2854 reflections (4.15%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7725	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.19 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 6.0683e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/3566	0.59	0/4843
1	B	0.50	0/3599	0.59	0/4888
All	All	0.50	0/7165	0.59	0/9731

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3482	0	3446	7	0
1	B	3511	0	3487	7	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	1	0	0	1	0
3	B	1	0	0	1	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	338	0	0	1	0
5	B	274	0	0	0	0
All	All	7725	0	7011	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:877:HOH:O	1:B:298:THR:HG21	1.96	0.65
1:B:184:GLU:HG2	1:B:297:LEU:HD22	1.77	0.65
1:A:184:GLU:HG2	1:A:297:LEU:HD22	1.87	0.56
1:A:84:ARG:HG3	1:A:215:ILE:HD12	1.90	0.53
1:B:318:PRO:HA	3:B:502:CL:CL	2.48	0.51
1:A:113:ARG:NH2	1:A:117:ILE:HD11	2.26	0.50
1:A:318:PRO:HA	3:A:502:CL:CL	2.49	0.49
1:B:84:ARG:HD2	1:B:215:ILE:HD12	1.97	0.46
1:A:100:ARG:H	1:A:100:ARG:HG3	1.57	0.45
1:A:295:LEU:HB3	1:A:315:PRO:HD2	2.01	0.42
1:B:319:PHE:CD2	1:B:370:TYR:HB2	2.55	0.42
1:A:438:LEU:HA	1:A:441:ILE:HD12	2.03	0.41
1:B:53:ILE:HD11	1:B:111:ARG:HH21	1.85	0.41
1:B:215:ILE:HG22	1:B:217:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/461 (97%)	437 (98%)	10 (2%)	0	100	100
1	B	453/461 (98%)	445 (98%)	6 (1%)	2 (0%)	38	25
All	All	900/922 (98%)	882 (98%)	16 (2%)	2 (0%)	51	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	102	ASP
1	B	377	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/372 (98%)	358 (98%)	6 (2%)	68	61
1	B	367/372 (99%)	356 (97%)	11 (3%)	46	32
All	All	731/744 (98%)	714 (98%)	17 (2%)	56	45

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ARG
1	A	113	ARG
1	A	158	ARG
1	A	381	THR
1	A	427	LEU
1	A	438	LEU
1	B	51	ARG
1	B	72	ARG
1	B	101	ASP
1	B	104	GLU
1	B	158	ARG
1	B	197	GLN
1	B	242	GLN
1	B	272	ASP
1	B	374	SER
1	B	427	LEU
1	B	438	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	GLN
1	B	197	GLN
1	B	242	GLN
1	B	269	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	501	-	51,58,58	1.02	5 (9%)	54,89,89	2.18	6 (11%)
4	GOL	A	503	-	5,5,5	0.14	0	5,5,5	0.21	0
2	FAD	B	501	-	51,58,58	1.14	5 (9%)	54,89,89	2.19	5 (9%)
4	GOL	B	503	-	5,5,5	0.11	0	5,5,5	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/28/50/50	0/6/6/6
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
2	FAD	B	501	-	-	0/28/50/50	0/6/6/6
4	GOL	B	503	-	-	0/4/4/4	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	C4-C4X	2.20	1.45	1.41
2	A	501	FAD	C9A-N10	2.30	1.41	1.38
2	A	501	FAD	C5X-N5	2.79	1.39	1.35
2	B	501	FAD	C5X-N5	2.85	1.39	1.35
2	B	501	FAD	C4X-C10	2.85	1.46	1.41
2	B	501	FAD	C4-C4X	2.90	1.46	1.41
2	A	501	FAD	C4X-C10	3.11	1.46	1.41
2	B	501	FAD	C4-N3	3.23	1.38	1.33
2	A	501	FAD	C4-N3	3.43	1.39	1.33
2	B	501	FAD	C9A-N10	4.12	1.44	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C4X-C4-N3	-6.47	114.27	123.48
2	B	501	FAD	C4X-C4-N3	-6.23	114.61	123.48
2	B	501	FAD	C4X-C10-N10	-5.41	116.76	120.52
2	A	501	FAD	C4X-C10-N10	-5.11	116.97	120.52
2	B	501	FAD	C4-C4X-C10	-3.15	117.42	119.96
2	A	501	FAD	C4-C4X-C10	-2.82	117.68	119.96
2	A	501	FAD	C1'-N10-C10	2.45	121.01	118.50
2	A	501	FAD	C10-C4X-N5	3.04	124.09	120.59
2	B	501	FAD	C10-C4X-N5	3.30	124.39	120.59
2	A	501	FAD	C4-N3-C2	12.18	125.81	115.16
2	B	501	FAD	C4-N3-C2	12.22	125.85	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/461 (97%)	-0.02	15 (3%) 47 56	14, 25, 61, 79	0
1	B	451/461 (97%)	0.15	27 (5%) 23 30	15, 31, 67, 95	0
All	All	899/922 (97%)	0.07	42 (4%) 32 41	14, 28, 65, 95	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	245	ALA	6.6
1	B	246	ALA	6.2
1	B	100	ARG	5.8
1	A	379	SER	5.0
1	B	247	GLN	4.8
1	B	377	VAL	4.7
1	B	245	ALA	4.7
1	B	378	ALA	4.2
1	A	49	ARG	4.1
1	A	90	GLY	3.9
1	B	382	TYR	3.9
1	B	257	VAL	3.6
1	B	379	SER	3.6
1	A	382	TYR	3.6
1	B	376	LYS	3.6
1	A	247	GLN	3.5
1	B	383	LEU	3.4
1	B	250	SER	3.2
1	B	101	ASP	3.2
1	B	150	VAL	3.2
1	B	102	ASP	3.1
1	A	249	ALA	3.0
1	B	249	ALA	2.9
1	B	103	SER	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	244	PRO	2.6
1	A	438	LEU	2.6
1	A	383	LEU	2.6
1	B	438	LEU	2.6
1	A	246	ALA	2.5
1	A	50	GLY	2.5
1	A	380	PRO	2.5
1	B	53	ILE	2.5
1	B	49	ARG	2.4
1	A	443	LEU	2.4
1	B	7	ALA	2.3
1	A	91	THR	2.3
1	B	287	HIS	2.2
1	B	51	ARG	2.2
1	B	448	HIS	2.1
1	B	386	ARG	2.1
1	B	248	PRO	2.1
1	B	394	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	A	503	6/6	0.90	0.16	2.00	33,38,39,40	0
4	GOL	B	503	6/6	0.89	0.14	1.01	29,35,36,38	0
3	CL	B	502	1/1	0.99	0.09	-0.12	24,24,24,24	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FAD	A	501	53/53	0.98	0.08	-0.68	12,15,19,21	0
2	FAD	B	501	53/53	0.98	0.07	-0.86	14,19,23,24	0
3	CL	A	502	1/1	0.99	0.06	-3.55	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.