



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 27, 2017 – 12:35 AM EDT

PDB ID : 5NCZ
Title : mPI3Kd IN COMPLEX WITH inh1
Authors : Petersen, J.
Deposited on : 2017-03-06
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3233 (1.96-1.92)
Clashscore	112137	3430 (1.96-1.92)
Ramachandran outliers	110173	3395 (1.96-1.92)
Sidechain outliers	110143	3395 (1.96-1.92)
RSRZ outliers	101464	3250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	936	<div> <div>7%</div> <div>81%</div> <div>7%</div> <div>12%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	827	6687	4287	1134	1211	55	3	3	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	497	ILE	HIS	conflict	UNP O35904
A	?	-	LEU	deletion	UNP O35904
A	?	-	GLN	deletion	UNP O35904

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 4-azanyl-6-[[[(1 {S})-1-[6-[3-[(dimethylamino)methyl]phenyl]-3-methyl-5-oxidanylidene-[1,3]thiazolo[3,2-a]pyridin-7-yl]ethyl]amino]pyrimidine-5-carbonitrile (three-letter code: 8TN) (formula: C₂₄H₂₅N₇OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			33	24	7	1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	257	Total	O	0	0
			257	257		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.65Å 64.84Å 116.60Å 90.00° 103.63° 90.00°	Depositor
Resolution (Å)	42.97 – 1.94 42.97 – 1.94	Depositor EDS
% Data completeness (in resolution range)	96.9 (42.97-1.94) 96.9 (42.97-1.94)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 1.94Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.225 , 0.259 0.232 , 0.267	Depositor DCC
R_{free} test set	3715 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	30.4	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6993	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 8TN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/6838	0.60	0/9223

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6687	0	6671	28	0
2	A	16	0	24	1	0
3	A	33	0	0	0	0
4	A	257	0	0	0	0
All	All	6993	0	6695	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:GLY:HA3	1:A:768:ALA:HB2	1.50	0.93
1:A:328:ILE:HB	1:A:472:VAL:HG23	1.74	0.68
1:A:549:LEU:HG	1:A:564:MET:CE	2.28	0.64
1:A:549:LEU:HG	1:A:564:MET:HE3	1.83	0.60
1:A:642:LYS:HG2	2:A:1103:EDO:H21	1.85	0.58
1:A:784:LEU:HD12	1:A:823:GLY:HA3	1.84	0.58
1:A:512:ARG:HG2	1:A:530:LEU:HD12	1.86	0.57
1:A:194:VAL:HG21	1:A:216:LEU:HD21	1.87	0.56
1:A:271:THR:O	1:A:273:HIS:HD2	1.90	0.55
1:A:331:ARG:HD2	1:A:470:ALA:HB3	1.89	0.53
1:A:859:LEU:HD21	1:A:905:GLY:HA2	1.90	0.53
1:A:247:HIS:CD2	1:A:740:LEU:HD21	2.45	0.52
1:A:515:LEU:HD12	1:A:516:GLU:HG3	1.95	0.48
1:A:329:GLU:HB2	1:A:369:PRO:O	2.15	0.46
1:A:549:LEU:HG	1:A:564:MET:HE1	1.98	0.45
1:A:213:PRO:O	1:A:217:MET:HG3	2.17	0.45
1:A:192:VAL:HG11	1:A:216:LEU:HD11	1.99	0.44
1:A:902:ARG:HD3	1:A:906:GLN:HB2	1.98	0.44
1:A:971:GLY:HA3	1:A:1004:LEU:HD21	2.00	0.44
1:A:324:SER:HB3	1:A:376:GLU:HG3	1.99	0.44
1:A:157:LEU:HD22	1:A:161:GLU:HB3	2.00	0.43
1:A:329:GLU:HG2	1:A:472:VAL:HG22	2.01	0.43
1:A:209:THR:HB	1:A:257:CYS:HB3	2.00	0.42
1:A:187:ASN:N	1:A:210:LYS:HZ2	2.18	0.42
1:A:637:ALA:HB1	1:A:644:GLY:HA2	2.02	0.41
1:A:617:GLN:HE21	1:A:984:ALA:HA	1.85	0.41
1:A:859:LEU:HA	1:A:859:LEU:HD23	1.94	0.40
1:A:342:VAL:HG22	1:A:362:GLU:HG2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	806/936 (86%)	787 (98%)	17 (2%)	2 (0%)	51 40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	319	LEU
1	A	328	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	735/824 (89%)	705 (96%)	30 (4%)	35 20

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	LEU
1	A	203	PHE
1	A	316	LEU
1	A	317	TRP
1	A	331	ARG
1	A	332	LYS
1	A	333	VAL
1	A	340	LYS
1	A	356	LYS
1	A	374	ARG
1	A	423	LEU
1	A	437	ARG
1	A	453	LEU
1	A	507	THR
1	A	511	LEU
1	A	512	ARG
1	A	530	LEU
1	A	553	THR
1	A	634	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	731	LEU
1	A	743	GLU
1	A	755	LYS
1	A	757	LYS
1	A	766	GLU
1	A	841	LYS
1	A	855	LEU
1	A	898	ASN
1	A	907	LEU
1	A	915	PHE
1	A	1004	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	247	HIS
1	A	273	HIS
1	A	278	HIS
1	A	291	ASN
1	A	344	GLN
1	A	349	HIS
1	A	610	GLN
1	A	617	GLN
1	A	696	ASN
1	A	710	GLN
1	A	898	ASN
1	A	943	GLN
1	A	976	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1101	-	3,3,3	0.17	0	2,2,2	0.13	0
2	EDO	A	1102	-	3,3,3	0.25	0	2,2,2	0.25	0
2	EDO	A	1103	-	3,3,3	0.16	0	2,2,2	0.14	0
2	EDO	A	1104	-	3,3,3	0.23	0	2,2,2	0.13	0
3	8TN	A	1105	-	33,36,36	1.84	9 (27%)	31,52,52	1.82	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1101	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1102	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1103	-	-	0/1/1/1	0/0/0/0
2	EDO	A	1104	-	-	0/1/1/1	0/0/0/0
3	8TN	A	1105	-	-	0/16/18/18	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1105	8TN	C12-C17	-4.86	1.33	1.40
3	A	1105	8TN	C7-C10	-3.06	1.38	1.42
3	A	1105	8TN	C14-C8	-2.67	1.46	1.49
3	A	1105	8TN	C14-C15	-2.33	1.36	1.40
3	A	1105	8TN	C7-C11	-2.33	1.39	1.42
3	A	1105	8TN	C18-C14	2.14	1.48	1.44
3	A	1105	8TN	C10-N29	2.35	1.40	1.34
3	A	1105	8TN	C11-N30	3.52	1.40	1.35
3	A	1105	8TN	C12-C15	5.60	1.45	1.36

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1105	8TN	C14-C18-N28	-5.39	115.71	119.47
3	A	1105	8TN	C19-C16-C13	-4.24	122.73	129.90
3	A	1105	8TN	C20-C24-C15	-3.41	106.62	111.63
3	A	1105	8TN	C11-C7-C1	2.16	122.44	119.32
3	A	1105	8TN	N30-C11-N27	2.46	121.77	117.87
3	A	1105	8TN	C12-C15-C14	2.74	121.11	118.04
3	A	1105	8TN	C8-C14-C15	3.39	123.47	119.69

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1103	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	827/936 (88%)	0.51	64 (7%) 14 20	14, 34, 69, 112	4 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	317	TRP	7.1
1	A	228	PHE	6.4
1	A	334	ASN	6.1
1	A	416	CYS	5.6
1	A	1027	TRP	4.9
1	A	397	VAL	4.6
1	A	445	VAL	4.6
1	A	203	PHE	4.4
1	A	226	THR	4.2
1	A	517	ARG	3.9
1	A	395	TYR	3.9
1	A	1017	VAL	3.8
1	A	1024	ARG	3.7
1	A	227	VAL	3.7
1	A	333	VAL	3.6
1	A	270	LEU	3.6
1	A	488	LEU	3.4
1	A	205	PHE	3.4
1	A	496	ARG	3.4
1	A	363	VAL	3.4
1	A	515	LEU	3.3
1	A	474	TYR	3.2
1	A	330	GLY	3.2
1	A	930	VAL	3.1
1	A	396	ALA	3.1
1	A	934	LEU	3.1
1	A	514	ILE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	398	VAL	3.0
1	A	1014	HIS	2.9
1	A	394	LEU	2.9
1	A	483	VAL	2.9
1	A	479	ALA	2.9
1	A	377	PHE	2.8
1	A	323	PHE	2.8
1	A	936	TYR	2.8
1	A	471	LEU	2.8
1	A	316	LEU	2.7
1	A	492	LEU	2.7
1	A	507	THR	2.7
1	A	497	ILE	2.7
1	A	919	PHE	2.7
1	A	847	THR	2.6
1	A	993	LYS	2.6
1	A	341	LEU	2.6
1	A	472	VAL	2.6
1	A	191	LEU	2.5
1	A	418	ILE	2.5
1	A	235	GLN	2.4
1	A	512	ARG	2.4
1	A	523	LEU	2.3
1	A	342	VAL	2.3
1	A	201	GLU	2.3
1	A	417	PRO	2.3
1	A	225	ALA	2.2
1	A	360	SER	2.2
1	A	319	LEU	2.2
1	A	745[A]	CYS	2.2
1	A	710	GLN	2.1
1	A	318	SER	2.1
1	A	493	GLU	2.1
1	A	229	ARG	2.1
1	A	913	GLY	2.1
1	A	200	GLU	2.0
1	A	393	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	EDO	A	1101	4/4	0.96	0.14	0.32	23,28,29,30	0
2	EDO	A	1104	4/4	0.95	0.13	0.21	21,23,24,24	0
3	8TN	A	1105	33/33	0.96	0.11	-0.79	14,19,34,35	0
2	EDO	A	1103	4/4	0.97	0.09	-0.88	22,25,28,30	0
2	EDO	A	1102	4/4	0.95	0.07	-2.98	26,30,32,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.