



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 12:35 am GMT

PDB ID : 1ND5  
Title : Crystal Structures of Human Prostatic Acid Phosphatase in Complex with a Phosphate Ion and alpha-Benzylaminobenzylphosphonic Acid Update the Mechanistic Picture and Offer New Insights into Inhibitor Design  
Authors : Ortlund, E.; LaCount, M.W.; Lebioda, L.  
Deposited on : 2002-12-07  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	354	
1	B	354	
1	C	354	
1	D	354	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	5001	X	-	-	-
2	NAG	A	5002	X	-	-	-
2	NAG	B	5005	X	-	X	-
2	NAG	C	5007	X	-	X	-
2	NAG	C	5008	X	-	-	-
3	NDG	B	5004	-	-	X	-
4	NAG	D	5009	X	-	-	-
5	NAG	D	5010	X	-	-	-

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11727 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called prostatic acid phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			
1	B	342	Total	C	N	O	S	0	0	0
			2796	1802	461	517	16			
1	C	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			
1	D	342	Total	C	N	O	S	0	0	0
			2800	1807	461	516	16			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			30	16	2	12		
2	A	2	Total	C	N	O	0	0
			30	16	2	12		
2	B	2	Total	C	N	O	0	0
			30	16	2	12		
2	C	2	Total	C	N	O	0	0
			30	16	2	12		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NDG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	3	Total	C	N	O	0	0
			42	22	2	18		

- Molecule 6 is ALPHA-BENZYL-AMINOBENZYL-PHOSPHONIC ACID (three-letter code: 2BF) (formula:  $C_{14}H_{16}NO_3P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			19	14	1	3	1		
6	B	1	Total	C	N	O	P	0	0
			19	14	1	3	1		
6	C	1	Total	C	N	O	P	0	0
			19	14	1	3	1		
6	D	1	Total	C	N	O	P	0	0
			19	14	1	3	1		

- Molecule 7 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			16	10	6		
7	B	1	Total	C	O	0	0
			16	10	6		
7	C	1	Total	C	O	0	0
			16	10	6		
7	D	1	Total	C	O	0	0
			16	10	6		
7	A	1	Total	C	O	0	0
			16	10	6		
7	B	1	Total	C	O	0	0
			16	10	6		
7	C	1	Total	C	O	0	0
			16	10	6		
7	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	47	Total	O	0	0
			47	47		
8	B	35	Total	O	0	0
			35	35		
8	C	22	Total	O	0	0
			22	22		
8	D	31	Total	O	0	0
			31	31		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: prostatic acid phosphatase



- Molecule 1: prostatic acid phosphatase









• Molecule 1: prostatic acid phosphatase

Chain C: 42% 49% 6%



• Molecule 1: prostatic acid phosphatase

Chain D: 43% 47% 6%



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.10Å 204.86Å 71.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.94 – 2.90	Depositor
% Data completeness (in resolution range)	78.9 (77.94-2.90)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.205 , 0.279	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11727	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2BF, MAN, NAG, NDG, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2882	0.64	0/3914
1	B	0.39	0/2876	0.64	0/3905
1	C	0.39	0/2882	0.64	0/3914
1	D	0.39	0/2882	0.62	0/3914
All	All	0.39	0/11522	0.64	0/15647

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	2	0
2	B	1	0
2	C	2	0
5	D	1	0
All	All	6	0

There are no bond length outliers.

There are no bond angle outliers.

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	5001	NAG	C1
2	A	5002	NAG	C1
2	B	5005	NAG	C1
2	C	5007	NAG	C1
2	C	5008	NAG	C1

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2749	200	0
1	B	2796	0	2745	180	0
1	C	2800	0	2747	196	0
1	D	2800	0	2746	195	0
2	A	60	0	55	8	0
2	B	30	0	28	9	0
2	C	30	0	27	8	0
3	B	15	0	15	10	0
4	D	15	0	14	6	0
5	D	42	0	38	4	0
6	A	19	0	14	3	0
6	B	19	0	14	2	0
6	C	19	0	14	2	0
6	D	19	0	14	0	0
7	A	48	0	66	3	0
7	B	32	0	44	5	0
7	C	32	0	44	2	0
7	D	16	0	22	1	0
8	A	47	0	0	4	0
8	B	35	0	0	4	0
8	C	22	0	0	1	0
8	D	31	0	0	8	0
All	All	11727	0	11396	767	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 767 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:D:5010:NAG:H61	5:D:5011:NAG:H82	1.37	1.07
1:B:1145:LYS:HA	1:B:1145:LYS:HE3	1.32	1.05
1:B:1186:ASN:HD21	3:B:5004:NDG:H1	1.22	1.03
1:B:1299:ASN:ND2	2:B:5005:NAG:H1	1.74	1.02
1:B:1299:ASN:HD21	2:B:5005:NAG:H1	1.23	0.98

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/354 (96%)	288 (85%)	39 (12%)	13 (4%)	4	15
1	B	338/354 (96%)	284 (84%)	43 (13%)	11 (3%)	4	18
1	C	340/354 (96%)	282 (83%)	41 (12%)	17 (5%)	2	8
1	D	340/354 (96%)	287 (84%)	37 (11%)	16 (5%)	3	10
All	All	1358/1416 (96%)	1141 (84%)	160 (12%)	57 (4%)	3	12

5 of 57 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	65	TYR
1	B	1064	TYR
1	B	1245	SER
1	C	2099	TRP

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/325 (97%)	293 (93%)	21 (7%)	19	48
1	B	314/325 (97%)	297 (95%)	17 (5%)	26	59
1	C	314/325 (97%)	293 (93%)	21 (7%)	19	48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	314/325 (97%)	295 (94%)	19 (6%)	22	53
All	All	1256/1300 (97%)	1178 (94%)	78 (6%)	21	52

5 of 78 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1241	THR
1	C	2072	ARG
1	D	3153	ASP
1	B	1281	LEU
1	C	2036	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1185	HIS
1	C	2061	ASN
1	D	3226	GLN
1	B	1186	ASN
1	B	1231	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

11 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	5000	1,2	15,15,15	0.52	0	21,21,21	0.90	1 (4%)
2	NAG	A	5001	2	15,15,15	0.43	0	21,21,21	0.50	0
2	NAG	A	5002	1,2	15,15,15	0.42	0	21,21,21	0.67	0
2	NAG	A	5003	2	15,15,15	0.44	0	21,21,21	1.06	2 (9%)
2	NAG	B	5005	2	15,15,15	0.42	0	21,21,21	0.58	0
2	NAG	B	5006	2	15,15,15	0.49	0	21,21,21	0.99	1 (4%)
2	NAG	C	5007	1,2	15,15,15	0.74	0	21,21,21	1.04	1 (4%)
2	NAG	C	5008	2	15,15,15	0.44	0	21,21,21	0.88	1 (4%)
5	NAG	D	5010	1,5	15,15,15	0.54	0	21,21,21	0.56	0
5	NAG	D	5011	5	15,15,15	0.46	0	21,21,21	1.04	1 (4%)
5	MAN	D	5012	5	12,12,12	0.60	0	17,17,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	5000	1,2	-	0/6/26/26	0/1/1/1
2	NAG	A	5001	2	1/1/6/7	0/6/26/26	0/1/1/1
2	NAG	A	5002	1,2	1/1/6/7	0/6/26/26	0/1/1/1
2	NAG	A	5003	2	-	0/6/26/26	0/1/1/1
2	NAG	B	5005	2	1/1/6/7	0/6/26/26	0/1/1/1
2	NAG	B	5006	2	-	0/6/26/26	0/1/1/1
2	NAG	C	5007	1,2	1/1/6/7	0/6/26/26	0/1/1/1
2	NAG	C	5008	2	1/1/6/7	0/6/26/26	0/1/1/1
5	NAG	D	5010	1,5	1/1/6/7	0/6/26/26	0/1/1/1
5	NAG	D	5011	5	-	0/6/26/26	0/1/1/1
5	MAN	D	5012	5	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	5003	NAG	O1-C1-C2	-3.52	101.91	109.22
5	D	5011	NAG	O1-C1-C2	-3.49	101.96	109.22
2	C	5007	NAG	C1-C2-C3	-3.30	106.05	110.54
2	B	5006	NAG	O1-C1-C2	-3.28	102.41	109.22
2	A	5000	NAG	O1-C1-C2	-2.92	103.16	109.22

5 of 6 chirality outliers are listed below:



Mol	Chain	Res	Type	Atom
2	C	5007	NAG	C1
2	B	5005	NAG	C1
2	A	5001	NAG	C1
2	C	5008	NAG	C1
2	A	5002	NAG	C1

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	5000	NAG	4	0
2	A	5001	NAG	2	0
2	A	5002	NAG	4	0
2	B	5005	NAG	9	0
2	C	5007	NAG	7	0
2	C	5008	NAG	4	0
5	D	5010	NAG	4	0
5	D	5011	NAG	3	0

## 5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	2BF	A	6000	-	20,20,20	4.08	15 (75%)	22,27,27	1.32	3 (13%)
7	1PE	A	7000	-	15,15,15	0.90	0	14,14,14	1.47	4 (28%)
7	1PE	A	7004	-	15,15,15	0.80	0	14,14,14	1.50	4 (28%)
7	1PE	A	7007	-	15,15,15	0.96	0	14,14,14	1.48	4 (28%)
3	NDG	B	5004	-	15,15,15	0.30	0	21,21,21	0.63	0
6	2BF	B	6001	-	20,20,20	3.90	16 (80%)	22,27,27	1.34	3 (13%)
7	1PE	B	7001	-	15,15,15	0.72	0	14,14,14	1.43	4 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	1PE	B	7005	-	15,15,15	0.84	0	14,14,14	1.49	4 (28%)
6	2BF	C	6002	-	20,20,20	3.96	15 (75%)	22,27,27	1.38	4 (18%)
7	1PE	C	7002	-	15,15,15	0.86	0	14,14,14	1.49	4 (28%)
7	1PE	C	7006	-	15,15,15	0.88	0	14,14,14	1.47	4 (28%)
4	NAG	D	5009	1	15,15,15	0.42	0	21,21,21	0.61	0
6	2BF	D	6003	-	20,20,20	4.09	16 (80%)	22,27,27	1.27	3 (13%)
7	1PE	D	7003	-	15,15,15	0.85	0	14,14,14	1.43	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	2BF	A	6000	-	-	0/15/15/15	0/2/2/2
7	1PE	A	7000	-	-	0/13/13/13	0/0/0/0
7	1PE	A	7004	-	-	0/13/13/13	0/0/0/0
7	1PE	A	7007	-	-	0/13/13/13	0/0/0/0
3	NDG	B	5004	-	-	0/6/26/26	0/1/1/1
6	2BF	B	6001	-	-	0/15/15/15	0/2/2/2
7	1PE	B	7001	-	-	0/13/13/13	0/0/0/0
7	1PE	B	7005	-	-	0/13/13/13	0/0/0/0
6	2BF	C	6002	-	-	0/15/15/15	0/2/2/2
7	1PE	C	7002	-	-	0/13/13/13	0/0/0/0
7	1PE	C	7006	-	-	0/13/13/13	0/0/0/0
4	NAG	D	5009	1	1/1/6/7	0/6/26/26	0/1/1/1
6	2BF	D	6003	-	-	0/15/15/15	0/2/2/2
7	1PE	D	7003	-	-	0/13/13/13	0/0/0/0

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	6001	2BF	P-C7	2.03	1.86	1.83
6	B	6001	2BF	C10-C9	2.03	1.42	1.38
6	D	6003	2BF	C10-C9	2.04	1.42	1.38
6	D	6003	2BF	C2-C1	2.12	1.43	1.38
6	C	6002	2BF	P-C7	2.23	1.86	1.83

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	6001	2BF	O1P-P-C7	-4.07	104.03	111.04
6	C	6002	2BF	O1P-P-C7	-3.76	104.56	111.04
6	A	6000	2BF	O1P-P-C7	-3.26	105.42	111.04
6	D	6003	2BF	C8-C19-N	-2.75	105.73	112.69
6	D	6003	2BF	C6-C5-C7	-2.66	117.89	120.72

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	5009	NAG	C1

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 34 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	6000	2BF	3	0
7	A	7000	1PE	1	0
7	A	7004	1PE	2	0
3	B	5004	NDG	10	0
6	B	6001	2BF	2	0
7	B	7001	1PE	1	0
7	B	7005	1PE	4	0
6	C	6002	2BF	2	0
7	C	7002	1PE	1	0
7	C	7006	1PE	1	0
4	D	5009	NAG	6	0
7	D	7003	1PE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.