



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:19 am GMT

PDB ID : 1NDL
Title : THE AWD NUCLEOTIDE DIPHOSPHATE KINASE FROM DROSOPHILA
Authors : Janin, J.; Chiadmi, M.; Dumas, C.; Lascu, I.; Lebras, G.; Morera, S.; Veron, M.
Deposited on : 1993-11-27
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	153	
1	B	153	
1	C	153	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NUCLEOSIDE DIPHOSPHATE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	152	Total	C	N	O	S	0	0	0
			1203	777	205	216	5			
1	B	152	Total	C	N	O	S	0	0	0
			1203	777	205	216	5			
1	C	152	Total	C	N	O	S	0	0	0
			1203	777	205	216	5			

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	116	Total	O	0	0
			116	116		
2	B	94	Total	O	0	0
			94	94		
2	C	105	Total	O	0	0
			105	105		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: NUCLEOSIDE DIPHOSPHATE KINASE



• Molecule 1: NUCLEOSIDE DIPHOSPHATE KINASE



• Molecule 1: NUCLEOSIDE DIPHOSPHATE KINASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	115.65Å 115.65Å 98.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.40)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.167 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3924	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/1233	1.64	17/1668 (1.0%)
1	B	0.79	0/1233	1.62	20/1668 (1.2%)
1	C	0.81	0/1233	1.80	15/1668 (0.9%)
All	All	0.80	0/3699	1.69	52/5004 (1.0%)

There are no bond length outliers.

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	28	ARG	CD-NE-CZ	36.93	175.31	123.60
1	A	106	ARG	CD-NE-CZ	22.95	155.73	123.60
1	A	106	ARG	NE-CZ-NH1	15.29	127.94	120.30
1	C	115	ARG	NE-CZ-NH1	12.90	126.75	120.30
1	B	28	ARG	NE-CZ-NH2	-10.42	115.09	120.30
1	C	106	ARG	NE-CZ-NH1	9.94	125.27	120.30
1	B	106	ARG	CD-NE-CZ	9.56	136.98	123.60
1	C	122	ASP	N-CA-CB	-9.29	93.87	110.60
1	B	122	ASP	N-CA-CB	-8.94	94.50	110.60
1	A	115	ARG	NE-CZ-NH2	-8.24	116.18	120.30
1	B	19	ARG	NE-CZ-NH1	-8.24	116.18	120.30
1	C	122	ASP	CB-CG-OD1	-8.20	110.92	118.30
1	B	121	SER	C-N-CA	8.18	142.16	121.70
1	B	19	ARG	NE-CZ-NH2	8.02	124.31	120.30
1	B	106	ARG	NE-CZ-NH2	-7.99	116.30	120.30
1	B	29	PHE	CB-CG-CD2	-7.50	115.55	120.80
1	C	59	ARG	NE-CZ-NH1	7.15	123.87	120.30
1	A	68	TYR	CB-CG-CD1	-6.97	116.82	121.00
1	A	122	ASP	CB-CA-C	-6.62	97.17	110.40
1	C	121	SER	C-N-CA	6.61	138.22	121.70
1	C	92	LEU	CA-CB-CG	6.49	130.24	115.30
1	B	115	ARG	NE-CZ-NH1	6.41	123.50	120.30
1	B	153	GLU	CA-CB-CG	6.36	127.39	113.40
1	B	121	SER	CA-C-O	6.34	133.42	120.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	31	GLN	CB-CA-C	-6.29	97.83	110.40
1	B	31	GLN	CB-CA-C	-6.24	97.92	110.40
1	A	116	ASN	C-N-CA	6.22	137.26	121.70
1	C	149	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	55	ASP	CB-CG-OD2	6.00	123.70	118.30
1	B	122	ASP	CB-CG-OD1	-5.99	112.91	118.30
1	B	122	ASP	CB-CG-OD2	-5.98	112.92	118.30
1	B	28	ARG	NH1-CZ-NH2	5.95	125.95	119.40
1	B	15	ASP	CB-CG-OD1	5.78	123.50	118.30
1	A	29	PHE	CB-CG-CD2	-5.76	116.76	120.80
1	B	122	ASP	OD1-CG-OD2	5.72	134.17	123.30
1	A	121	SER	C-N-CA	5.67	135.88	121.70
1	B	153	GLU	CG-CD-OE1	-5.65	107.00	118.30
1	A	68	TYR	CB-CG-CD2	5.61	124.36	121.00
1	C	115	ARG	CD-NE-CZ	5.59	131.43	123.60
1	C	55	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	106	ARG	NH1-CZ-NH2	-5.41	113.45	119.40
1	A	92	LEU	CA-CB-CG	5.38	127.69	115.30
1	A	43	TRP	CA-C-O	-5.37	108.82	120.10
1	A	92	LEU	CB-CG-CD2	5.34	120.08	111.00
1	A	31	GLN	CA-CB-CG	5.21	124.87	113.40
1	C	121	SER	CA-C-O	5.18	130.98	120.10
1	B	109	PHE	N-CA-C	5.14	124.89	111.00
1	A	29	PHE	CB-CG-CD1	5.12	124.39	120.80
1	C	121	SER	N-CA-CB	5.11	118.17	110.50
1	B	104	THR	CA-CB-OG1	-5.07	98.36	109.00
1	A	75	VAL	CA-CB-CG2	5.05	118.48	110.90
1	C	31	GLN	CA-CB-CG	5.04	124.49	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1203	0	1207	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1203	0	1206	18	0
1	C	1203	0	1206	19	0
2	A	116	0	0	3	0
2	B	94	0	0	4	0
2	C	105	0	0	2	0
All	All	3924	0	3619	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:122:ASP:HB2	2:B:167:HOH:O	1.69	0.91
1:C:4:ASN:HD22	1:C:82:LEU:HB2	1.43	0.82
1:B:4:ASN:HD22	1:B:82:LEU:HB2	1.53	0.72
1:B:8:THR:HG23	1:B:10:ILE:HD13	1.75	0.69
1:C:4:ASN:ND2	1:C:82:LEU:HB2	2.06	0.68
1:A:98:ALA:HB2	2:A:254:HOH:O	1.93	0.67
1:B:62:PHE:N	1:B:63:PRO:HD2	2.12	0.63
1:A:40:LYS:O	1:A:76:PRO:HD2	1.99	0.62
1:C:40:LYS:O	1:C:76:PRO:HD2	1.98	0.62
1:B:112:GLN:HE22	1:C:153:GLU:HB3	1.67	0.59
1:A:62:PHE:HB3	1:A:63:PRO:HD3	1.84	0.58
1:B:122:ASP:HB3	1:B:126:SER:OG	2.04	0.58
1:A:153:GLU:HG2	1:C:114:GLY:HA3	1.87	0.56
1:A:6:GLU:HA	2:A:234:HOH:O	2.03	0.56
1:C:8:THR:HG23	1:C:10:ILE:HD12	1.87	0.56
1:C:122:ASP:HB2	2:C:169:HOH:O	2.05	0.56
1:A:127:ALA:O	1:A:131:ILE:HG13	2.04	0.56
1:C:62:PHE:HB3	1:C:63:PRO:HD3	1.88	0.54
1:C:122:ASP:HB3	1:C:126:SER:OG	2.08	0.54
1:A:92:LEU:HD22	1:A:118:ILE:HD13	1.90	0.52
1:C:27:GLU:O	1:C:31:GLN:HG3	2.09	0.52
1:B:48:LEU:HD11	1:B:133:LEU:HG	1.92	0.50
1:A:13:LYS:HB3	1:A:14:PRO:HD2	1.92	0.50
1:B:106:ARG:NH2	2:B:188:HOH:O	2.45	0.49
1:A:106:ARG:HD3	1:A:116:ASN:HB2	1.95	0.49
1:A:62:PHE:HB3	1:A:63:PRO:CD	2.44	0.48
1:C:4:ASN:ND2	1:C:82:LEU:HD13	2.29	0.48
1:B:125:GLU:O	1:B:129:LYS:HD3	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:40:LYS:HE2	1:C:135:PHE:CE1	2.50	0.47
1:C:106:ARG:NH2	2:C:190:HOH:O	2.47	0.47
1:A:13:LYS:HB3	1:A:14:PRO:CD	2.44	0.46
1:C:45:SER:HB2	1:C:47:GLU:OE1	2.16	0.46
1:A:47:GLU:HG2	1:A:51:LYS:HE3	1.99	0.45
1:B:4:ASN:ND2	1:B:82:LEU:HB2	2.26	0.45
1:A:2:ALA:HB3	1:A:5:LYS:HG3	1.99	0.44
1:A:27:GLU:O	1:A:31:GLN:HB2	2.18	0.44
1:B:40:LYS:O	1:B:76:PRO:HD2	2.19	0.43
1:A:14:PRO:HA	2:A:257:HOH:O	2.18	0.43
1:C:13:LYS:HB3	1:C:14:PRO:CD	2.48	0.42
1:B:32:LYS:HG2	1:B:34:PHE:CE2	2.54	0.42
1:C:48:LEU:CD1	1:C:133:LEU:HD22	2.49	0.42
1:B:106:ARG:HD3	1:B:116:ASN:HB2	2.00	0.42
1:B:106:ARG:NH1	1:B:118:ILE:O	2.53	0.42
1:A:28:ARG:HH11	1:A:28:ARG:HD2	1.64	0.41
1:B:142:THR:HA	2:B:224:HOH:O	2.20	0.41
1:B:62:PHE:N	1:B:63:PRO:CD	2.79	0.41
1:B:8:THR:HG23	1:B:10:ILE:CD1	2.46	0.41
2:B:244:HOH:O	1:C:91:MET:HG2	2.20	0.41
1:C:56:LEU:O	1:C:62:PHE:HB2	2.22	0.40
1:B:121:SER:OG	1:B:127:ALA:HA	2.21	0.40
1:A:31:GLN:HG2	1:C:109:PHE:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/153 (98%)	146 (97%)	3 (2%)	1 (1%)	25	37
1	B	150/153 (98%)	143 (95%)	6 (4%)	1 (1%)	25	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	150/153 (98%)	145 (97%)	4 (3%)	1 (1%)	25	37
All	All	450/459 (98%)	434 (96%)	13 (3%)	3 (1%)	25	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	117	ILE
1	B	117	ILE
1	C	117	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/127 (99%)	122 (97%)	4 (3%)	44	65
1	B	126/127 (99%)	125 (99%)	1 (1%)	85	93
1	C	126/127 (99%)	119 (94%)	7 (6%)	25	39
All	All	378/381 (99%)	366 (97%)	12 (3%)	44	65

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	MET
1	A	89	ARG
1	A	122	ASP
1	A	142	THR
1	B	128	GLU
1	C	28	ARG
1	C	47	GLU
1	C	83	ASN
1	C	126	SER
1	C	133	LEU
1	C	142	THR
1	C	153	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	83	ASN
1	A	90	GLN
1	A	136	ASN
1	B	4	ASN
1	B	112	GLN
1	C	4	ASN
1	C	83	ASN
1	C	136	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.