



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:51 am GMT

PDB ID : 1NE7
Title : HUMAN GLUCOSAMINE-6-PHOSPHATE DEAMINASE ISOMERASE AT
1.75 Å RESOLUTION COMPLEXED WITH N-ACETYL-GLUCOSAMINE
-6-PHOSPHATE AND 2-DEOXY-2-AMINO-GLUCITOL-6-PHOSPHATE
Authors : Arreola, R.; Valderrama, B.; Morante, M.L.; Horjales, E.
Deposited on : 2002-12-10
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1762 (1.76-1.76)
Clashscore	112137	1889 (1.76-1.76)
Ramachandran outliers	110173	1868 (1.76-1.76)
Sidechain outliers	110143	1868 (1.76-1.76)
RSRZ outliers	101464	1770 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>• •</div> </div> </div>
1	B	289	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>• 5%</div> </div> </div>
1	C	289	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>• 5%</div> </div> </div>
1	D	289	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>9%</div> <div>• 5%</div> </div> </div>
1	E	289	<div> <div>7%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>• 5%</div> </div> </div>
1	F	289	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>• 5%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	C	295	-	-	-	X
2	GLC	E	295	-	-	-	X
5	SO4	C	4297[B]	-	-	-	X
6	AGP	A	2298[A]	-	-	-	X
6	AGP	B	3298[A]	-	-	-	X
6	AGP	C	4298[A]	-	-	-	X
6	AGP	D	5298[A]	-	-	-	X
6	AGP	E	6298[A]	-	-	-	X
6	AGP	F	7298[A]	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 16013 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosamine-6-phosphate isomerase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	C	N	O	S	21	7	0
			2294	1480	382	419	13			
1	B	274	Total	C	N	O	S	5	3	0
			2203	1425	364	400	14			
1	C	274	Total	C	N	O	S	7	8	0
			2244	1451	372	406	15			
1	D	274	Total	C	N	O	S	4	7	0
			2231	1443	369	406	13			
1	E	274	Total	C	N	O	S	17	7	0
			2237	1446	372	405	14			
1	F	274	Total	C	N	O	S	9	6	0
			2227	1438	371	405	13			

- Molecule 2 is SUGAR (GLU-GLU) (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	C	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			11	6	5		
2	D	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			11	6	5		
2	E	1	Total	C	O	0	0
			12	6	6		
2	E	1	Total	C	O	0	0
			11	6	5		
2	F	1	Total	C	O	0	0
			12	6	6		
2	F	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE-6-PHOSPHATE) (three-letter code: 16G) (formula: C₈H₁₆NO₉P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			19	8	1	9	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
3	A	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
3	E	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
3	F	1	Total	C	N	O	P	0	0
			19	8	1	9	1		
3	D	1	Total	C	N	O	P	0	0
			19	8	1	9	1		

- Molecule 4 is a polymer of unknown type called SUGAR (GLU-GLU).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	2	Total	C	O	0	0
			23	12	11		

- Molecule 5 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	1
			5	4	1		
5	B	1	Total	O	S	0	1
			5	4	1		
5	C	1	Total	O	S	0	1
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	1
			5	4	1		
5	D	1	Total	O	S	0	1
			5	4	1		
5	E	1	Total	O	S	0	1
			5	4	1		
5	F	1	Total	O	S	0	1
			5	4	1		

- Molecule 6 is 2-DEOXY-2-AMINO GLUCITOL-6-PHOSPHATE (three-letter code: AGP) (formula: C₆H₁₆NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 16	C 6	N 1	O 8	P 1	3	1
6	B	1	Total 16	C 6	N 1	O 8	P 1	2	1
6	C	1	Total 16	C 6	N 1	O 8	P 1	2	1
6	D	1	Total 16	C 6	N 1	O 8	P 1	2	1
6	E	1	Total 16	C 6	N 1	O 8	P 1	2	1
6	F	1	Total 16	C 6	N 1	O 8	P 1	0	1


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	376	Total 376	O 376	0	0
7	B	364	Total 364	O 364	0	0
7	C	406	Total 406	O 406	0	0
7	D	366	Total 366	O 366	0	0
7	E	305	Total 305	O 305	0	0
7	F	377	Total 377	O 377	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glucosamine-6-phosphate isomerase




• Molecule 1: Glucosamine-6-phosphate isomerase




• Molecule 1: Glucosamine-6-phosphate isomerase





• Molecule 1: Glucosamine-6-phosphate isomerase







● Molecule 1: Glucosamine-6-phosphate isomerase



● Molecule 1: Glucosamine-6-phosphate isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	109.88Å 110.89Å 180.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 1.75 29.54 – 1.75	Depositor EDS
% Data completeness (in resolution range)	94.3 (19.97-1.75) 94.1 (29.54-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.86 (at 1.75Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.193 , 0.217 0.203 , 0.225	Depositor DCC
R_{free} test set	20802 reflections (9.95%)	DCC
Wilson B-factor (Å ²)	20.5	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16013	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, 16G, SO4, AGP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/2351	0.66	4/3178 (0.1%)
1	B	0.30	0/2259	0.58	0/3057
1	C	0.32	0/2300	0.60	0/3111
1	D	0.31	0/2288	0.58	0/3096
1	E	0.30	0/2293	0.57	0/3101
1	F	0.30	0/2283	0.57	0/3088
All	All	0.31	0/13774	0.60	4/18631 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	172[A]	ARG	NE-CZ-NH1	-8.57	116.01	120.30
1	A	172[B]	ARG	NE-CZ-NH1	-8.57	116.01	120.30
1	A	172[A]	ARG	NE-CZ-NH2	6.42	123.51	120.30
1	A	172[B]	ARG	NE-CZ-NH2	6.42	123.51	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2294	0	2284	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2203	0	2189	42	0
1	C	2244	0	2234	37	0
1	D	2231	0	2217	29	0
1	E	2237	0	2228	35	0
1	F	2227	0	2213	33	0
2	A	23	0	16	0	0
2	C	23	0	16	0	0
2	D	23	0	16	0	0
2	E	23	0	16	0	0
2	F	23	0	16	0	0
3	A	19	0	14	0	0
3	B	19	0	14	0	0
3	C	19	0	14	0	0
3	D	19	0	14	0	0
3	E	19	0	14	0	0
3	F	19	0	14	0	0
4	B	23	0	16	0	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
5	C	10	0	0	0	0
5	D	5	0	0	0	0
5	E	5	0	0	0	0
5	F	5	0	0	1	0
6	A	16	0	14	1	0
6	B	16	0	13	0	0
6	C	16	0	14	0	0
6	D	16	0	14	0	0
6	E	16	0	14	0	0
6	F	16	0	14	5	0
7	A	376	0	0	2	0
7	B	364	0	0	3	0
7	C	406	0	0	1	0
7	D	366	0	0	3	0
7	E	305	0	0	1	0
7	F	377	0	0	8	0
All	All	16013	0	13628	204	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 204 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:7298[A]:AGP:H61	7:F:7673:HOH:O	1.47	1.12
6:A:2298[A]:AGP:O3	7:A:4587:HOH:O	1.68	1.10
1:A:7:GLU:HB3	1:A:11:GLN:HE22	1.25	1.00
1:E:7:GLU:HB3	1:E:11:GLN:HE22	1.36	0.90
1:A:7:GLU:HB3	1:A:11:GLN:NE2	1.90	0.85

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/289 (99%)	276 (96%)	8 (3%)	2 (1%)	25	9
1	B	275/289 (95%)	266 (97%)	8 (3%)	1 (0%)	38	18
1	C	280/289 (97%)	273 (98%)	6 (2%)	1 (0%)	38	18
1	D	279/289 (96%)	272 (98%)	6 (2%)	1 (0%)	38	18
1	E	279/289 (96%)	269 (96%)	9 (3%)	1 (0%)	38	18
1	F	278/289 (96%)	270 (97%)	7 (2%)	1 (0%)	38	18
All	All	1677/1734 (97%)	1626 (97%)	44 (3%)	7 (0%)	38	18

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	THR
1	F	40	PRO
1	A	40	PRO
1	D	40	PRO
1	E	40	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/249 (100%)	242 (98%)	6 (2%)	54	30
1	B	237/249 (95%)	230 (97%)	7 (3%)	46	21
1	C	242/249 (97%)	236 (98%)	6 (2%)	53	28
1	D	241/249 (97%)	235 (98%)	6 (2%)	53	28
1	E	241/249 (97%)	235 (98%)	6 (2%)	53	28
1	F	240/249 (96%)	234 (98%)	6 (2%)	53	28
All	All	1449/1494 (97%)	1412 (97%)	37 (3%)	51	27

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	143	HIS
1	D	134	PHE
1	F	180	LYS
1	C	212	LEU
1	C	222	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	58	ASN
1	C	100	HIS
1	F	11	GLN
1	C	8	HIS
1	C	11	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GLC	B	294	4	12,12,12	3.18	5 (41%)	17,17,17	3.33	6 (35%)
4	GLC	B	295	4	11,11,12	3.21	5 (45%)	13,15,17	1.65	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	B	294	4	-	0/2/22/22	0/1/1/1
4	GLC	B	295	4	-	0/2/19/22	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	295	GLC	O2-C2	-8.36	1.25	1.43
4	B	294	GLC	O3-C3	-8.08	1.24	1.43
4	B	295	GLC	O6-C6	-4.31	1.24	1.42
4	B	294	GLC	O6-C6	-4.26	1.24	1.42
4	B	295	GLC	C4-C3	2.02	1.57	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	294	GLC	O2-C2-C1	-6.75	95.73	109.75
4	B	294	GLC	O1-C1-O5	-3.90	98.67	110.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	294	GLC	C1-C2-C3	-2.58	105.99	110.65
4	B	294	GLC	O6-C6-C5	2.36	119.30	111.34
4	B	295	GLC	O6-C6-C5	2.42	119.50	111.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry [i](#)

29 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	2296[B]	-	4,4,4	0.59	0	6,6,6	0.31	0
6	AGP	A	2298[A]	-	15,15,15	0.94	0	16,21,21	1.23	2 (12%)
2	GLC	A	294	2	12,12,12	3.19	6 (50%)	17,17,17	3.33	6 (35%)
2	GLC	A	295	2	11,11,12	3.27	5 (45%)	13,15,17	1.69	3 (23%)
3	16G	A	4299	-	19,19,19	2.05	5 (26%)	28,28,28	2.39	6 (21%)
3	16G	B	2299	-	19,19,19	2.09	6 (31%)	28,28,28	2.38	7 (25%)
5	SO4	B	3296[B]	-	4,4,4	0.58	0	6,6,6	0.30	0
6	AGP	B	3298[A]	-	15,15,15	1.09	0	16,21,21	1.62	2 (12%)
2	GLC	C	294	2	12,12,12	3.17	6 (50%)	17,17,17	3.36	6 (35%)
2	GLC	C	295	2	11,11,12	3.22	5 (45%)	13,15,17	1.65	2 (15%)
3	16G	C	3299	-	19,19,19	2.05	7 (36%)	28,28,28	2.37	5 (17%)
5	SO4	C	4296[B]	-	4,4,4	0.58	0	6,6,6	0.30	0
5	SO4	C	4297[B]	-	4,4,4	0.62	0	6,6,6	0.27	0
6	AGP	C	4298[A]	-	15,15,15	1.14	1 (6%)	16,21,21	2.74	3 (18%)
2	GLC	D	294	2	12,12,12	3.19	6 (50%)	17,17,17	3.37	6 (35%)
2	GLC	D	295	2	11,11,12	3.25	5 (45%)	13,15,17	1.71	3 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	D	5296[B]	-	4,4,4	0.60	0	6,6,6	0.30	0
6	AGP	D	5298[A]	-	15,15,15	0.99	0	16,21,21	1.63	2 (12%)
3	16G	D	7299	-	19,19,19	2.08	8 (42%)	28,28,28	2.32	7 (25%)
2	GLC	E	294	2	12,12,12	3.17	6 (50%)	17,17,17	3.36	6 (35%)
2	GLC	E	295	2	11,11,12	3.28	5 (45%)	13,15,17	1.74	3 (23%)
3	16G	E	5299	-	19,19,19	2.11	6 (31%)	28,28,28	2.42	6 (21%)
5	SO4	E	6296[B]	-	4,4,4	0.58	0	6,6,6	0.31	0
6	AGP	E	6298[A]	-	15,15,15	1.10	1 (6%)	16,21,21	3.93	3 (18%)
2	GLC	F	294	2	12,12,12	3.23	6 (50%)	17,17,17	3.36	6 (35%)
2	GLC	F	295	2	11,11,12	3.24	5 (45%)	13,15,17	1.67	2 (15%)
3	16G	F	6299	-	19,19,19	2.05	6 (31%)	28,28,28	2.41	6 (21%)
5	SO4	F	7296[B]	-	4,4,4	0.52	0	6,6,6	0.30	0
6	AGP	F	7298[A]	-	15,15,15	1.14	0	16,21,21	4.12	4 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SO4	A	2296[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	A	2298[A]	-	-	0/20/20/20	0/0/0/0
2	GLC	A	294	2	-	0/2/22/22	0/1/1/1
2	GLC	A	295	2	-	0/2/19/22	0/1/1/1
3	16G	A	4299	-	-	0/10/30/30	0/1/1/1
3	16G	B	2299	-	-	0/10/30/30	0/1/1/1
5	SO4	B	3296[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	B	3298[A]	-	-	0/20/20/20	0/0/0/0
2	GLC	C	294	2	-	0/2/22/22	0/1/1/1
2	GLC	C	295	2	-	0/2/19/22	0/1/1/1
3	16G	C	3299	-	-	0/10/30/30	0/1/1/1
5	SO4	C	4296[B]	-	-	0/0/0/0	0/0/0/0
5	SO4	C	4297[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	C	4298[A]	-	-	0/20/20/20	0/0/0/0
2	GLC	D	294	2	-	0/2/22/22	0/1/1/1
2	GLC	D	295	2	-	0/2/19/22	0/1/1/1
5	SO4	D	5296[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	D	5298[A]	-	-	0/20/20/20	0/0/0/0
3	16G	D	7299	-	-	0/10/30/30	0/1/1/1
2	GLC	E	294	2	-	0/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	295	2	-	0/2/19/22	0/1/1/1
3	16G	E	5299	-	-	0/10/30/30	0/1/1/1
5	SO4	E	6296[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	E	6298[A]	-	-	0/20/20/20	0/0/0/0
2	GLC	F	294	2	-	0/2/22/22	0/1/1/1
2	GLC	F	295	2	-	0/2/19/22	0/1/1/1
3	16G	F	6299	-	-	0/10/30/30	0/1/1/1
5	SO4	F	7296[B]	-	-	0/0/0/0	0/0/0/0
6	AGP	F	7298[A]	-	-	0/20/20/20	0/0/0/0

The worst 5 of 95 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	295	GLC	O2-C2	-8.60	1.24	1.43
2	E	295	GLC	O2-C2	-8.57	1.24	1.43
2	D	295	GLC	O2-C2	-8.52	1.24	1.43
2	F	295	GLC	O2-C2	-8.51	1.24	1.43
2	C	295	GLC	O2-C2	-8.47	1.24	1.43

The worst 5 of 96 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	6298[A]	AGP	O3-C3-C2	-14.89	81.34	110.09
6	F	7298[A]	AGP	O3-C3-C2	-12.75	85.47	110.09
6	C	4298[A]	AGP	O3-C3-C4	-7.19	92.27	109.46
2	C	294	GLC	O2-C2-C1	-6.99	95.23	109.75
2	E	294	GLC	O2-C2-C1	-6.97	95.26	109.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	2298[A]	AGP	1	0
5	F	7296[B]	SO4	1	0
6	F	7298[A]	AGP	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	281/289 (97%)	0.09	16 (5%) 24 31	13, 21, 40, 53	7 (2%)
1	B	274/289 (94%)	0.18	16 (5%) 24 30	13, 22, 38, 51	2 (0%)
1	C	274/289 (94%)	-0.08	4 (1%) 74 81	13, 18, 30, 50	3 (1%)
1	D	274/289 (94%)	0.10	10 (3%) 43 50	14, 21, 33, 52	3 (1%)
1	E	274/289 (94%)	0.24	21 (7%) 14 19	14, 24, 43, 53	5 (1%)
1	F	274/289 (94%)	0.17	8 (2%) 52 59	14, 22, 34, 51	3 (1%)
All	All	1651/1734 (95%)	0.12	75 (4%) 34 40	13, 21, 37, 53	23 (1%)

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	274	GLU	7.5
1	B	178	LEU	7.4
1	F	274	GLU	7.2
1	D	274	GLU	6.9
1	E	274	GLU	6.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GLC	B	295	11/12	0.91	0.10	0.89	22,23,25,26	11
4	GLC	B	294	12/12	0.84	0.14	-	23,26,28,28	12

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	AGP	C	4298[A]	16/16	0.78	0.28	4.99	41,44,46,47	16
2	GLC	E	295	11/12	0.82	0.14	3.44	28,30,32,33	11
6	AGP	D	5298[A]	16/16	0.90	0.20	3.28	33,38,40,40	16
6	AGP	A	2298[A]	16/16	0.73	0.38	3.01	13,24,32,32	16
6	AGP	E	6298[A]	16/16	0.90	0.21	2.70	37,38,41,42	16
2	GLC	C	295	11/12	0.90	0.12	2.59	21,22,23,26	11
6	AGP	F	7298[A]	16/16	0.89	0.23	2.53	36,37,38,39	16
6	AGP	B	3298[A]	16/16	0.85	0.25	2.36	36,39,42,43	16
5	SO4	C	4297[B]	5/5	0.95	0.11	2.28	34,36,37,38	0
2	GLC	F	295	11/12	0.83	0.15	1.78	23,25,27,32	11
2	GLC	D	295	11/12	0.86	0.13	1.78	26,28,29,31	11
5	SO4	A	2296[B]	5/5	0.88	0.31	1.49	52,52,53,53	5
5	SO4	C	4296[B]	5/5	0.92	0.15	0.91	61,61,61,62	5
2	GLC	A	295	11/12	0.90	0.10	0.15	22,23,24,27	11
5	SO4	B	3296[B]	5/5	0.90	0.13	0.12	55,56,56,56	5
3	16G	D	7299	19/19	0.97	0.08	0.04	16,17,18,19	0
3	16G	E	5299	19/19	0.97	0.08	0.01	19,21,27,27	0
3	16G	B	2299	19/19	0.97	0.08	-0.06	16,18,21,22	0
5	SO4	D	5296[B]	5/5	0.95	0.10	-0.09	44,44,45,45	5
5	SO4	F	7296[B]	5/5	0.96	0.11	-0.16	44,44,44,44	5
3	16G	F	6299	19/19	0.97	0.08	-0.17	18,20,22,24	0
3	16G	A	4299	19/19	0.97	0.07	-0.48	15,18,19,19	0
3	16G	C	3299	19/19	0.98	0.07	-0.51	14,16,18,19	0
5	SO4	E	6296[B]	5/5	0.96	0.07	-0.69	48,49,49,49	5
2	GLC	A	294	12/12	0.83	0.15	-	24,25,28,29	12
2	GLC	D	294	12/12	0.74	0.22	-	27,30,34,34	12
2	GLC	C	294	12/12	0.83	0.12	-	20,24,27,27	12
2	GLC	F	294	12/12	0.86	0.13	-	26,28,28,29	12
2	GLC	E	294	12/12	0.79	0.18	-	30,32,34,34	12

6.5 Other polymers [i](#)

There are no such residues in this entry.