



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:55 pm GMT

PDB ID : 4NHF  
Title : Crystal structure of the soluble domain of TrwG Type IV secretion machinery from *Bartonella grahamii*  
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2013-11-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	<div> <div style="width: 20%; background-color: red;"></div> <div style="width: 76%; background-color: green;"></div> <div style="width: 4%; background-color: yellow;"></div> <div style="width: 20%; background-color: grey;"></div> </div> <div>20%</div>
1	B	180	<div> <div style="width: 68%; background-color: green;"></div> <div style="width: 8%; background-color: yellow;"></div> <div style="width: 23%; background-color: grey;"></div> </div> <div>23%</div>
1	C	180	<div> <div style="width: 2%; background-color: red;"></div> <div style="width: 73%; background-color: green;"></div> <div style="width: 6%; background-color: yellow;"></div> <div style="width: 21%; background-color: grey;"></div> </div> <div>21%</div>
1	D	180	<div> <div style="width: 76%; background-color: green;"></div> <div style="width: 21%; background-color: grey;"></div> </div> <div>21%</div>
1	E	180	<div> <div style="width: 2%; background-color: red;"></div> <div style="width: 77%; background-color: green;"></div> <div style="width: 21%; background-color: grey;"></div> </div> <div>21%</div>
1	F	180	<div> <div style="width: 75%; background-color: green;"></div> <div style="width: 21%; background-color: grey;"></div> </div> <div>21%</div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TrwG protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	144	Total	C	N	O	S	0	5	0
			1161	727	202	231	1			
1	B	138	Total	C	N	O		0	2	0
			1094	688	193	213				
1	C	142	Total	C	N	O		0	2	0
			1110	697	191	222				
1	D	142	Total	C	N	O	S	0	1	0
			1121	702	193	225	1			
1	E	142	Total	C	N	O		0	1	0
			1106	693	191	222				
1	F	143	Total	C	N	O	S	0	1	0
			1127	710	195	221	1			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	54	MET	-	EXPRESSION TAG	UNP C6AAT5
A	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
A	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	61	HIS	-	EXPRESSION TAG	UNP C6AAT5
A	62	MET	-	EXPRESSION TAG	UNP C6AAT5
B	54	MET	-	EXPRESSION TAG	UNP C6AAT5
B	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
B	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
B	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
B	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
B	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
B	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
B	61	HIS	-	EXPRESSION TAG	UNP C6AAT5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	62	MET	-	EXPRESSION TAG	UNP C6AAT5
C	54	MET	-	EXPRESSION TAG	UNP C6AAT5
C	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
C	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	61	HIS	-	EXPRESSION TAG	UNP C6AAT5
C	62	MET	-	EXPRESSION TAG	UNP C6AAT5
D	54	MET	-	EXPRESSION TAG	UNP C6AAT5
D	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
D	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	61	HIS	-	EXPRESSION TAG	UNP C6AAT5
D	62	MET	-	EXPRESSION TAG	UNP C6AAT5
E	54	MET	-	EXPRESSION TAG	UNP C6AAT5
E	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
E	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	61	HIS	-	EXPRESSION TAG	UNP C6AAT5
E	62	MET	-	EXPRESSION TAG	UNP C6AAT5
F	54	MET	-	EXPRESSION TAG	UNP C6AAT5
F	55	ALA	-	EXPRESSION TAG	UNP C6AAT5
F	56	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	57	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	58	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	59	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	60	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	61	HIS	-	EXPRESSION TAG	UNP C6AAT5
F	62	MET	-	EXPRESSION TAG	UNP C6AAT5

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
-----	-------	----------	-------	---------	---------

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	144	Total O 144 144	0	0
3	B	121	Total O 121 121	0	0
3	C	73	Total O 73 73	0	0
3	D	106	Total O 106 106	0	0
3	E	76	Total O 76 76	0	0
3	F	92	Total O 92 92	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TrwG protein




- Molecule 1: TrwG protein




- Molecule 1: TrwG protein



- Molecule 1: TrwG protein




- Molecule 1: TrwG protein





● Molecule 1: TrwG protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.95Å 87.65Å 191.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 45.12 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (50.00-2.00) 99.1 (45.12-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.54 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.195 , 0.240 0.204 , 0.248	Depositor DCC
$R_{free}$ test set	3620 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtrriage
Anisotropy	0.243	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7335	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.39 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.1531e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.78	0/1191	0.86	2/1622 (0.1%)
1	B	0.78	0/1119	0.85	1/1523 (0.1%)
1	C	0.61	0/1136	0.76	0/1551
1	D	0.62	0/1141	0.77	1/1554 (0.1%)
1	E	0.64	0/1129	0.80	4/1542 (0.3%)
1	F	0.63	0/1150	0.74	0/1564
All	All	0.68	0/6866	0.80	8/9356 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1
1	F	0	1
All	All	0	2

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	99[A]	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	E	99[B]	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	E	99[A]	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	E	99[B]	ARG	NE-CZ-NH1	5.94	123.27	120.30
1	B	109	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	D	155	ARG	NE-CZ-NH2	-5.43	117.59	120.30
1	A	109	ARG	NE-CZ-NH1	5.31	122.96	120.30
1	A	98	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	91	THR	Peptide
1	F	166	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1161	0	1117	5	0
1	B	1094	0	1044	8	0
1	C	1110	0	1054	8	0
1	D	1121	0	1075	0	0
1	E	1106	0	1042	1	0
1	F	1127	0	1097	3	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	144	0	0	1	0
3	B	121	0	0	1	0
3	C	73	0	0	0	0
3	D	106	0	0	0	0
3	E	76	0	0	0	0
3	F	92	0	0	0	0
All	All	7335	0	6429	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:174[A]:ARG:HG2	1:C:194:ILE:HD12	1.77	0.67
1:B:170:GLN:NE2	3:B:497:HOH:O	2.31	0.62
1:A:174:ARG:HG2	1:A:194:ILE:HD12	1.83	0.61

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:95:GLU:HG2	1:B:99[A]:ARG:NH2	2.16	0.59
1:A:174:ARG:HG2	1:A:194:ILE:CD1	2.34	0.58
1:C:174[A]:ARG:HG2	1:C:194:ILE:CD1	2.37	0.54
1:B:166:ASN:HD21	1:B:172:THR:HG23	1.72	0.53
1:C:181:ASP:OD2	1:C:185:ALA:HB3	2.11	0.51
1:A:155[B]:ARG:NH1	3:A:305:HOH:O	2.45	0.49
1:B:107:LEU:O	1:B:111:ASN:HB2	2.16	0.45
1:B:126:LEU:HD11	1:B:210:ASP:HB3	1.99	0.45
1:B:95:GLU:HG3	1:B:165:LEU:HD12	1.99	0.44
1:C:163:ILE:HG12	1:C:173:VAL:HG22	1.98	0.44
1:F:107:LEU:O	1:F:111:ASN:HB2	2.18	0.44
1:C:107:LEU:O	1:C:111:ASN:HB2	2.19	0.43
1:C:99:ARG:NH1	1:C:163:ILE:O	2.42	0.43
1:A:187:THR:OG1	1:A:187:THR:O	2.32	0.43
1:B:99[B]:ARG:NH2	1:C:231:LEU:HD21	2.34	0.43
1:B:95:GLU:CG	1:B:99[A]:ARG:NH2	2.81	0.42
1:E:107:LEU:O	1:E:111:ASN:HB2	2.20	0.42
1:A:155[A]:ARG:HH21	1:A:155[A]:ARG:HG2	1.86	0.41
1:C:181:ASP:C	1:C:181:ASP:OD1	2.58	0.41
1:F:166:ASN:N	1:F:167:GLY:HA2	2.35	0.41
1:F:114:TYR:O	1:F:117[A]:ILE:HG22	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/180 (82%)	146 (99%)	1 (1%)	0	100	100
1	B	136/180 (76%)	131 (96%)	4 (3%)	1 (1%)	25	18
1	C	142/180 (79%)	139 (98%)	3 (2%)	0	100	100
1	D	141/180 (78%)	138 (98%)	2 (1%)	1 (1%)	25	18

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	141/180 (78%)	139 (99%)	2 (1%)	0	100	100
1	F	142/180 (79%)	138 (97%)	4 (3%)	0	100	100
All	All	849/1080 (79%)	831 (98%)	16 (2%)	2 (0%)	51	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	167	GLY
1	D	151	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/158 (79%)	121 (97%)	4 (3%)	44	42
1	B	114/158 (72%)	110 (96%)	4 (4%)	41	39
1	C	116/158 (73%)	114 (98%)	2 (2%)	66	70
1	D	120/158 (76%)	117 (98%)	3 (2%)	53	54
1	E	115/158 (73%)	114 (99%)	1 (1%)	82	87
1	F	120/158 (76%)	118 (98%)	2 (2%)	66	70
All	All	710/948 (75%)	694 (98%)	16 (2%)	60	58

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92[A]	SER
1	A	92[B]	SER
1	A	155[A]	ARG
1	A	155[B]	ARG
1	B	92	SER
1	B	126	LEU
1	B	174	ARG
1	B	231	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	182	SER
1	C	231	LEU
1	D	182	SER
1	D	183	SER
1	D	207	LYS
1	E	208	SER
1	F	183	SER
1	F	208	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	166	ASN
1	D	164	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	144/180 (80%)	-0.15	1 (0%) 87 87	12, 22, 42, 58	0
1	B	138/180 (76%)	-0.17	0 100 100	13, 24, 49, 65	0
1	C	142/180 (78%)	0.08	4 (2%) 53 53	17, 36, 65, 87	0
1	D	142/180 (78%)	-0.26	0 100 100	19, 30, 54, 75	0
1	E	142/180 (78%)	0.01	4 (2%) 53 53	17, 34, 61, 83	0
1	F	143/180 (79%)	-0.19	0 100 100	20, 31, 57, 74	0
All	All	851/1080 (78%)	-0.11	9 (1%) 80 80	12, 29, 59, 87	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	144	ASN	3.3
1	C	149	VAL	3.1
1	E	146	ARG	3.1
1	E	167	GLY	2.5
1	E	140	TYR	2.4
1	C	150	LEU	2.2
1	E	147	ASP	2.2
1	C	151	SER	2.1
1	A	167	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CA	E	301	1/1	0.99	0.09	-1.46	45,45,45,45	0
2	CA	B	301	1/1	0.99	0.05	-2.92	31,31,31,31	0
2	CA	F	301	1/1	0.99	0.07	-	42,42,42,42	0
2	CA	C	301	1/1	0.96	0.06	-	46,46,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.