



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:44 pm GMT

PDB ID : 4NKU
Title : Structure of Cid1 in complex with its short product ApU
Authors : Munoz-Tello, P.; Gabus, C.; Thore, S.
Deposited on : 2013-11-13
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3233 (1.96-1.92)
Clashscore	112137	3430 (1.96-1.92)
Ramachandran outliers	110173	3395 (1.96-1.92)
Sidechain outliers	110143	3395 (1.96-1.92)
RSRZ outliers	101464	3250 (1.96-1.92)
RNA backbone	2435	1003 (2.64-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>6%</div> </div> </div>
1	B	341	<div> <div>4%</div> <div> <div></div> <div>76%</div> <div>13%</div> <div>9%</div> </div> </div>
2	D	2	<div> <div></div> <div>100%</div> </div>
2	H	2	<div> <div></div> <div>50%</div> <div>50%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly(A) RNA polymerase protein cid1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	3	0
			2605	1691	431	472	11			
1	B	310	Total	C	N	O	S	0	2	0
			2524	1644	412	457	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	37	GLY	-	EXPRESSION TAG	UNP O13833
A	38	ASP	-	EXPRESSION TAG	UNP O13833
A	39	MET	-	EXPRESSION TAG	UNP O13833
A	160	ALA	ASP	ENGINEERED MUTATION	UNP O13833
B	37	GLY	-	EXPRESSION TAG	UNP O13833
B	38	ASP	-	EXPRESSION TAG	UNP O13833
B	39	MET	-	EXPRESSION TAG	UNP O13833
B	160	ALA	ASP	ENGINEERED MUTATION	UNP O13833

- Molecule 2 is a RNA chain called 5'-R(*AP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	2	Total	C	N	O	P	0	0	0
			39	19	7	12	1			
2	H	2	Total	C	N	O	P	0	0	0
			39	19	7	12	1			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Br 1	0	0
4	A	1	Total 1	Br 1	0	0
4	D	1	Total 1	Br 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	195	Total 195	O 195	0	0
5	B	152	Total 152	O 152	0	0
5	D	6	Total 6	O 6	0	0
5	H	3	Total 3	O 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Poly(A) RNA polymerase protein cid1



- Molecule 1: Poly(A) RNA polymerase protein cid1



- Molecule 2: 5'-R(*AP*U)-3'



- Molecule 2: 5'-R(*AP*U)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.02Å 77.09Å 81.83Å 90.00° 90.71° 90.00°	Depositor
Resolution (Å)	45.34 – 1.94 45.34 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.34-1.94) 99.6 (45.34-1.94)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.94Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.179 , 0.226 0.180 , 0.226	Depositor DCC
R_{free} test set	2002 reflections (4.04%)	DCC
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5569	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/2675	0.54	0/3606
1	B	0.39	0/2588	0.53	0/3489
2	D	0.37	0/43	0.53	0/65
2	H	0.42	0/43	0.64	0/65
All	All	0.40	0/5349	0.54	0/7225

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	366	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2605	0	2625	19	0
1	B	2524	0	2545	25	0
2	D	39	0	23	2	0
2	H	39	0	23	2	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	1	0
4	D	1	0	0	0	0
5	A	195	0	0	7	0
5	B	152	0	0	7	1
5	D	6	0	0	2	0
5	H	3	0	0	2	0
All	All	5569	0	5216	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:402:BR:BR	5:B:593:HOH:O	2.01	1.28
1:A:79:ILE:HD11	1:A:124:GLU:HG3	1.52	0.88
1:A:337:ASN:O	1:A:340:ARG:HG3	1.86	0.75
1:B:370:TYR:OH	5:B:535:HOH:O	2.06	0.73
1:A:323:ARG:N	5:A:637:HOH:O	2.22	0.72
1:B:39:MET:HG2	1:B:40:SER:H	1.55	0.71
2:H:59:U:OP1	5:H:102:HOH:O	2.11	0.69
2:H:59:U:O3'	5:H:103:HOH:O	2.10	0.69
2:D:59:U:OP1	5:D:206:HOH:O	2.11	0.68
1:A:38:ASP:N	5:A:606:HOH:O	2.32	0.62
1:A:293:ARG:NH2	5:A:525:HOH:O	2.34	0.60
1:A:138:ALA:O	5:A:616:HOH:O	2.17	0.60
1:B:108:MET:SD	1:B:117:ILE:HG21	2.45	0.56
1:B:167:LEU:HD21	1:B:246:ASP:HB2	1.87	0.56
1:A:175:LEU:O	1:A:179:THR:HG23	2.06	0.56
1:B:39:MET:N	5:B:650:HOH:O	2.40	0.54
1:A:197:LYS:NZ	5:A:670:HOH:O	2.39	0.54
1:B:175:LEU:O	1:B:179[A]:THR:HG23	2.08	0.52
1:B:84:GLU:HG3	1:B:107:LEU:HB2	1.91	0.52
1:B:78:ARG:HB2	5:B:529:HOH:O	2.10	0.52
2:D:58:A:N6	5:D:202:HOH:O	2.43	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:SER:O	1:B:179[B]:THR:HG22	2.10	0.51
1:B:91:LEU:HB2	5:B:501:HOH:O	2.11	0.51
1:A:235:LEU:HD23	1:A:293:ARG:HH21	1.77	0.49
1:B:325:ILE:HG22	1:B:347:LEU:HD22	1.93	0.49
1:B:307:THR:OG1	1:B:308:SER:N	2.45	0.48
1:A:286:ARG:HH22	1:B:375:GLU:HA	1.78	0.48
1:B:76:LEU:HD11	1:B:117:ILE:HD11	1.95	0.48
1:A:370:TYR:OH	5:A:629:HOH:O	2.19	0.47
1:A:109:ASP:N	5:A:679:HOH:O	2.48	0.47
1:B:39:MET:HG2	1:B:40:SER:N	2.27	0.47
1:B:124:GLU:HG2	5:B:529:HOH:O	2.14	0.47
1:B:65:LYS:HD3	1:B:91:LEU:HD11	1.97	0.47
1:B:116:THR:O	1:B:120:GLN:HG2	2.16	0.46
1:A:68:ALA:HA	1:A:153:PHE:CZ	2.51	0.46
1:B:46:LYS:NZ	1:B:50:GLU:OE1	2.49	0.45
1:A:291:THR:HG21	1:A:299:LEU:HB2	1.98	0.45
1:A:204:PRO:HG3	1:A:210:SER:HA	1.99	0.44
1:A:294:ARG:NH2	1:A:303:GLU:OE1	2.41	0.44
1:B:372:SER:HA	1:B:375:GLU:HG3	2.00	0.43
1:A:108:MET:HE3	1:A:108:MET:HB2	1.87	0.43
1:A:366:TYR:HB3	1:A:367:PRO:HD2	2.00	0.43
1:B:59:ASP:O	1:B:63:LYS:HD2	2.18	0.43
1:B:157:PHE:N	5:B:641:HOH:O	2.51	0.43
1:B:193:LYS:HE2	1:B:211:SER:HB3	2.01	0.43
1:B:310:THR:N	1:B:322:ASP:O	2.48	0.43
1:B:66:ARG:HD2	1:B:66:ARG:HA	1.93	0.42
1:A:365:SER:OG	1:A:366:TYR:N	2.50	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:592:HOH:O	5:B:644:HOH:O[2_656]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/341 (93%)	312 (98%)	4 (1%)	1 (0%)	44	33
1	B	302/341 (89%)	299 (99%)	3 (1%)	0	100	100
All	All	619/682 (91%)	611 (99%)	7 (1%)	1 (0%)	51	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	367	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/301 (95%)	268 (93%)	19 (7%)	19	6
1	B	279/301 (93%)	259 (93%)	20 (7%)	17	5
All	All	566/602 (94%)	527 (93%)	39 (7%)	18	6

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	GLU
1	A	66	ARG
1	A	116	THR
1	A	120	GLN
1	A	128	GLU
1	A	131	GLU
1	A	136	GLN
1	A	165	ASN
1	A	167	LEU
1	A	200	GLN
1	A	246	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	LYS
1	A	293	ARG
1	A	299	LEU
1	A	302	GLN
1	A	308	SER
1	A	340	ARG
1	A	366	TYR
1	A	376	GLU
1	B	39	MET
1	B	63	LYS
1	B	66	ARG
1	B	101	ASP
1	B	117	ILE
1	B	119	LEU
1	B	123	GLU
1	B	135	LEU
1	B	140	ILE
1	B	200	GLN
1	B	239	LEU
1	B	277	ARG
1	B	286	ARG
1	B	287	GLU
1	B	299	LEU
1	B	310	THR
1	B	323	ARG
1	B	344	SER
1	B	371	ASP
1	B	376	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	1/2 (50%)	0	0
2	H	1/2 (50%)	0	0
All	All	2/4 (50%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/341 (93%)	0.10	6 (1%) 67 75	16, 31, 49, 76	0
1	B	310/341 (90%)	0.21	13 (4%) 37 45	18, 33, 55, 69	0
2	D	2/2 (100%)	-0.24	0 100 100	32, 32, 32, 37	2 (100%)
2	H	2/2 (100%)	0.73	0 100 100	35, 35, 35, 38	2 (100%)
All	All	634/686 (92%)	0.16	19 (2%) 51 60	16, 32, 53, 76	4 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	140	ILE	6.7
1	A	366	TYR	5.5
1	B	377	ALA	4.4
1	A	139	ARG	3.5
1	B	126	ILE	3.0
1	B	310	THR	2.9
1	B	157	PHE	2.9
1	A	206	PHE	2.8
1	B	39	MET	2.7
1	B	136	GLN	2.7
1	A	364	ARG	2.5
1	B	348	TYR	2.5
1	B	119	LEU	2.4
1	B	308	SER	2.4
1	B	147	SER	2.1
1	B	307	THR	2.1
1	A	377	ALA	2.1
1	A	109	ASP	2.1
1	B	205	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	BR	D	101	1/1	0.99	0.12	1.19	30,30,30,30	1
4	BR	B	402	1/1	0.99	0.11	0.10	38,38,38,38	1
3	MG	A	401	1/1	0.96	0.10	-0.49	42,42,42,42	0
3	MG	A	402	1/1	0.96	0.08	-0.96	38,38,38,38	0
3	MG	B	401	1/1	0.98	0.09	-1.20	31,31,31,31	0
4	BR	A	403	1/1	0.99	0.04	-	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.