



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2017 – 09:56 AM EDT

PDB ID : 1NPZ  
Title : Crystal structures of Cathepsin S inhibitor complexes  
Authors : Pauly, T.A.; Sulea, T.; Ammirati, M.; Sivaraman, J.; Danley, D.E.; Griffor, M.C.; Kamath, A.V.; Wang, I.K.; Laird, E.R.; Seddon, A.P.; Menard, R.; Cygler, M.; Rath, V.L.  
Deposited on : unknown  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div> <div>4%</div> <div>86%</div> <div>14%</div> </div>
1	B	217	<div> <div>3%</div> <div>84%</div> <div>15%</div> <div>.</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3623 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cathepsin S.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1687	1065	286	324	12			
1	B	217	Total	C	N	O	S	0	0	0
			1687	1065	286	324	12			

- Molecule 2 is N 2 -(morpholin-4-ylcarbonyl)-N-[(3S)-1-phenyl-5-(phenylsulfonyl)pentan-3-yl]-L-leucinamide (three-letter code: C1P) (formula: C<sub>28</sub>H<sub>39</sub>N<sub>3</sub>O<sub>5</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			37	28	3	5	1		
2	B	1	Total	C	N	O	S	0	0
			37	28	3	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total 87	O 87	0	0
3	B	88	Total 88	O 88	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cathepsin S



#### • Molecule 1: Cathepsin S



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.08Å 108.08Å 105.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.00 29.44 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-2.00) 93.6 (29.44-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.45 (at 2.00Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.192 , 0.219 0.192 , 0.217	Depositor DCC
$R_{free}$ test set	1416 reflections (4.90%)	DCC
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.023 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3623	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C1P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.59	1/1729 (0.1%)	0.70	0/2337
1	B	0.63	0/1729	0.72	0/2337
All	All	0.61	1/3458 (0.0%)	0.71	0/4674

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	175	ASN	CB-CG	6.00	1.64	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	195	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1687	0	1610	23	0
1	B	1687	0	1605	23	0
2	A	37	0	38	1	0
2	B	37	0	38	1	0
3	A	87	0	0	6	0
3	B	88	0	0	1	0
All	All	3623	0	3291	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:427:LYS:HA	1:B:427:LYS:HE2	1.26	1.10
1:B:366:LEU:HB2	1:B:462:PRO:HG2	1.41	1.01
1:A:116:LEU:HB2	1:A:212:PRO:HG2	1.48	0.95
1:A:173:ASP:OD2	3:A:507:HOH:O	2.00	0.80
3:A:507:HOH:O	1:B:438:HIS:ND1	2.17	0.77
1:A:177:LYS:HE2	1:A:177:LYS:HA	1.65	0.76
1:B:251:LEU:HD11	1:B:376:GLU:HG2	1.69	0.73
1:B:410:GLN:NE2	1:B:454:ASN:HD22	1.91	0.67
1:A:14:THR:HB	1:A:47:THR:HG22	1.76	0.67
1:B:427:LYS:HA	1:B:427:LYS:CE	2.15	0.65
1:B:427:LYS:CA	1:B:427:LYS:HE2	2.12	0.61
1:A:14:THR:CB	1:A:47:THR:HG22	2.32	0.59
1:A:1:LEU:HD11	1:A:126:GLU:HG2	1.86	0.57
1:A:93:LYS:HG3	1:A:97:GLN:NE2	2.21	0.56
1:A:31:VAL:O	1:A:35:GLU:HG3	2.08	0.54
1:A:174:LEU:HA	1:B:265:GLU:HG2	1.90	0.54
3:A:504:HOH:O	1:B:265:GLU:HB3	2.07	0.53
1:B:368:TYR:OH	1:B:410:GLN:HG2	2.10	0.52
1:A:164:HIS:ND1	2:A:468:C1P:H17	2.25	0.51
1:A:81:ASN:O	1:A:82:LYS:HB2	2.11	0.50
1:A:174:LEU:HD13	1:B:265:GLU:HG3	1.95	0.49
1:B:414:HIS:ND1	2:B:469:C1P:H17	2.28	0.49
1:A:93:LYS:HG3	1:A:97:GLN:HE22	1.76	0.49
1:A:100:GLN:O	1:A:100:GLN:HG3	2.14	0.47
1:B:310:LYS:HB3	3:B:658:HOH:O	2.14	0.47
1:B:261:GLY:O	1:B:295:LEU:HD22	2.15	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:255:VAL:HG23	1:B:420:GLY:HA2	1.97	0.46
1:B:258:ARG:HD3	1:B:445:TYR:CZ	2.52	0.45
1:B:264:THR:HG21	1:B:297:THR:HG22	1.99	0.45
1:B:389:ASP:OD2	1:B:392:HIS:HE1	2.01	0.44
1:A:102:ASP:OD1	1:A:104:LYS:HE2	2.18	0.44
1:B:307:SER:O	1:B:313:ASN:HB2	2.17	0.43
1:B:442:GLU:OE1	1:B:447:ARG:NH2	2.52	0.43
1:A:146:PHE:HB2	3:A:558:HOH:O	2.17	0.43
1:A:86:SER:HB3	1:A:107:ALA:HB2	2.01	0.43
1:B:429:TYR:CD1	1:B:429:TYR:C	2.91	0.43
1:A:100:GLN:HB3	3:A:566:HOH:O	2.18	0.42
1:A:102:ASP:OD1	1:A:104:LYS:HG3	2.19	0.42
1:A:102:ASP:OD2	1:A:104:LYS:HE2	2.20	0.42
1:B:281:VAL:O	1:B:285:GLU:HG3	2.19	0.42
1:B:306:CYS:CB	1:B:349:CYS:SG	3.08	0.42
1:A:14:THR:CG2	1:A:47:THR:HG22	2.50	0.41
1:A:100:GLN:CB	3:A:566:HOH:O	2.68	0.41
1:A:148:TYR:CE1	1:A:196:ILE:HG13	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/217 (99%)	208 (97%)	7 (3%)	0	100	100
1	B	215/217 (99%)	207 (96%)	8 (4%)	0	100	100
All	All	430/434 (99%)	415 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	178/178 (100%)	173 (97%)	5 (3%)	49	49
1	B	178/178 (100%)	172 (97%)	6 (3%)	42	40
All	All	356/356 (100%)	345 (97%)	11 (3%)	45	44

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	90	TYR
1	A	133	PRO
1	A	158	CYS
1	A	160	GLN
1	B	302	ASN
1	B	340	TYR
1	B	383	PRO
1	B	408	CYS
1	B	410	GLN
1	B	438	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	GLN
1	A	130	ASN
1	B	326	GLN
1	B	380	ASN
1	B	392	HIS
1	B	410	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	C1P	A	468	1	39,39,39	2.24	15 (38%)	50,52,52	1.92	12 (24%)
2	C1P	B	469	1	39,39,39	2.58	14 (35%)	50,52,52	1.96	13 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C1P	A	468	1	-	0/37/45/45	0/3/3/3
2	C1P	B	469	1	-	0/37/45/45	0/3/3/3

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	469	C1P	C27-C22	2.14	1.43	1.38
2	A	468	C1P	O12-C13	2.19	1.51	1.42
2	A	468	C1P	C23-C22	2.29	1.43	1.38
2	A	468	C1P	C17-S33	2.42	1.83	1.78
2	A	468	C1P	C30-C31	2.43	1.43	1.38
2	B	469	C1P	C26-C25	2.47	1.44	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	468	C1P	C27-C22	2.49	1.44	1.38
2	B	469	C1P	C36-C32	2.50	1.42	1.38
2	A	468	C1P	C24-C23	2.50	1.43	1.38
2	B	469	C1P	C23-C22	2.51	1.44	1.38
2	A	468	C1P	C26-C25	2.61	1.44	1.38
2	B	469	C1P	C37-C36	2.62	1.43	1.38
2	B	469	C1P	C7-C6	2.67	1.66	1.51
2	A	468	C1P	C36-C32	2.70	1.43	1.38
2	A	468	C1P	C26-C27	2.74	1.44	1.38
2	B	469	C1P	C17-S33	2.87	1.84	1.78
2	A	468	C1P	C37-C36	3.00	1.44	1.38
2	B	469	C1P	C26-C27	3.00	1.44	1.38
2	B	469	C1P	C24-C23	3.65	1.45	1.38
2	A	468	C1P	C31-C32	3.69	1.44	1.38
2	A	468	C1P	C14-N9	3.85	1.53	1.47
2	A	468	C1P	C18-C17	4.15	1.57	1.52
2	B	469	C1P	C14-N9	4.16	1.54	1.47
2	B	469	C1P	C18-C17	4.81	1.57	1.52
2	B	469	C1P	C31-C32	4.82	1.46	1.38
2	A	468	C1P	O35-S33	5.29	1.52	1.44
2	A	468	C1P	O34-S33	5.92	1.53	1.44
2	B	469	C1P	O35-S33	6.67	1.54	1.44
2	B	469	C1P	O34-S33	7.58	1.56	1.44

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	469	C1P	O35-S33-O34	-6.92	109.70	118.42
2	A	468	C1P	O35-S33-O34	-6.86	109.78	118.42
2	A	468	C1P	O16-C15-N9	-4.99	114.88	121.73
2	B	469	C1P	O34-S33-C32	-4.28	103.83	108.39
2	A	468	C1P	O34-S33-C32	-3.80	104.34	108.39
2	B	469	C1P	O16-C15-N9	-3.08	117.50	121.73
2	B	469	C1P	O35-S33-C32	-2.98	105.21	108.39
2	A	468	C1P	C20-C19-N28	-2.62	106.78	110.58
2	B	469	C1P	C31-C32-C36	-2.53	116.82	120.43
2	A	468	C1P	C31-C32-C36	-2.41	116.98	120.43
2	B	469	C1P	C3-C2-N1	-2.03	105.62	111.20
2	B	469	C1P	C10-N9-C14	2.05	116.49	112.61
2	A	468	C1P	C10-N9-C14	2.09	116.56	112.61
2	A	468	C1P	C13-C14-N9	2.10	114.40	109.86
2	A	468	C1P	C17-S33-C32	2.12	107.98	105.03

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	469	C1P	C31-C32-S33	2.22	121.99	119.50
2	A	468	C1P	C30-C31-C32	2.33	121.42	118.96
2	B	469	C1P	C13-C14-N9	2.33	114.91	109.86
2	A	468	C1P	O34-S33-C17	2.51	112.31	108.18
2	B	469	C1P	O34-S33-C17	2.66	112.56	108.18
2	B	469	C1P	C30-C31-C32	2.70	121.80	118.96
2	B	469	C1P	N1-C15-N9	2.96	122.74	117.20
2	A	468	C1P	N1-C15-N9	3.99	124.66	117.20
2	A	468	C1P	O35-S33-C17	4.01	114.77	108.18
2	B	469	C1P	O35-S33-C17	5.57	117.35	108.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	468	C1P	1	0
2	B	469	C1P	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	217/217 (100%)	0.05	9 (4%) 38 38	17, 26, 38, 43	0
1	B	217/217 (100%)	-0.11	6 (2%) 53 53	15, 23, 34, 38	0
All	All	434/434 (100%)	-0.03	15 (3%) 44 45	15, 24, 37, 43	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	ARG	3.8
1	B	391	ARG	3.6
1	A	104	LYS	3.4
1	B	425	ASN	3.3
1	A	105	TYR	3.3
1	A	100	GLN	3.0
1	B	350	GLN	2.9
1	B	355	TYR	2.4
1	A	44	LYS	2.4
1	A	188	HIS	2.3
1	A	15	GLU	2.3
1	A	211	PHE	2.2
1	A	41	LYS	2.2
1	B	354	LYS	2.1
1	B	265	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	C1P	A	468	37/37	0.95	0.12	-0.03	20,22,23,23	0
2	C1P	B	469	37/37	0.96	0.11	-0.03	18,20,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.