



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:27 am GMT

PDB ID : 4NQF  
Title : Crystal structure of HEPN domain protein  
Authors : Padavannil, A.; Jobichen, C.; Sivaraman, J.  
Deposited on : 2013-11-25  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	158	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 16%, green 75%, grey 8%);"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> <span style="display: inline-block; width: 1%;"></span> <span style="display: inline-block; width: 1%;"></span> <span style="display: inline-block; width: 14%;"></span> <span style="display: inline-block; width: 64%;"></span> <span style="display: inline-block; width: 1%;"></span> <span style="display: inline-block; width: 1%;"></span> </div> </div> </div>
1	B	158	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%; text-align: center;">2%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 17%, green 78%, grey 2%);"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> <span style="display: inline-block; width: 2%;"></span> <span style="display: inline-block; width: 2%;"></span> <span style="display: inline-block; width: 15%;"></span> <span style="display: inline-block; width: 61%;"></span> <span style="display: inline-block; width: 2%;"></span> <span style="display: inline-block; width: 2%;"></span> </div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleotidyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	Se	0	0	0
			1197	759	217	216	2	3			
1	B	151	Total	C	N	O	S	Se	0	0	0
			1230	779	224	222	2	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	168	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19
A	219	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19
A	251	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19
B	168	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19
B	219	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19
B	251	MSE	LEU	ENGINEERED MUTATION	UNP Q7CW19


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	O	0	0
			12	12		
2	B	14	Total	O	0	0
			14	14		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Nucleotidyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.10Å 71.57Å 83.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.04 – 2.70 36.04 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.0 (36.04-2.70) 99.1 (36.04-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	30.52 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.218 , 0.280 0.223 , 0.279	Depositor DCC
$R_{free}$ test set	1075 reflections (11.12%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.7	Xtriage
Anisotropy	0.350	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2453	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.90 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.9071e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/1223	0.68	0/1652
1	B	0.60	0/1257	0.74	1/1700 (0.1%)
All	All	0.58	0/2480	0.71	1/3352 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	229	ARG	NE-CZ-NH1	5.34	122.97	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1197	0	1161	19	1
1	B	1230	0	1183	14	1
2	A	12	0	0	2	0
2	B	14	0	0	0	0
All	All	2453	0	2344	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:159:VAL:HG11	1:A:211:ASN:HD21	1.42	0.82
1:B:159:VAL:HG11	1:B:211:ASN:HD21	1.43	0.81
1:A:168:MSE:CE	1:A:204:CYS:SG	2.73	0.77
1:A:168:MSE:HE2	1:A:204:CYS:SG	2.24	0.77
1:A:151:LEU:N	2:A:406:HOH:O	2.23	0.70
1:B:214:PRO:O	1:B:215:ALA:HB3	1.94	0.68
1:A:214:PRO:O	1:A:215:ALA:HB3	1.98	0.64
1:B:254:ALA:O	1:B:258:ALA:HB3	2.03	0.58
1:A:254:ALA:O	1:A:258:ALA:HB3	2.03	0.57
1:B:293:ILE:HD12	1:B:293:ILE:O	2.04	0.57
1:A:293:ILE:O	1:A:293:ILE:HD12	2.05	0.56
1:B:214:PRO:O	1:B:215:ALA:CB	2.54	0.56
1:A:214:PRO:O	1:A:215:ALA:CB	2.56	0.54
1:A:168:MSE:HE3	1:A:204:CYS:SG	2.48	0.53
1:A:159:VAL:HG11	1:A:211:ASN:ND2	2.18	0.53
1:B:206:LEU:HD12	1:B:222:LEU:HD22	1.91	0.52
1:A:290:ARG:NH1	2:A:411:HOH:O	2.43	0.51
1:B:159:VAL:HG11	1:B:211:ASN:ND2	2.19	0.51
1:A:243:ARG:HD2	1:A:247:TRP:CH2	2.46	0.51
1:A:236:ILE:O	1:A:277:ARG:HD3	2.13	0.48
1:B:263:ARG:HD2	1:B:263:ARG:O	2.14	0.48
1:B:243:ARG:HD2	1:B:247:TRP:CH2	2.49	0.47
1:B:294:GLU:OE1	1:B:294:GLU:N	2.48	0.47
1:B:168:MSE:N	1:B:169:PRO:CD	2.77	0.46
1:A:290:ARG:O	1:A:293:ILE:HG23	2.15	0.46
1:B:236:ILE:O	1:B:277:ARG:HD3	2.15	0.45
1:B:263:ARG:CD	1:B:263:ARG:O	2.64	0.45
1:B:290:ARG:O	1:B:293:ILE:HG23	2.16	0.45
1:A:214:PRO:HG2	1:A:222:LEU:HD12	1.99	0.45
1:A:168:MSE:N	1:A:169:PRO:CD	2.80	0.44
1:A:217:HIS:HA	1:A:222:LEU:HD21	2.01	0.43
1:A:159:VAL:CG1	1:A:211:ASN:HD21	2.24	0.41
1:A:151:LEU:O	1:A:152:SER:CB	2.68	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:ARG:NH2	1:B:162:ASP:OD1[2_445]	2.17	0.03

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/158 (90%)	135 (94%)	6 (4%)	2 (1%)	13	33
1	B	149/158 (94%)	137 (92%)	9 (6%)	3 (2%)	9	22
All	All	292/316 (92%)	272 (93%)	15 (5%)	5 (2%)	11	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	SER
1	B	152	SER
1	B	300	ALA
1	B	257	LYS
1	A	257	LYS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/131 (95%)	117 (94%)	7 (6%)	25	51
1	B	125/131 (95%)	119 (95%)	6 (5%)	30	59
All	All	249/262 (95%)	236 (95%)	13 (5%)	27	55

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	165	LEU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	253	GLU
1	A	256	VAL
1	A	257	LYS
1	A	261	SER
1	A	266	VAL
1	A	284	LEU
1	B	151	LEU
1	B	165	LEU
1	B	263	ARG
1	B	266	VAL
1	B	295	LYS
1	B	296	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	211	ASN
1	B	163	HIS
1	B	211	ASN
1	B	275	GLN
1	B	298	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/158 (89%)	0.18	2 (1%) 75 76	38, 70, 109, 127	0
1	B	148/158 (93%)	0.02	3 (2%) 65 66	34, 58, 94, 111	0
All	All	290/316 (91%)	0.10	5 (1%) 70 72	34, 64, 104, 127	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	262	LYS	2.4
1	B	301	GLY	2.3
1	B	269	GLU	2.3
1	A	263	ARG	2.2
1	A	151	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.