



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 3, 2018 – 11:35 AM EST

PDB ID : 5NQ7
Title : Crystal structure of laccases from *Pycnoporus sanguineus*, isoform I
Authors : Orlikowska, M.; de J.Rostro-Alanis, M.; Bujacz, A.; Hernandez-Luna, C.; Rubio, R.; Parra, R.; Bujacz, G.
Deposited on : 2017-04-19
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3666 (2.80-2.72)
Clashscore	112137	4174 (2.80-2.72)
Ramachandran outliers	110173	4103 (2.80-2.72)
Sidechain outliers	110143	4106 (2.80-2.72)
RSRZ outliers	101464	3697 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	607	-	-	-	X
3	BMA	A	611	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Laccase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	497	3820	2430	647	733	10	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	8	LEU	PHE	conflict	UNP D7F485
A	9	CYS	PHE	conflict	UNP D7F485
A	13	VAL	ALA	conflict	UNP D7F485
A	185	LEU	VAL	conflict	UNP D7F485
A	241	ILE	VAL	conflict	UNP D7F485
A	515	PRO	LEU	conflict	UNP D7F485

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			10	6	4		
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	Cu	0	0
			4	4		

- Molecule 6 is PEROXIDE ION (three-letter code: PER) (formula: O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O 2 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	163	Total O 163 163	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Laccase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₃ 3 2	Depositor
Cell constants a, b, c, α , β , γ	185.55Å 185.55Å 185.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.39 – 2.75 46.39 – 2.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.39-2.75) 100.0 (46.39-2.75)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.49 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.161 , 0.207 0.170 , 0.210	Depositor DCC
R_{free} test set	1449 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	53.0	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4182	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CU, NAG, PER, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/3942	0.72	0/5415

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3820	0	3617	13	0
2	A	84	0	72	1	0
3	A	65	0	52	1	0
4	A	44	0	39	0	0
5	A	4	0	0	0	0
6	A	2	0	0	0	0
7	A	163	0	0	2	0
All	All	4182	0	3780	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 14 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:ALA:HB3	7:A:740:HOH:O	2.01	0.60
1:A:100:VAL:HG11	1:A:469:PHE:CD2	2.36	0.59
1:A:171:ASP:HB2	1:A:187:ALA:HB2	1.86	0.57
1:A:52:THR:HA	1:A:53:PRO:C	2.25	0.56
1:A:229:ASN:O	1:A:285:PRO:CD	2.55	0.54

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	496/518 (96%)	472 (95%)	22 (4%)	2 (0%)	38 69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	ASP
1	A	79	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	420/436 (96%)	408 (97%)	12 (3%)	48 78

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	264	ARG
1	A	319	LYS
1	A	469	PHE
1	A	226	CYS
1	A	414	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 21 ligands modelled in this entry, 4 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	601	1,2	14,14,15	0.33	0	15,19,21	0.58	0
2	NAG	A	602	3,2	14,14,15	0.36	0	15,19,21	0.64	0
3	BMA	A	603	3,2,4	11,11,12	0.26	0	13,15,17	0.41	0
4	MAN	A	604	3,4	11,11,12	0.36	0	13,15,17	0.74	0
4	MAN	A	605	4	11,11,12	0.41	0	13,15,17	0.59	0
3	BMA	A	606	3	11,11,12	0.48	0	13,15,17	0.76	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	607	1,2	14,14,15	0.68	0	15,19,21	0.78	1 (6%)
2	NAG	A	608	3,2	14,14,15	0.63	0	15,19,21	0.91	0
3	BMA	A	609	3,2	11,11,12	0.37	0	13,15,17	0.42	0
3	BMA	A	610	3,4	10,10,12	0.37	0	10,13,17	1.36	2 (20%)
3	BMA	A	611	3	11,11,12	0.48	0	13,15,17	0.93	1 (7%)
4	MAN	A	612	3	11,11,12	0.33	0	13,15,17	0.54	0
2	NAG	A	613	1,2	14,14,15	0.53	0	15,19,21	0.49	0
2	NAG	A	614	3,2	14,14,15	0.56	0	15,19,21	0.72	1 (6%)
3	BMA	A	615	2,4	11,11,12	0.44	0	13,15,17	0.96	1 (7%)
4	MAN	A	616	3	11,11,12	0.31	0	13,15,17	0.57	0
6	PER	A	621	5	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	602	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	603	3,2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	604	3,4	-	0/2/19/22	0/1/1/1
4	MAN	A	605	4	-	0/2/19/22	1/1/1/1
3	BMA	A	606	3	-	0/2/19/22	0/1/1/1
2	NAG	A	607	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	608	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	609	3,2	-	0/2/19/22	0/1/1/1
3	BMA	A	610	3,4	-	0/2/16/22	0/1/1/1
3	BMA	A	611	3	-	0/2/19/22	0/1/1/1
4	MAN	A	612	3	-	0/2/19/22	0/1/1/1
2	NAG	A	613	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	614	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	615	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	616	3	-	0/2/19/22	1/1/1/1
6	PER	A	621	5	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	614	NAG	O5-C1-C2	-2.28	108.30	111.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	610	BMA	O5-C1-C2	-2.02	106.64	110.71
3	A	615	BMA	C3-C4-C5	2.08	113.88	110.22
3	A	611	BMA	C1-O5-C5	2.40	115.47	112.17
2	A	607	NAG	C4-C3-C2	2.48	114.65	111.02

There are no chirality outliers.

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	605	MAN	C1-C2-C3-C4-C5-O5
4	A	616	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	608	NAG	1	0
3	A	609	BMA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	497/518 (95%)	-0.35	0 100 100	37, 49, 68, 89	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	BMA	A	611	11/12	0.79	0.42	15.04	84,118,125,135	0
2	NAG	A	607	14/15	0.78	0.29	7.42	93,132,142,143	0
2	NAG	A	601	14/15	0.98	0.23	0.67	44,51,61,61	0
2	NAG	A	613	14/15	0.97	0.20	0.40	49,65,71,83	0
2	NAG	A	602	14/15	0.96	0.23	0.19	49,60,71,90	0
5	CU	A	618	1/1	1.00	0.13	0.18	44,44,44,44	0
5	CU	A	617	1/1	0.99	0.13	-0.64	47,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	CU	A	620	1/1	0.99	0.11	-1.06	50,50,50,50	0
6	PER	A	621	2/2	0.99	0.10	-2.92	45,45,45,57	0
3	BMA	A	610	10/12	0.74	0.26	-	112,122,129,132	0
3	BMA	A	606	11/12	0.77	0.45	-	97,126,136,140	0
5	CU	A	619	1/1	0.99	0.12	-	45,45,45,45	0
4	MAN	A	605	11/12	0.74	0.32	-	103,121,129,132	0
2	NAG	A	614	14/15	0.91	0.31	-	71,90,98,108	0
3	BMA	A	615	11/12	0.88	0.45	-	116,128,136,143	0
4	MAN	A	616	11/12	0.74	0.44	-	104,126,143,150	0
2	NAG	A	608	14/15	0.76	0.28	-	110,141,153,161	0
3	BMA	A	603	11/12	0.87	0.37	-	100,108,129,130	0
4	MAN	A	604	11/12	0.85	0.40	-	103,129,140,144	0
4	MAN	A	612	11/12	0.62	0.36	-	105,129,149,162	0
3	BMA	A	609	11/12	0.78	0.26	-	123,147,159,162	0

6.5 Other polymers [i](#)

There are no such residues in this entry.