



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 04:53 am GMT

PDB ID : 4NRU
Title : Murine Norovirus RNA-dependent-RNA-polymerase in complex with Compound 6, a suramin derivative
Authors : Milani, M.; Croci, R.; Pezzullo, M.; Tarantino, D.; Mastrangelo, E.; Bolognesi, M.
Deposited on : 2013-11-27
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>7%</div> </div> </div>
1	B	515	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>
1	C	515	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>7%</div> </div> </div>
1	D	515	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>11%</div> <div>7%</div> </div> </div>
1	E	515	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>7%</div> </div> </div>
1	F	515	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>14%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 24851 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	477	Total	C	N	O	S	0	4	0
			3821	2421	674	701	25			
1	B	482	Total	C	N	O	S	0	1	0
			3835	2424	678	709	24			
1	C	478	Total	C	N	O	S	0	0	0
			3803	2406	670	703	24			
1	D	479	Total	C	N	O	S	0	2	0
			3820	2415	674	707	24			
1	E	477	Total	C	N	O	S	0	2	0
			3807	2408	672	702	25			
1	F	473	Total	C	N	O	S	0	0	0
			3761	2381	659	697	24			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7
A	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
A	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
A	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
A	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
A	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
A	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
A	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7
B	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
B	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
B	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
C	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
C	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7
D	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
D	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
D	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7
E	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
E	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
E	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	508	LEU	-	EXPRESSION TAG	UNP S4V9Y7
F	509	GLU	-	EXPRESSION TAG	UNP S4V9Y7
F	510	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	511	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	512	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	513	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	514	HIS	-	EXPRESSION TAG	UNP S4V9Y7
F	515	HIS	-	EXPRESSION TAG	UNP S4V9Y7

- Molecule 2 is 4-(4-METHYL-3-[(3-NITROBENZOYL)AMINO]BENZOYL)AMINO)NAPHTHALENE-1,5-DISULFONIC ACID (three-letter code: 2NG) (formula: C₂₅H₁₉N₃O₁₀S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	B	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	C	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	D	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	E	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	E	1	Total 40	C 25	N 3	O 10	S 2	0	0
2	F	1	Total 40	C 25	N 3	O 10	S 2	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Mg 1 1	0	0
3	E	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	384	Total 385	O 385	0	1
4	B	331	Total 332	O 332	0	1
4	C	304	Total 305	O 305	0	1
4	D	302	Total 303	O 303	0	1
4	E	248	Total 248	O 248	0	1
4	F	145	Total 145	O 145	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: RNA dependent RNA polymerase




• Molecule 1: RNA dependent RNA polymerase





• Molecule 1: RNA dependent RNA polymerase









• Molecule 1: RNA dependent RNA polymerase



• Molecule 1: RNA dependent RNA polymerase



• Molecule 1: RNA dependent RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.22Å 162.42Å 122.96Å 90.00° 97.04° 90.00°	Depositor
Resolution (Å)	61.09 – 2.30 61.02 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (61.09-2.30) 99.8 (61.02-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.191 , 0.254 0.194 , 0.254	Depositor DCC
R_{free} test set	9440 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	39.1	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	24851	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 2NG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3928	0.71	0/5319
1	B	0.49	0/3932	0.70	2/5324 (0.0%)
1	C	0.49	0/3898	0.69	0/5281
1	D	0.49	0/3918	0.70	0/5305
1	E	0.47	0/3906	0.69	0/5290
1	F	0.45	0/3853	0.65	0/5218
All	All	0.49	0/23435	0.69	2/31737 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	412	ASP	CB-CG-OD1	5.85	123.56	118.30
1	B	413	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3821	0	3792	36	0
1	B	3835	0	3794	25	0
1	C	3803	0	3761	51	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3820	0	3784	33	0
1	E	3807	0	3770	34	0
1	F	3761	0	3720	41	0
2	A	40	0	19	5	0
2	B	40	0	19	2	0
2	C	40	0	19	2	0
2	D	40	0	19	1	0
2	E	80	0	38	5	0
2	F	40	0	19	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	A	385	0	0	2	0
4	B	332	0	0	1	0
4	C	305	0	0	3	0
4	D	303	0	0	3	0
4	E	248	0	0	3	0
4	F	145	0	0	1	0
All	All	24851	0	22754	229	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 229 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:364:ARG:HB2	1:F:364:ARG:NH1	1.24	1.47
2:F:1101:2NG:NBL	2:F:1101:2NG:OAJ	1.73	1.19
1:F:364:ARG:CB	1:F:364:ARG:NH1	2.11	1.13
1:F:364:ARG:CB	1:F:364:ARG:HH11	1.68	1.04
1:C:435:GLN:CB	1:C:437:PRO:HD3	1.90	1.00

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/515 (92%)	465 (98%)	10 (2%)	0	100	100
1	B	477/515 (93%)	465 (98%)	12 (2%)	0	100	100
1	C	474/515 (92%)	465 (98%)	8 (2%)	1 (0%)	51	63
1	D	475/515 (92%)	463 (98%)	11 (2%)	1 (0%)	51	63
1	E	473/515 (92%)	460 (97%)	12 (2%)	1 (0%)	51	63
1	F	467/515 (91%)	448 (96%)	19 (4%)	0	100	100
All	All	2841/3090 (92%)	2766 (97%)	72 (2%)	3 (0%)	55	67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	436	ARG
1	D	436	ARG
1	E	38	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/441 (93%)	408 (99%)	4 (1%)	80	90
1	B	412/441 (93%)	396 (96%)	16 (4%)	37	51
1	C	409/441 (93%)	387 (95%)	22 (5%)	26	35
1	D	411/441 (93%)	386 (94%)	25 (6%)	22	29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	410/441 (93%)	387 (94%)	23 (6%)	25	33
1	F	405/441 (92%)	384 (95%)	21 (5%)	27	36
All	All	2459/2646 (93%)	2348 (96%)	111 (4%)	32	44

5 of 111 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	169	LEU
1	D	440	LEU
1	F	316	LEU
1	D	216	ARG
1	D	294	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	238	HIS
1	D	66	GLN
1	F	148	ASN
1	C	313	HIS
1	C	439	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 6 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	2NG	A	1101	-	42,43,43	2.85	8 (19%)	60,65,65	1.31	6 (10%)
2	2NG	B	1101	-	42,43,43	2.88	9 (21%)	60,65,65	1.12	5 (8%)
2	2NG	C	1101	-	42,43,43	2.95	7 (16%)	60,65,65	1.48	9 (15%)
2	2NG	D	1101	-	42,43,43	3.50	10 (23%)	60,65,65	1.56	11 (18%)
2	2NG	E	1101	-	42,43,43	2.90	8 (19%)	60,65,65	1.43	9 (15%)
2	2NG	E	1102	-	42,43,43	3.09	12 (28%)	60,65,65	1.73	13 (21%)
2	2NG	F	1101	-	42,43,43	3.56	8 (19%)	60,65,65	1.69	10 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2NG	A	1101	-	-	0/30/32/32	0/4/4/4
2	2NG	B	1101	-	-	0/30/32/32	0/4/4/4
2	2NG	C	1101	-	-	0/30/32/32	0/4/4/4
2	2NG	D	1101	-	-	0/30/32/32	0/4/4/4
2	2NG	E	1101	-	-	0/30/32/32	0/4/4/4
2	2NG	E	1102	-	-	0/30/32/32	0/4/4/4
2	2NG	F	1101	-	-	0/30/32/32	0/4/4/4

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1102	2NG	CBG-NBL	-7.06	1.31	1.45
2	D	1101	2NG	CBG-NBL	-6.61	1.32	1.45
2	C	1101	2NG	CBG-NBL	-6.42	1.33	1.45
2	A	1101	2NG	CBG-NBL	-6.17	1.33	1.45
2	B	1101	2NG	CBG-NBL	-6.12	1.33	1.45

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	1101	2NG	CAU-CBJ-CBH	-4.53	117.82	123.63
2	A	1101	2NG	CAU-CBJ-CBH	-3.83	118.72	123.63
2	C	1101	2NG	CAU-CBJ-CBH	-3.74	118.84	123.63
2	E	1101	2NG	CAU-CBJ-CBH	-3.73	118.84	123.63
2	B	1101	2NG	CAU-CBJ-CBH	-3.61	119.00	123.63

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	2NG	5	0
2	B	1101	2NG	2	0
2	C	1101	2NG	2	0
2	D	1101	2NG	1	0
2	E	1101	2NG	3	0
2	E	1102	2NG	2	0
2	F	1101	2NG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	477/515 (92%)	0.26	15 (3%) 49 56	20, 33, 74, 113	6 (1%)
1	B	482/515 (93%)	-0.03	4 (0%) 86 89	21, 36, 66, 101	8 (1%)
1	C	478/515 (92%)	0.04	10 (2%) 64 70	23, 37, 75, 106	9 (1%)
1	D	479/515 (93%)	0.00	9 (1%) 67 73	25, 39, 71, 101	7 (1%)
1	E	477/515 (92%)	0.24	15 (3%) 49 56	27, 44, 88, 111	5 (1%)
1	F	473/515 (91%)	0.54	30 (6%) 21 27	33, 54, 94, 116	7 (1%)
All	All	2866/3090 (92%)	0.18	83 (2%) 52 59	20, 40, 80, 116	42 (1%)

The worst 5 of 83 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	462	VAL	6.4
1	F	456	ARG	5.2
1	F	475	VAL	5.1
1	F	474	VAL	5.1
1	F	463	SER	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	2NG	F	1101	40/40	0.90	0.29	1.45	62,84,112,117	0
2	2NG	A	1101	40/40	0.92	0.22	1.42	48,76,111,114	0
2	2NG	C	1101	40/40	0.95	0.18	1.14	40,71,96,117	0
2	2NG	E	1101	40/40	0.94	0.23	1.08	59,88,103,111	0
2	2NG	B	1101	40/40	0.97	0.15	0.63	30,55,80,91	0
2	2NG	D	1101	40/40	0.96	0.15	0.05	39,61,93,100	0
2	2NG	E	1102	40/40	0.92	0.17	-0.31	29,48,70,90	0
3	MG	D	1102	1/1	0.95	0.11	-0.94	50,50,50,50	0
3	MG	E	1103	1/1	0.69	0.11	-1.33	48,48,48,48	0
3	MG	A	1102	1/1	0.97	0.15	-1.52	34,34,34,34	0
3	MG	B	1102	1/1	0.98	0.10	-2.03	31,31,31,31	0
3	MG	F	1102	1/1	0.82	0.08	-2.62	56,56,56,56	0
3	MG	C	1102	1/1	0.95	0.07	-6.40	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.