



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2017 – 04:34 AM EDT

PDB ID : 3NS7  
Title : Succinic Acid Amides as P2-P3 Replacements for Inhibitors of Interleukin-1beta Converting Enzyme (ICE or Caspase 1)  
Authors : Galatsis, P.  
Deposited on : unknown  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	
2	B	88	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Caspase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	1	0	0
			1277	801	222	243	11			

- Molecule 2 is a protein called Caspase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	88	Total	C	N	O	S	0	0	0
			719	457	126	129	7			

- Molecule 3 is (3S)-4-hydroxy-3-{[(2S)-4-{[2-(2-methyl-1H-benzimidazol-1-yl)ethyl]amino}-2-(1-methylethyl)-4-oxobutanoyl]amino}butanoic acid (three-letter code: 3NS) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>4</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			30	21	4	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total 32	O 32	0	0
4	B	15	Total 15	O 15	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Caspase-1



#### • Molecule 2: Caspase-1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.95Å 63.95Å 158.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.60 24.89 – 2.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (25.00-2.60) 97.5 (24.89-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.85 (at 2.60Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.188 , (Not available) 0.281 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	32.7	Xtriage
Anisotropy	0.349	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	2073	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 3NS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	1/1299 (0.1%)	1.29	1/1749 (0.1%)
2	B	0.85	0/739	1.34	1/995 (0.1%)
All	All	0.80	1/2038 (0.0%)	1.31	2/2744 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
2	B	0	2
All	All	0	7

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	297	ASP	C-OXT	6.32	1.35	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	391	ARG	NE-CZ-NH1	-6.21	117.19	120.30
1	A	163	ARG	NE-CZ-NH2	-5.30	117.65	120.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	137	SER	Peptide
1	A	150	ALA	Peptide
1	A	178	ARG	Sidechain
1	A	240	ARG	Sidechain
1	A	291	GLY	Peptide
2	B	362	CYS	Peptide
2	B	396	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1277	0	1288	27	0
2	B	719	0	693	17	0
3	B	30	0	28	2	0
4	A	32	0	0	7	3
4	B	15	0	0	3	0
All	All	2073	0	2009	40	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:385:GLN:HA	4:B:15:HOH:O	1.31	1.28
1:A:158:LYS:HG2	4:A:41:HOH:O	1.67	0.93
1:A:258:LEU:HB2	4:A:7:HOH:O	1.71	0.89
1:A:206:LEU:HG	1:A:210:ASP:HB2	1.68	0.74
2:B:362:CYS:HB2	2:B:402:PRO:HD2	1.72	0.72
2:B:327:PHE:O	4:B:29:HOH:O	2.10	0.69
1:A:155:ILE:HD13	4:A:24:HOH:O	1.93	0.67
2:B:393:THR:N	4:B:29:HOH:O	2.24	0.65
1:A:273:LEU:HB3	1:A:278:LYS:HE3	1.79	0.63
1:A:163:ARG:HD2	1:A:197:GLY:O	1.98	0.63
2:B:383:ARG:HD3	3:B:1:3NS:C18	2.33	0.59
1:A:258:LEU:HD23	1:A:282:ILE:HD11	1.84	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:HIS:HA	1:A:252:VAL:O	2.05	0.56
1:A:148:LYS:NZ	4:A:6:HOH:O	2.38	0.56
1:A:157:ASP:O	1:A:161:ARG:HG3	2.06	0.56
1:A:258:LEU:CB	4:A:7:HOH:O	2.41	0.56
1:A:235:MET:HA	1:A:283:GLN:O	2.07	0.55
2:B:341:ARG:HA	2:B:347:SER:HA	1.89	0.53
1:A:167:ILE:HA	1:A:233:VAL:HG13	1.92	0.51
1:A:155:ILE:HD11	2:B:398:PHE:HE2	1.77	0.50
1:A:170:ASN:O	1:A:179:ARG:HD3	2.14	0.48
1:A:286:ARG:HH22	2:B:390:GLU:CD	2.16	0.48
1:A:268:LYS:HE3	1:A:269:ASN:OD1	2.15	0.47
2:B:357:MET:O	2:B:361:ALA:HB2	2.16	0.45
2:B:366:VAL:HG22	2:B:398:PHE:HB3	1.98	0.45
2:B:342:HIS:HA	2:B:343:PRO:HD2	1.87	0.45
1:A:176:ILE:HB	1:A:177:PRO:HD2	1.98	0.45
1:A:279:VAL:HB	2:B:328:ILE:HG23	1.99	0.45
1:A:155:ILE:CD1	4:A:24:HOH:O	2.57	0.44
2:B:365:ASP:O	2:B:369:ILE:HG13	2.18	0.44
2:B:385:GLN:O	2:B:387:PRO:HD3	2.18	0.43
1:A:139:GLU:HA	4:A:47:HOH:O	2.19	0.43
1:A:185:ASP:HA	2:B:350:ILE:HG21	2.01	0.43
1:A:212:THR:O	1:A:216:GLU:HB2	2.20	0.42
1:A:204:LYS:O	1:A:206:LEU:HD13	2.19	0.41
1:A:139:GLU:HB2	1:A:140:GLU:H	1.74	0.41
1:A:147:GLN:HG2	1:A:148:LYS:HD3	2.02	0.41
2:B:379:GLN:HA	2:B:380:PRO:HD3	1.95	0.41
1:A:258:LEU:HD21	1:A:262:PHE:HE1	1.86	0.41
2:B:341:ARG:NH1	3:B:1:3NS:O2	2.55	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2:HOH:O	4:A:39:HOH:O[4_555]	0.37	1.83
4:A:1:HOH:O	4:A:45:HOH:O[6_455]	0.45	1.75
4:A:6:HOH:O	4:A:6:HOH:O[8_665]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/162 (99%)	136 (85%)	20 (12%)	4 (2%)	6	11
2	B	86/88 (98%)	75 (87%)	8 (9%)	3 (4%)	4	6
All	All	246/250 (98%)	211 (86%)	28 (11%)	7 (3%)	6	9

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	281	ILE
1	A	140	GLU
1	A	209	SER
2	B	343	PRO
2	B	401	PHE
1	A	150	ALA
2	B	328	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/145 (100%)	123 (85%)	22 (15%)	3	5
2	B	79/79 (100%)	71 (90%)	8 (10%)	9	16
All	All	224/224 (100%)	194 (87%)	30 (13%)	4	8

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	139	GLU
1	A	147	GLN
1	A	148	LYS
1	A	157	ASP
1	A	164	LEU
1	A	180	THR
1	A	184	VAL
1	A	190	THR
1	A	192	LEU
1	A	193	LEU
1	A	203	LYS
1	A	206	LEU
1	A	209	SER
1	A	210	ASP
1	A	226	THR
1	A	233	VAL
1	A	250	GLU
1	A	256	LEU
1	A	259	ASN
1	A	273	LEU
1	A	275	ASP
1	A	295	PHE
2	B	343	PRO
2	B	348	VAL
2	B	353	LEU
2	B	362	CYS
2	B	388	THR
2	B	390	GLU
2	B	392	VAL
2	B	393	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	3NS	B	1	1	26,31,31	0.92	1 (3%)	28,42,42	0.99	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3NS	B	1	1	-	0/26/28/28	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1	3NS	C15-C14	-2.66	1.37	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1	3NS	C4-C3-N1	-2.60	112.05	115.92
3	B	1	3NS	C5-C7-N2	-2.26	113.28	116.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1	3NS	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/162 (100%)	0.57	12 (7%)	15	11	5, 22, 58, 83	1 (0%)
2	B	88/88 (100%)	0.21	3 (3%)	46	38	2, 15, 34, 55	0
All	All	250/250 (100%)	0.44	15 (6%)	23	17	2, 19, 53, 83	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	252	VAL	5.0
1	A	139	GLU	4.8
1	A	147	GLN	3.6
1	A	148	LYS	3.4
1	A	241	GLU	3.1
1	A	172	GLU	2.7
2	B	381	ASP	2.7
1	A	250	GLU	2.6
1	A	145	TRP	2.5
2	B	379	GLN	2.4
1	A	150	ALA	2.4
1	A	149	SER	2.1
2	B	317	ALA	2.1
1	A	178	ARG	2.1
1	A	159	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	3NS	B	1	30/30	0.80	0.29	1.75	15,20,28,29	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.