



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 10, 2017 – 06:36 PM EDT

PDB ID : 5NTN  
Title : Structural states of RORgt: X-ray elucidation of molecular mechanisms and binding interactions for natural and synthetic compounds  
Authors : Kallen, J.  
Deposited on : unknown  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div>0.1%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>
1	B	257	<div> <div>2%</div> <div> <div></div> <div>93%</div> <div>5%</div> </div> </div>
1	C	257	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>5%</div> <div>5%</div> </div> </div>
1	D	257	<div> <div>7%</div> <div> <div></div> <div>91%</div> <div>5%</div> </div> </div>
2	P	20	<div> <div></div> <div> <div>40%</div> <div>60%</div> </div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	Q	20	 40% 60%
2	R	20	 40% 60%
2	S	20	 35% 10% 55%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear receptor ROR-gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	3	0
			2014	1285	357	359	13			
1	B	244	Total	C	N	O	S	0	0	0
			1998	1275	356	354	13			
1	C	245	Total	C	N	O	S	0	1	0
			2008	1281	357	357	13			
1	D	244	Total	C	N	O	S	0	4	0
			2013	1286	357	357	13			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	262	GLY	-	expression tag	UNP P51449
A	455	SER	CYS	engineered mutation	UNP P51449
B	262	GLY	-	expression tag	UNP P51449
B	455	SER	CYS	engineered mutation	UNP P51449
C	262	GLY	-	expression tag	UNP P51449
C	455	SER	CYS	engineered mutation	UNP P51449
D	262	GLY	-	expression tag	UNP P51449
D	455	SER	CYS	engineered mutation	UNP P51449

- Molecule 2 is a protein called Nuclear receptor-interacting protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	8	Total	C	N	O	0	0	0
			60	41	9	10			
2	Q	8	Total	C	N	O	0	0	0
			60	41	9	10			
2	R	8	Total	C	N	O	0	0	0
			60	41	9	10			
2	S	9	Total	C	N	O	0	0	0
			70	47	12	11			

- Molecule 3 is (5R,10S,13R,14R,17R)-17-((R,E)-7-hydroxy-6-methylhept-5-en-2-yl)-4,4,10,13,14-pentamethyl-1,2,5,6,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthrene-3,7(4H)-dione (three-letter code: 98H) (formula: C<sub>30</sub>H<sub>46</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			33	30	3		
3	B	1	Total	C	O	0	0
			33	30	3		
3	C	1	Total	C	O	0	0
			33	30	3		
3	D	1	Total	C	O	0	0
			33	30	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	218	Total	O	0	0
			218	218		
4	B	176	Total	O	0	0
			176	176		
4	C	128	Total	O	0	0
			128	128		
4	D	122	Total	O	0	0
			122	122		
4	P	2	Total	O	0	0
			2	2		
4	Q	1	Total	O	0	0
			1	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	S	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 1: Nuclear receptor ROR-gamma



- Molecule 1: Nuclear receptor ROR-gamma




- Molecule 2: Nuclear receptor-interacting protein 1




- Molecule 2: Nuclear receptor-interacting protein 1

Chain Q:  40% 60%




- Molecule 2: Nuclear receptor-interacting protein 1

Chain R:  40% 60%



- Molecule 2: Nuclear receptor-interacting protein 1

Chain S:  35% 10% 55%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.24Å 69.68Å 97.21Å 90.00° 107.94° 90.00°	Depositor
Resolution (Å)	19.92 – 1.90 19.92 – 1.90	Depositor EDS
% Data completeness (in resolution range)	89.3 (19.92-1.90) 89.4 (19.92-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.32 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, $R_{free}$	0.200 , 0.225 0.199 , 0.224	Depositor DCC
$R_{free}$ test set	3880 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.4	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9068	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.41 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5573e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 98H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	0/2067	0.48	0/2785
1	B	0.38	0/2042	0.46	0/2751
1	C	0.37	0/2055	0.47	0/2769
1	D	0.38	0/2069	0.47	0/2788
2	P	0.41	0/59	0.52	0/79
2	Q	0.37	0/59	0.45	0/79
2	R	0.39	0/59	0.55	0/79
2	S	0.36	0/70	0.45	0/94
All	All	0.38	0/8480	0.47	0/11424

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2014	0	2021	4	0
1	B	1998	0	1999	3	0
1	C	2008	0	2011	6	0
1	D	2013	0	2024	5	0
2	P	60	0	72	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	Q	60	0	72	0	0
2	R	60	0	72	0	0
2	S	70	0	79	2	0
3	A	33	0	0	0	0
3	B	33	0	0	0	0
3	C	33	0	0	0	0
3	D	33	0	0	0	0
4	A	218	0	0	0	0
4	B	176	0	0	1	0
4	C	128	0	0	0	0
4	D	122	0	0	0	0
4	P	2	0	0	0	0
4	Q	1	0	0	0	0
4	S	6	0	0	1	0
All	All	9068	0	8350	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:323:HIS:HE1	1:C:379:GLU:OE2	1.76	0.69
1:B:264:TYR:N	4:B:702:HOH:O	2.39	0.55
1:A:383:GLY:HA3	1:A:387:LEU:HD11	1.92	0.52
1:C:487:GLN:HG3	1:C:494:VAL:HG21	1.92	0.52
1:D:383:GLY:HA3	1:D:387:LEU:HD11	1.93	0.51
2:S:499:THR:N	4:S:601:HOH:O	2.47	0.47
1:D:478:GLN:O	1:D:482:ARG:HG2	2.16	0.46
1:C:412:PHE:HB3	1:C:417:ILE:HG13	1.97	0.45
1:B:346:GLN:O	1:B:350:ILE:HG12	2.17	0.44
1:D:490:HIS:HB3	1:D:493[B]:VAL:HG22	1.99	0.43
1:C:383:GLY:HA3	1:C:387:LEU:HD11	2.01	0.43
1:D:277:VAL:HG22	1:D:338:LEU:HD11	2.00	0.43
1:C:496:ALA:HA	2:S:500:LEU:HD22	2.01	0.43
1:B:412:PHE:HB3	1:B:417:ILE:HG13	1.99	0.42
1:A:427:ASN:O	1:A:430:ARG:HG2	2.21	0.41
1:A:412:PHE:HB3	1:A:417:ILE:HG13	2.02	0.41
1:A:383:GLY:HA3	1:A:387:LEU:CD1	2.50	0.41
1:D:463:LEU:HA	1:D:466:LEU:HD13	2.02	0.41
1:C:452:HIS:NE2	1:C:456:LYS:HE3	2.35	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/257 (96%)	243 (99%)	3 (1%)	0	100	100
1	B	242/257 (94%)	239 (99%)	3 (1%)	0	100	100
1	C	244/257 (95%)	239 (98%)	4 (2%)	1 (0%)	38	26
1	D	246/257 (96%)	242 (98%)	4 (2%)	0	100	100
2	P	6/20 (30%)	6 (100%)	0	0	100	100
2	Q	6/20 (30%)	6 (100%)	0	0	100	100
2	R	6/20 (30%)	6 (100%)	0	0	100	100
2	S	7/20 (35%)	7 (100%)	0	0	100	100
All	All	1003/1108 (90%)	988 (98%)	14 (1%)	1 (0%)	55	45

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	458	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/229 (97%)	221 (100%)	1 (0%)	91	91

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	218/229 (95%)	218 (100%)	0	100	100
1	C	220/229 (96%)	218 (99%)	2 (1%)	82	82
1	D	222/229 (97%)	221 (100%)	1 (0%)	91	91
2	P	7/19 (37%)	7 (100%)	0	100	100
2	Q	7/19 (37%)	7 (100%)	0	100	100
2	R	7/19 (37%)	7 (100%)	0	100	100
2	S	8/19 (42%)	8 (100%)	0	100	100
All	All	911/992 (92%)	907 (100%)	4 (0%)	93	93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	404	SER
1	C	434	GLN
1	C	457	THR
1	D	450	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	GLN
1	B	322	HIS
1	C	322	HIS
1	C	323	HIS
1	D	322	HIS
1	D	460	GLN
2	S	502	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	98H	A	601	-	34,36,36	1.26	2 (5%)	52,59,59	1.02	4 (7%)
3	98H	B	601	-	34,36,36	1.21	2 (5%)	52,59,59	1.07	5 (9%)
3	98H	C	601	-	34,36,36	1.20	2 (5%)	52,59,59	1.08	6 (11%)
3	98H	D	601	-	34,36,36	1.30	2 (5%)	52,59,59	1.08	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	98H	A	601	-	-	0/12/87/87	0/4/4/4
3	98H	B	601	-	-	0/12/87/87	0/4/4/4
3	98H	C	601	-	-	0/12/87/87	0/4/4/4
3	98H	D	601	-	-	0/12/87/87	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	98H	C68-C19	2.05	1.58	1.54
3	C	601	98H	C37-C29	2.08	1.58	1.54
3	A	601	98H	C68-C19	2.12	1.58	1.54
3	B	601	98H	C15-C17	2.18	1.58	1.54
3	C	601	98H	C15-C17	2.18	1.58	1.54
3	D	601	98H	C68-C19	2.22	1.58	1.54

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	98H	C15-C17	2.37	1.58	1.54
3	D	601	98H	C15-C17	2.53	1.59	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	98H	C64-C20-C21	-2.38	104.80	113.64
3	A	601	98H	C64-C20-C21	-2.36	104.88	113.64
3	B	601	98H	C64-C20-C21	-2.25	105.27	113.64
3	D	601	98H	C64-C20-C21	-2.10	105.84	113.64
3	A	601	98H	C34-C37-C29	2.09	117.45	113.42
3	A	601	98H	C49-C52-C27	2.10	120.36	117.15
3	C	601	98H	C19-C17-C15	2.12	122.08	119.26
3	C	601	98H	C64-C20-C19	2.13	119.77	114.03
3	C	601	98H	C34-C37-C29	2.13	117.54	113.42
3	B	601	98H	C49-C30-C29	2.13	112.81	110.34
3	C	601	98H	C49-C30-C29	2.14	112.81	110.34
3	B	601	98H	C9-C12-C15	2.15	118.50	114.56
3	B	601	98H	C64-C20-C19	2.29	120.20	114.03
3	D	601	98H	C64-C20-C19	2.30	120.23	114.03
3	C	601	98H	C49-C52-C27	2.31	120.69	117.15
3	A	601	98H	C19-C17-C15	2.43	122.49	119.26
3	B	601	98H	C49-C52-C27	2.58	121.10	117.15
3	D	601	98H	C19-C17-C15	2.64	122.77	119.26
3	D	601	98H	C49-C52-C27	2.93	121.63	117.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	245/257 (95%)	-0.11	2 (0%) 86 87	14, 21, 31, 44	0
1	B	244/257 (94%)	0.08	5 (2%) 65 69	17, 26, 37, 43	0
1	C	245/257 (95%)	0.28	12 (4%) 30 34	17, 28, 52, 68	0
1	D	244/257 (94%)	0.33	17 (6%) 17 19	19, 31, 45, 53	0
2	P	8/20 (40%)	-0.15	0 100 100	25, 28, 32, 33	0
2	Q	8/20 (40%)	0.39	0 100 100	29, 34, 39, 40	0
2	R	8/20 (40%)	0.51	0 100 100	32, 35, 41, 42	0
2	S	9/20 (45%)	-0.04	0 100 100	26, 28, 33, 34	0
All	All	1011/1108 (91%)	0.15	36 (3%) 43 47	14, 27, 43, 68	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	464	ALA	4.8
1	D	302	ARG	4.6
1	D	294	ARG	3.9
1	C	458	HIS	3.8
1	D	272	HIS	3.5
1	D	484	GLN	3.3
1	C	294	ARG	3.3
1	A	469	LYS	3.2
1	C	469	LYS	3.2
1	D	371	ALA	2.8
1	C	272	HIS	2.8
1	D	437	ARG	2.7
1	D	464	ALA	2.7
1	D	288	ARG	2.7
1	C	264	TYR	2.6
1	D	429	HIS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	268	THR	2.6
1	D	493[A]	VAL	2.6
1	C	456	LYS	2.5
1	C	461	SER	2.5
1	D	290	GLU	2.5
1	D	372	ASP	2.5
1	D	264	TYR	2.5
1	C	437	ARG	2.4
1	D	389	ARG	2.4
1	B	290	GLU	2.3
1	B	294	ARG	2.3
1	B	469	LYS	2.3
1	C	429	HIS	2.3
1	B	470	GLY	2.2
1	B	473	ARG	2.2
1	D	461	SER	2.1
1	D	268	THR	2.1
1	D	466	LEU	2.1
1	A	303	GLU	2.0
1	C	457	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	98H	D	601	33/33	0.91	0.12	1.11	25,27,31,31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	98H	C	601	33/33	0.92	0.11	0.57	20,21,26,26	0
3	98H	B	601	33/33	0.93	0.11	0.41	20,21,25,26	0
3	98H	A	601	33/33	0.95	0.11	0.23	18,18,20,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.