



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 11:21 pm GMT

PDB ID : 2NUA
Title : C123aV Mutant of E. coli Succinyl-CoA Synthetase
Authors : Fraser, M.E.
Deposited on : 2006-11-08
Resolution : 2.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28683
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2395 (3.00-2.92)
Clashscore	112137	2773 (3.00-2.92)
Ramachandran outliers	110173	2680 (3.00-2.92)
Sidechain outliers	110143	2683 (3.00-2.92)
RSRZ outliers	101464	2421 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	<div> <div>2%</div> <div>64%</div> <div>34%</div> <div>•</div> </div>
1	D	288	<div> <div>5%</div> <div>65%</div> <div>32%</div> <div>•</div> </div>
2	B	388	<div> <div>6%</div> <div>54%</div> <div>41%</div> <div>•</div> </div>
2	E	388	<div> <div>10%</div> <div>57%</div> <div>37%</div> <div>5%</div> <div>•</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10105 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinyl-CoA ligase [ADP-forming] subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	0	0	0
			2070	1315	346	399	10			
1	D	287	Total	C	N	O	S	0	0	0
			2070	1315	346	399	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	VAL	CYS	ENGINEERED	UNP P0AGE9
D	123	VAL	CYS	ENGINEERED	UNP P0AGE9

- Molecule 2 is a protein called Succinyl-CoA synthetase beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	388	Total	C	N	O	S	0	0	0
			2908	1836	509	550	13			
2	E	385	Total	C	N	O	S	0	0	0
			2885	1823	505	544	13			

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	D	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
5	B	1	Total 18	C 11	N 2	O 4	S 1		0	0
5	D	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
5	E	1	Total 18	C 11	N 2	O 4	S 1		0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	4	Total	O	0	0
			4	4		
6	B	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha




- Molecule 1: Succinyl-CoA ligase [ADP-forming] subunit alpha



- Molecule 2: Succinyl-CoA synthetase beta chain





• Molecule 2: Succinyl-CoA synthetase beta chain

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	97.51Å 97.51Å 390.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.51 – 2.95 97.51 – 2.95	Depositor EDS
% Data completeness (in resolution range)	90.8 (97.51-2.95) 90.8 (97.51-2.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.96Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.221 , 0.264 0.231 , 0.273	Depositor DCC
R_{free} test set	4191 reflections (11.28%)	DCC
Wilson B-factor (Å ²)	75.3	Xtriage
Anisotropy	0.181	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10105	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, PO4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/2104	0.66	0/2851
1	D	0.37	0/2104	0.66	0/2851
2	B	0.41	0/2950	0.64	0/3989
2	E	0.39	0/2927	0.64	0/3961
All	All	0.39	0/10085	0.65	0/13652

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2070	0	2133	84	1
1	D	2070	0	2133	88	1
2	B	2908	0	2962	162	0
2	E	2885	0	2940	157	0
3	A	5	0	0	0	0
3	D	5	0	0	0	0
4	A	5	0	0	0	0
4	B	10	0	0	0	0
4	E	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	48	0	32	4	0
5	B	18	0	20	4	0
5	D	48	0	32	3	0
5	E	18	0	20	6	0
6	A	4	0	0	1	0
6	B	1	0	0	0	0
All	All	10105	0	10272	473	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 473 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:LYS:HD3	1:A:240:LYS:H	0.96	1.10
2:E:275:ILE:HD13	2:E:374:LEU:HD12	1.46	0.97
1:A:240:LYS:HD3	1:A:240:LYS:N	1.81	0.96
1:A:240:LYS:HA	1:A:251:ILE:HG23	1.48	0.96
2:B:275:ILE:HD13	2:B:374:LEU:HD12	1.49	0.95

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:113:GLU:OE2	1:D:113:GLU:OE2[7_555]	2.06	0.14
1:A:113:GLU:OE2	1:A:113:GLU:OE2[7_555]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/288 (99%)	267 (94%)	17 (6%)	1 (0%)	38	75
1	D	285/288 (99%)	257 (90%)	22 (8%)	6 (2%)	8	35
2	B	386/388 (100%)	344 (89%)	32 (8%)	10 (3%)	6	28
2	E	383/388 (99%)	345 (90%)	30 (8%)	8 (2%)	8	35
All	All	1339/1352 (99%)	1213 (91%)	101 (8%)	25 (2%)	9	38

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	257	GLY
2	B	293	ALA
1	D	249	ALA
1	D	253	GLY
1	D	256	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/218 (100%)	211 (97%)	6 (3%)	49	80
1	D	217/218 (100%)	210 (97%)	7 (3%)	44	78
2	B	298/298 (100%)	282 (95%)	16 (5%)	26	61
2	E	296/298 (99%)	280 (95%)	16 (5%)	26	61
All	All	1028/1032 (100%)	983 (96%)	45 (4%)	33	69

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	374	LEU
1	D	240	LYS
2	E	350	GLU
1	D	117	ARG
1	D	242	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	279	HIS
1	D	186	ASN
2	E	342	ASN
2	B	352	ASN
1	D	137	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	COA	A	1300	-	43,50,50	0.82	1 (2%)	48,75,75	1.28	5 (10%)
3	PO4	A	1600	-	4,4,4	1.29	0	6,6,6	0.38	0
4	SO4	A	1700	-	4,4,4	0.58	0	6,6,6	0.08	0
5	COA	B	1325	2	14,17,50	1.49	1 (7%)	16,22,75	4.20	11 (68%)
4	SO4	B	1400	-	4,4,4	0.38	0	6,6,6	0.10	0
4	SO4	B	1500	-	4,4,4	0.35	0	6,6,6	0.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	COA	D	1301	-	43,50,50	0.89	1 (2%)	48,75,75	1.45	5 (10%)
3	PO4	D	1601	-	4,4,4	1.38	0	6,6,6	0.41	0
5	COA	E	1325	2	14,17,50	1.30	1 (7%)	16,22,75	2.96	9 (56%)
4	SO4	E	1401	-	4,4,4	0.37	0	6,6,6	0.15	0
4	SO4	E	1501	-	4,4,4	0.35	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	COA	A	1300	-	-	0/44/64/64	0/3/3/3
3	PO4	A	1600	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1700	-	-	0/0/0/0	0/0/0/0
5	COA	B	1325	2	-	0/23/23/64	0/0/0/3
4	SO4	B	1400	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1500	-	-	0/0/0/0	0/0/0/0
5	COA	D	1301	-	-	0/44/64/64	0/3/3/3
3	PO4	D	1601	-	-	0/0/0/0	0/0/0/0
5	COA	E	1325	2	-	0/23/23/64	0/0/0/3
4	SO4	E	1401	-	-	0/0/0/0	0/0/0/0
4	SO4	E	1501	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	1301	COA	CCP-CBP	-3.11	1.48	1.52
5	A	1300	COA	CCP-CBP	-2.99	1.48	1.52
5	E	1325	COA	C6P-C5P	2.63	1.56	1.51
5	B	1325	COA	C6P-C5P	3.88	1.58	1.51

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1325	COA	CEP-CBP-CCP	-6.04	97.93	109.06
5	D	1301	COA	O5P-C5P-C6P	-4.38	113.78	122.01
5	B	1325	COA	O5P-C5P-N4P	-3.64	116.02	122.97
5	E	1325	COA	O9P-C9P-N8P	-3.60	116.15	123.07
5	E	1325	COA	O5P-C5P-N4P	-3.56	116.19	122.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1300	COA	4	0
5	B	1325	COA	4	0
5	D	1301	COA	3	0
5	E	1325	COA	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	287/288 (99%)	0.45	5 (1%) 70 51	35, 54, 91, 99	0
1	D	287/288 (99%)	0.67	13 (4%) 34 21	44, 69, 95, 100	0
2	B	388/388 (100%)	0.72	25 (6%) 20 11	35, 65, 93, 100	0
2	E	385/388 (99%)	0.77	39 (10%) 8 4	41, 70, 99, 100	0
All	All	1347/1352 (99%)	0.67	82 (6%) 22 13	35, 65, 97, 100	0

The worst 5 of 82 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	384	ALA	5.5
2	E	347	VAL	4.8
2	E	369	ILE	4.6
2	B	259	ILE	4.2
2	E	371	ALA	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	COA	B	1325	18/48	0.58	0.31	0.36	74,84,89,89	0
5	COA	A	1300	48/48	0.96	0.23	0.01	51,57,65,69	0
5	COA	D	1301	48/48	0.95	0.23	-0.01	49,59,65,73	0
4	SO4	E	1501	5/5	0.91	0.23	-0.06	98,99,100,100	0
5	COA	E	1325	18/48	0.58	0.25	-0.44	85,92,94,94	0
4	SO4	E	1401	5/5	0.92	0.16	-0.61	90,90,91,91	0
4	SO4	A	1700	5/5	0.92	0.18	-1.07	98,100,100,100	0
4	SO4	B	1500	5/5	0.94	0.15	-1.48	99,99,100,100	0
3	PO4	A	1600	5/5	0.97	0.15	-1.71	71,71,72,73	0
3	PO4	D	1601	5/5	0.97	0.14	-1.82	78,79,79,80	0
4	SO4	B	1400	5/5	0.96	0.13	-1.98	83,84,84,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.