



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 06:05 am GMT

PDB ID : 2NVX  
Title : RNA polymerase II elongation complex in 5 mM Mg+2 with 2'-dUTP  
Authors : Wang, D.; Bushnell, D.A.; Westover, K.D.; Kaplan, C.D.; Kornberg, R.D.  
Deposited on : 2006-11-13  
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1026 (3.74-3.46)
Clashscore	112137	1036 (3.70-3.50)
Ramachandran outliers	110173	1030 (3.72-3.48)
Sidechain outliers	110143	1030 (3.72-3.48)
RSRZ outliers	101464	1051 (3.74-3.46)
RNA backbone	2435	1002 (4.30-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	R	10	
2	N	14	
3	T	28	
4	A	1733	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	B	1224	
6	C	318	
7	E	215	
8	F	155	
9	H	146	
10	I	122	
11	J	70	
12	K	120	
13	L	70	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
16	DUT	B	1308[A]	-	-	-	X

## 2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 29436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(\*AP\*UP\*CP\*GP\*AP\*GP\*AP\*GP\*GP\*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	R	10	Total	C	N	O	P	0	0	0
			216	98	45	64	9			

- Molecule 2 is a DNA chain called 5'-D(\*CP\*TP\*GP\*CP\*TP\*TP\*AP\*TP\*CP\*GP\*GP\*TP\*AP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	14	Total	C	N	O	P	0	0	0
			284	137	49	85	13			

- Molecule 3 is a DNA chain called 28-MER DNA template strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	28	Total	C	N	O	P	0	0	0
			566	271	104	164	27			

- Molecule 4 is a protein called DNA-directed RNA polymerase II largest subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	1402	Total	C	N	O	S	0	0	0
			11028	6950	1934	2083	61			

- Molecule 5 is a protein called DNA-directed RNA polymerase II 140 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	B	1114	Total	C	N	O	S	0	0	0
			8856	5605	1553	1644	54			

- Molecule 6 is a protein called DNA-directed RNA polymerase II 45 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	C	266	Total	C	N	O	S	0	0	0
			2095	1317	348	417	13			

- Molecule 7 is a protein called DNA-directed RNA polymerases I, II, and III 27 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III 23 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	F	88	Total	C	N	O	S	0	0	0
			712	455	120	134	3			

- Molecule 9 is a protein called DNA-directed RNA polymerases I, II, and III 14.5 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	H	134	Total	C	N	O	S	0	0	0
			1076	678	181	212	5			

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	I	119	Total	C	N	O	S	0	0	0
			971	596	179	186	10			

- Molecule 11 is a protein called DNA-directed RNA polymerases I/II/III subunit 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 12 is a protein called DNA-directed RNA polymerase II 13.6 kDa polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 13 is a protein called DNA-directed RNA polymerases I, II, and III 7.7 kDa polypep-

tide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	L	46	Total	C	N	O	S	0	0	0
			364	224	72	64	4			


- Molecule 14 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	J	1	Total	Zn	0	0
			1	1		
14	B	1	Total	Zn	0	0
			1	1		
14	I	2	Total	Zn	0	0
			2	2		
14	C	1	Total	Zn	0	0
			1	1		
14	A	2	Total	Zn	0	0
			2	2		
14	L	1	Total	Zn	0	0
			1	1		

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	1	Total	Mg	0	0
			1	1		

- Molecule 16 is DEOXYURIDINE-5'-TRIPHOSPHATE (three-letter code: DUT) (formula: C<sub>9</sub>H<sub>15</sub>N<sub>2</sub>O<sub>14</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
16	B	1	Total	C	N	O	P	0	1
			56	18	4	28	6		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-R(\*AP\*UP\*CP\*GP\*AP\*GP\*AP\*GP\*GP\*A)-3'

Chain R: 




- Molecule 2: 5'-D(\*CP\*TP\*GP\*CP\*TP\*TP\*AP\*TP\*CP\*GP\*GP\*TP\*AP\*G)-3'

Chain N: 





- Molecule 3: 28-MER DNA template strand

Chain T: 



- Molecule 4: DNA-directed RNA polymerase II largest subunit

Chain A: 







TYR  
SER  
PRO  
LYS  
GLN  
ASP  
GLU  
GLN  
LYS  
HIS  
ASN  
GLU  
ASN  
GLU  
ASN  
ARG


• Molecule 5: DNA-directed RNA polymerase II 140 kDa polypeptide

Chain B: 







• Molecule 6: DNA-directed RNA polymerase II 45 kDa polypeptide




• Molecule 7: DNA-directed RNA polymerases I, II, and III 27 kDa polypeptide




• Molecule 8: DNA-directed RNA polymerases I, II, and III 23 kDa polypeptide



• Molecule 9: DNA-directed RNA polymerases I, II, and III 14.5 kDa polypeptide






- Molecule 10: DNA-directed RNA polymerase II subunit 9



- Molecule 11: DNA-directed RNA polymerases I/II/III subunit 10



- Molecule 12: DNA-directed RNA polymerase II 13.6 kDa polypeptide



- Molecule 13: DNA-directed RNA polymerases I, II, and III 7.7 kDa polypeptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	168.72Å 222.41Å 193.07Å 90.00° 101.29° 90.00°	Depositor
Resolution (Å)	50.00 – 3.60 44.25 – 3.60	Depositor EDS
% Data completeness (in resolution range)	91.5 (50.00-3.60) 91.6 (44.25-3.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 3.57Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.285 , 0.304 0.282 , 0.296	Depositor DCC
$R_{free}$ test set	3721 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	104.2	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 59.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	29436	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	109.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DUT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	R	1.16	1/243 (0.4%)	1.75	6/378 (1.6%)
2	N	0.74	0/317	1.27	0/488
3	T	1.18	4/634 (0.6%)	1.76	18/975 (1.8%)
4	A	0.68	8/11224 (0.1%)	0.67	2/15176 (0.0%)
5	B	0.82	11/9027 (0.1%)	0.73	3/12172 (0.0%)
6	C	0.71	0/2133	0.69	0/2891
7	E	0.58	1/1788 (0.1%)	0.61	0/2406
8	F	0.60	0/724	0.72	0/977
9	H	0.57	0/1094	0.65	0/1480
10	I	0.75	2/989 (0.2%)	0.68	0/1331
11	J	0.69	0/541	0.68	0/727
12	K	0.66	0/937	0.65	0/1265
13	L	0.70	0/366	0.78	0/485
All	All	0.74	27/30017 (0.1%)	0.76	29/40751 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	A	0	32
5	B	0	12
8	F	0	1
All	All	0	45

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	764	SER	CB-OG	13.86	1.60	1.42
5	B	490	SER	CB-OG	9.83	1.55	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	879	GLU	CD-OE1	7.99	1.34	1.25
3	T	19	DT	C5-C7	7.74	1.54	1.50
5	B	404	LYS	CE-NZ	7.54	1.67	1.49

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	19	DT	C4-C5-C7	10.26	125.16	119.00
3	T	19	DT	C6-C5-C7	-9.91	116.95	122.90
3	T	19	DT	N3-C2-O2	-9.91	116.36	122.30
3	T	16	DC	O4'-C1'-N1	9.72	114.80	108.00
3	T	28	DT	O4'-C1'-N1	8.42	113.89	108.00

There are no chirality outliers.

5 of 45 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	A	153	PRO	Peptide
4	A	70	CYS	Peptide
4	A	71	GLN	Peptide
4	A	79	GLY	Peptide
4	A	80	HIS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	R	216	0	109	7	0
2	N	284	0	161	1	0
3	T	566	0	316	11	0
4	A	11028	0	11120	901	0
5	B	8856	0	8897	724	0
6	C	2095	0	2051	80	0
7	E	1752	0	1776	33	0
8	F	712	0	738	47	0
9	H	1076	0	1052	40	0
10	I	971	0	930	21	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	J	532	0	542	41	0
12	K	919	0	929	20	0
13	L	364	0	387	20	0
14	A	2	0	0	1	0
14	B	1	0	0	1	0
14	C	1	0	0	0	0
14	I	2	0	0	0	0
14	J	1	0	0	0	0
14	L	1	0	0	0	0
15	A	1	0	0	0	0
16	B	56	0	22	1	0
All	All	29436	0	29030	1745	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 1745 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:404:LYS:NZ	5:B:404:LYS:CE	1.67	1.52
4:A:1287:TYR:CD2	4:A:1305:VAL:HB	1.59	1.36
4:A:1229:SER:HB2	4:A:1236:LEU:CD1	1.59	1.33
4:A:1229:SER:CB	4:A:1236:LEU:HD12	1.64	1.27
5:B:203:PHE:CA	5:B:204:ILE:HD12	1.63	1.27

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	A	1392/1733 (80%)	1118 (80%)	182 (13%)	92 (7%)	<b>1</b> <b>20</b>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	B	1096/1224 (90%)	897 (82%)	124 (11%)	75 (7%)	1	19
6	C	264/318 (83%)	229 (87%)	28 (11%)	7 (3%)	6	42
7	E	212/215 (99%)	190 (90%)	15 (7%)	7 (3%)	4	38
8	F	86/155 (56%)	75 (87%)	6 (7%)	5 (6%)	2	23
9	H	130/146 (89%)	105 (81%)	18 (14%)	7 (5%)	2	24
10	I	117/122 (96%)	96 (82%)	18 (15%)	3 (3%)	6	43
11	J	63/70 (90%)	53 (84%)	8 (13%)	2 (3%)	5	38
12	K	112/120 (93%)	107 (96%)	5 (4%)	0	100	100
13	L	44/70 (63%)	29 (66%)	13 (30%)	2 (4%)	3	29
All	All	3516/4173 (84%)	2899 (82%)	417 (12%)	200 (6%)	2	23

5 of 200 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	A	55	ASP
4	A	56	PRO
4	A	74	MET
4	A	157	ASP
4	A	158	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	1225/1520 (81%)	1128 (92%)	97 (8%)	14	51
5	B	967/1061 (91%)	886 (92%)	81 (8%)	13	49
6	C	234/274 (85%)	211 (90%)	23 (10%)	9	42
7	E	196/197 (100%)	181 (92%)	15 (8%)	15	52
8	F	78/137 (57%)	74 (95%)	4 (5%)	28	66
9	H	118/128 (92%)	107 (91%)	11 (9%)	10	45
10	I	113/116 (97%)	103 (91%)	10 (9%)	12	47

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
11	J	60/65 (92%)	52 (87%)	8 (13%)	4	28
12	K	99/102 (97%)	87 (88%)	12 (12%)	6	32
13	L	40/57 (70%)	36 (90%)	4 (10%)	9	41
All	All	3130/3657 (86%)	2865 (92%)	265 (8%)	12	48

5 of 265 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	B	388	CYS
5	B	806	THR
11	J	13	VAL
5	B	465	ASN
5	B	567	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 68 such sidechains are listed below:

Mol	Chain	Res	Type
5	B	363	HIS
5	B	592	ASN
9	H	137	GLN
5	B	366	GLN
5	B	494	HIS

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	R	8/10 (80%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 11 ligands modelled in this entry, 9 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
16	DUT	B	1308[A]	-	22,29,29	0.84	2 (9%)	24,45,45	2.01	3 (12%)
16	DUT	B	1308[B]	15	22,29,29	0.71	0	24,45,45	2.09	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	DUT	B	1308[A]	-	-	0/18/34/34	0/2/2/2
16	DUT	B	1308[B]	15	-	0/18/34/34	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	B	1308[A]	DUT	C4-N3	2.13	1.36	1.33
16	B	1308[A]	DUT	PG-O3B	2.33	1.63	1.60

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	B	1308[B]	DUT	C2'-C1'-N1	-2.32	108.75	114.23
16	B	1308[A]	DUT	O3G-PG-O2G	2.03	115.81	107.61
16	B	1308[A]	DUT	O4'-C1'-N1	2.05	111.24	107.78
16	B	1308[B]	DUT	O4'-C1'-N1	3.04	112.90	107.78
16	B	1308[B]	DUT	C4-N3-C2	7.94	120.95	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	B	1308[A]	DUT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	R	10/10 (100%)	-0.58	0 <span>100</span> <span>100</span>	85, 109, 154, 170	0
2	N	14/14 (100%)	1.46	3 (21%) <span>1</span> <span>1</span>	178, 182, 198, 202	0
3	T	28/28 (100%)	0.74	7 (25%) <span>1</span> <span>1</span>	93, 173, 189, 192	0
4	A	1402/1733 (80%)	-0.04	21 (1%) <span>74</span> <span>61</span>	75, 103, 150, 163	0
5	B	1114/1224 (91%)	-0.04	17 (1%) <span>74</span> <span>61</span>	26, 102, 133, 143	0
6	C	266/318 (83%)	-0.23	1 (0%) <span>92</span> <span>87</span>	81, 100, 128, 144	0
7	E	214/215 (99%)	0.07	5 (2%) <span>61</span> <span>46</span>	91, 131, 165, 167	0
8	F	88/155 (56%)	-0.12	0 <span>100</span> <span>100</span>	84, 104, 130, 137	0
9	H	134/146 (91%)	0.12	3 (2%) <span>62</span> <span>48</span>	100, 119, 152, 155	0
10	I	119/122 (97%)	-0.08	1 (0%) <span>86</span> <span>75</span>	87, 104, 122, 138	0
11	J	65/70 (92%)	-0.28	0 <span>100</span> <span>100</span>	92, 102, 122, 125	0
12	K	114/120 (95%)	-0.27	0 <span>100</span> <span>100</span>	82, 105, 119, 121	0
13	L	46/70 (65%)	0.32	3 (6%) <span>20</span> <span>14</span>	108, 145, 156, 157	0
All	All	3614/4225 (85%)	-0.04	61 (1%) <span>70</span> <span>57</span>	26, 104, 150, 202	0

The worst 5 of 61 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	A	69	THR	5.0
4	A	72	GLU	4.7
5	B	338	GLY	4.0
7	E	46	TYR	3.9
4	A	44	THR	3.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
16	DUT	B	1308[A]	28/28	0.79	0.35	3.85	131,133,149,150	28
16	DUT	B	1308[B]	28/28	0.79	0.35	1.75	57,59,61,62	28
14	ZN	I	204	1/1	0.98	0.09	-1.08	107,107,107,107	0
14	ZN	A	1735	1/1	0.98	0.09	-1.49	142,142,142,142	0
14	ZN	L	105	1/1	0.97	0.06	-1.58	181,181,181,181	0
14	ZN	C	319	1/1	0.99	0.06	-1.84	104,104,104,104	0
14	ZN	J	101	1/1	0.98	0.12	-1.93	149,149,149,149	0
14	ZN	I	203	1/1	0.93	0.07	-2.02	97,97,97,97	0
14	ZN	B	1307	1/1	0.96	0.09	-2.40	126,126,126,126	0
14	ZN	A	1734	1/1	0.89	0.05	-2.92	159,159,159,159	0
15	MG	A	2000	1/1	0.97	0.22	-	74,74,74,74	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.