



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:28 am GMT

PDB ID : 1NY6
Title : Crystal structure of sigm54 activator (AAA+ ATPase) in the active state
Authors : Lee, S.Y.; de la Torre, A.; Kustu, S.; Nixon, B.T.; Wemmer, D.E.
Deposited on : 2003-02-11
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1001 (3.12-3.08)
Clashscore	112137	1099 (3.12-3.08)
Ramachandran outliers	110173	1057 (3.12-3.08)
Sidechain outliers	110143	1057 (3.12-3.08)
RSRZ outliers	101464	1006 (3.12-3.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	<div> <div></div> <div>28% 54% 10% 7%</div> </div>
1	B	267	<div> <div>3%</div> <div>26% 52% 12% 9%</div> </div>
1	C	267	<div> <div>25% 54% 14% 7%</div> </div>
1	D	267	<div> <div>23% 52% 17% 7%</div> </div>
1	E	267	<div> <div>3%</div> <div>16% 57% 19% 7%</div> </div>
1	F	267	<div> <div>3%</div> <div>21% 54% 16% 7%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	267	<div><div><div></div><div></div><div></div><div></div></div><div>3%23%59%10%7%</div></div>
1	H	267	<div><div><div></div><div></div><div></div><div></div></div><div>5%22%57%12%8%</div></div>
1	I	267	<div><div><div></div><div></div><div></div><div></div></div><div>%28%51%13%7%</div></div>
1	J	267	<div><div><div></div><div></div><div></div><div></div></div><div>5%36%48%8%7%</div></div>
1	K	267	<div><div><div></div><div></div><div></div><div></div></div><div>2%27%52%13%7%</div></div>
1	L	267	<div><div><div></div><div></div><div></div><div></div></div><div>%13%58%19%8%</div></div>
1	M	267	<div><div><div></div><div></div><div></div><div></div></div><div>%21%56%14%7%</div></div>
1	N	267	<div><div><div></div><div></div><div></div><div></div></div><div>%27%52%11%7%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 28041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called transcriptional regulator (NtrC family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	0	0
			1979	1278	333	364	4			
1	B	243	Total	C	N	O	S	0	0	0
			1940	1250	329	357	4			
1	C	248	Total	C	N	O	S	0	0	0
			1988	1283	334	367	4			
1	D	247	Total	C	N	O	S	0	0	0
			1979	1278	333	364	4			
1	E	248	Total	C	N	O	S	0	0	0
			1988	1283	334	367	4			
1	F	248	Total	C	N	O	S	0	0	0
			1988	1283	334	367	4			
1	G	248	Total	C	N	O	S	0	0	0
			1988	1283	334	367	4			
1	H	245	Total	C	N	O	S	0	0	0
			1954	1259	331	360	4			
1	I	247	Total	C	N	O	S	0	0	0
			1966	1268	333	361	4			
1	J	247	Total	C	N	O	S	0	0	0
			1975	1276	333	362	4			
1	K	248	Total	C	N	O	S	0	0	0
			1988	1283	334	367	4			
1	L	246	Total	C	N	O	S	0	0	0
			1972	1273	332	363	4			
1	M	247	Total	C	N	O	S	0	0	0
			1979	1278	333	364	4			
1	N	247	Total	C	N	O	S	0	0	0
			1979	1278	333	364	4			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	121	MET	-	INITIATING METHIONINE	UNP O67198

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	121	MET	-	INITIATING METHIONINE	UNP O67198
C	121	MET	-	INITIATING METHIONINE	UNP O67198
D	121	MET	-	INITIATING METHIONINE	UNP O67198
E	121	MET	-	INITIATING METHIONINE	UNP O67198
F	121	MET	-	INITIATING METHIONINE	UNP O67198
G	121	MET	-	INITIATING METHIONINE	UNP O67198
H	121	MET	-	INITIATING METHIONINE	UNP O67198
I	121	MET	-	INITIATING METHIONINE	UNP O67198
J	121	MET	-	INITIATING METHIONINE	UNP O67198
K	121	MET	-	INITIATING METHIONINE	UNP O67198
L	121	MET	-	INITIATING METHIONINE	UNP O67198
M	121	MET	-	INITIATING METHIONINE	UNP O67198
N	121	MET	-	INITIATING METHIONINE	UNP O67198

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	G	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	I	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	J	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	K	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	L	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	M	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	N	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	H	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: transcriptional regulator (NtrC family)





- Molecule 1: transcriptional regulator (NtrC family)







- Chain C:  25% 54% 14% 7%




- Molecule 1: transcriptional regulator (NtrC family)









- Molecule 1: transcriptional regulator (NtrC family)




- Molecule 1: transcriptional regulator (NtrC family)








- Molecule 1: transcriptional regulator (NtrC family)



- Molecule 1: transcriptional regulator (NtrC family)



- Molecule 1: transcriptional regulator (NtrC family)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	106.79Å 108.27Å 110.02Å 70.25° 85.90° 73.27°	Depositor
Resolution (Å)	19.98 – 3.10 62.44 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.6 (19.98-3.10) 86.2 (62.44-3.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 3.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.266 , 0.329 0.259 , 0.317	Depositor DCC
R_{free} test set	7844 reflections (10.01%)	DCC
Wilson B-factor (Å ²)	94.2	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 82.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	28041	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/2013	0.65	0/2700
1	B	0.48	0/1972	0.67	0/2644
1	C	0.55	0/2022	0.83	1/2712 (0.0%)
1	D	0.55	0/2013	0.78	1/2700 (0.0%)
1	E	0.62	0/2022	0.93	1/2712 (0.0%)
1	F	0.53	0/2022	0.73	0/2712
1	G	0.47	0/2022	0.69	1/2712 (0.0%)
1	H	0.45	0/1987	0.96	2/2665 (0.1%)
1	I	0.45	0/1999	0.68	0/2681
1	J	0.42	0/2009	0.63	0/2695
1	K	0.47	0/2022	0.70	1/2712 (0.0%)
1	L	0.63	0/2006	0.87	1/2690 (0.0%)
1	M	0.52	0/2013	0.74	0/2700
1	N	0.45	0/2013	0.69	1/2700 (0.0%)
All	All	0.51	0/28135	0.76	9/37735 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	2

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	143	SER	C-N-CD	-34.04	45.71	120.60
1	E	190	PRO	CA-N-CD	-20.80	82.37	111.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	190	PRO	CA-N-CD	-16.92	87.81	111.50
1	H	142	GLU	O-C-N	-9.85	106.93	122.70
1	D	262	ARG	NE-CZ-NH2	-5.88	117.36	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	142	GLU	Mainchain,Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1979	0	2040	214	0
1	B	1940	0	2007	249	0
1	C	1988	0	2046	268	0
1	D	1979	0	2040	276	0
1	E	1988	0	2046	359	0
1	F	1988	0	2046	296	0
1	G	1988	0	2046	245	0
1	H	1954	0	2009	281	0
1	I	1966	0	2022	250	0
1	J	1975	0	2036	193	0
1	K	1988	0	2046	246	0
1	L	1972	0	2031	382	0
1	M	1979	0	2040	301	0
1	N	1979	0	2040	271	0
2	A	27	0	12	2	0
2	B	27	0	12	1	0
2	C	27	0	12	4	0
2	D	27	0	12	6	0
2	E	27	0	12	3	0
2	F	27	0	12	7	0
2	G	27	0	12	0	0
2	H	27	0	12	3	0
2	I	27	0	12	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	J	27	0	12	1	0
2	K	27	0	12	1	0
2	L	27	0	12	6	0
2	M	27	0	12	6	0
2	N	27	0	12	2	0
All	All	28041	0	28663	3697	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 65.

The worst 5 of 3697 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:235:LEU:HD12	1:D:236:PHE:N	1.44	1.30
1:L:334:LYS:NZ	1:L:367:VAL:HG13	1.51	1.24
1:E:177:ALA:CA	1:E:180:ILE:HD12	1.69	1.21
1:E:240:ILE:CD1	1:E:277:ALA:HB1	1.72	1.18
1:F:318:ILE:HG13	1:F:319:PRO:HD3	1.23	1.17

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/267 (92%)	192 (78%)	39 (16%)	14 (6%)	2	12
1	B	241/267 (90%)	155 (64%)	64 (27%)	22 (9%)	1	4
1	C	246/267 (92%)	180 (73%)	49 (20%)	17 (7%)	1	8
1	D	245/267 (92%)	165 (67%)	55 (22%)	25 (10%)	1	4
1	E	246/267 (92%)	165 (67%)	56 (23%)	25 (10%)	1	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	246/267 (92%)	179 (73%)	39 (16%)	28 (11%)	0	2
1	G	246/267 (92%)	177 (72%)	49 (20%)	20 (8%)	1	6
1	H	243/267 (91%)	171 (70%)	55 (23%)	17 (7%)	1	8
1	I	245/267 (92%)	174 (71%)	48 (20%)	23 (9%)	1	4
1	J	245/267 (92%)	182 (74%)	48 (20%)	15 (6%)	2	11
1	K	246/267 (92%)	173 (70%)	53 (22%)	20 (8%)	1	6
1	L	244/267 (91%)	158 (65%)	53 (22%)	33 (14%)	0	1
1	M	245/267 (92%)	164 (67%)	57 (23%)	24 (10%)	1	4
1	N	245/267 (92%)	169 (69%)	57 (23%)	19 (8%)	1	6
All	All	3428/3738 (92%)	2404 (70%)	722 (21%)	302 (9%)	1	5

5 of 302 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	144	PRO
1	A	221	SER
1	B	157	SER
1	B	217	THR
1	B	381	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/232 (91%)	187 (88%)	25 (12%)	6	25
1	B	208/232 (90%)	185 (89%)	23 (11%)	7	29
1	C	213/232 (92%)	188 (88%)	25 (12%)	6	26
1	D	212/232 (91%)	180 (85%)	32 (15%)	3	15
1	E	213/232 (92%)	176 (83%)	37 (17%)	2	10
1	F	213/232 (92%)	175 (82%)	38 (18%)	2	9
1	G	213/232 (92%)	188 (88%)	25 (12%)	6	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	208/232 (90%)	187 (90%)	21 (10%)	9	33
1	I	209/232 (90%)	187 (90%)	22 (10%)	8	31
1	J	211/232 (91%)	193 (92%)	18 (8%)	12	44
1	K	213/232 (92%)	188 (88%)	25 (12%)	6	26
1	L	211/232 (91%)	171 (81%)	40 (19%)	2	7
1	M	212/232 (91%)	186 (88%)	26 (12%)	5	23
1	N	212/232 (91%)	185 (87%)	27 (13%)	5	21
All	All	2960/3248 (91%)	2576 (87%)	384 (13%)	5	21

5 of 384 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	357	ARG
1	H	252	LEU
1	M	361	ASN
1	F	383	LEU
1	G	274	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	195	ASN
1	H	344	GLN
1	M	361	ASN
1	G	181	HIS
1	I	195	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	6	-	25,29,29	1.56	5 (20%)	24,45,45	2.74	3 (12%)
2	ADP	B	7	-	25,29,29	1.49	5 (20%)	24,45,45	2.72	4 (16%)
2	ADP	C	1	-	25,29,29	1.44	3 (12%)	24,45,45	2.63	2 (8%)
2	ADP	D	2	-	25,29,29	1.54	5 (20%)	24,45,45	2.74	4 (16%)
2	ADP	E	3	-	25,29,29	1.54	6 (24%)	24,45,45	2.79	1 (4%)
2	ADP	F	4	-	25,29,29	1.47	4 (16%)	24,45,45	2.66	4 (16%)
2	ADP	G	5	-	25,29,29	1.46	4 (16%)	24,45,45	2.74	3 (12%)
2	ADP	H	14	-	25,29,29	1.48	5 (20%)	24,45,45	2.71	3 (12%)
2	ADP	I	8	-	25,29,29	1.44	4 (16%)	24,45,45	2.69	3 (12%)
2	ADP	J	9	-	25,29,29	1.48	7 (28%)	24,45,45	2.71	4 (16%)
2	ADP	K	10	-	25,29,29	1.48	5 (20%)	24,45,45	2.78	3 (12%)
2	ADP	L	11	-	25,29,29	1.52	5 (20%)	24,45,45	2.73	3 (12%)
2	ADP	M	12	-	25,29,29	1.42	3 (12%)	24,45,45	2.63	4 (16%)
2	ADP	N	13	-	25,29,29	1.45	5 (20%)	24,45,45	2.68	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	6	-	-	0/12/32/32	0/3/3/3
2	ADP	B	7	-	-	0/12/32/32	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	C	1	-	-	0/12/32/32	0/3/3/3
2	ADP	D	2	-	-	0/12/32/32	0/3/3/3
2	ADP	E	3	-	-	0/12/32/32	0/3/3/3
2	ADP	F	4	-	-	0/12/32/32	0/3/3/3
2	ADP	G	5	-	-	0/12/32/32	0/3/3/3
2	ADP	H	14	-	-	0/12/32/32	0/3/3/3
2	ADP	I	8	-	-	0/12/32/32	0/3/3/3
2	ADP	J	9	-	-	0/12/32/32	0/3/3/3
2	ADP	K	10	-	-	0/12/32/32	0/3/3/3
2	ADP	L	11	-	-	0/12/32/32	0/3/3/3
2	ADP	M	12	-	-	0/12/32/32	0/3/3/3
2	ADP	N	13	-	-	0/12/32/32	0/3/3/3

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	ADP	PB-O3A	-3.16	1.55	1.60
2	A	6	ADP	C5-N7	-2.92	1.29	1.39
2	E	3	ADP	PB-O3A	-2.87	1.55	1.60
2	C	1	ADP	C5-N7	-2.75	1.30	1.39
2	M	12	ADP	C5-N7	-2.71	1.30	1.39

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	3	ADP	N3-C2-N1	-12.81	117.70	128.86
2	K	10	ADP	N3-C2-N1	-12.45	118.01	128.86
2	G	5	ADP	N3-C2-N1	-12.41	118.05	128.86
2	D	2	ADP	N3-C2-N1	-12.33	118.12	128.86
2	J	9	ADP	N3-C2-N1	-12.27	118.17	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

13 monomers are involved in 47 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	6	ADP	2	0
2	B	7	ADP	1	0
2	C	1	ADP	4	0
2	D	2	ADP	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	3	ADP	3	0
2	F	4	ADP	7	0
2	H	14	ADP	3	0
2	I	8	ADP	5	0
2	J	9	ADP	1	0
2	K	10	ADP	1	0
2	L	11	ADP	6	0
2	M	12	ADP	6	0
2	N	13	ADP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/267 (92%)	-0.05	4 (1%) 72 51	29, 107, 142, 151	0
1	B	243/267 (91%)	0.06	7 (2%) 52 28	18, 101, 150, 151	0
1	C	248/267 (92%)	-0.13	0 100 100	43, 85, 128, 142	0
1	D	247/267 (92%)	-0.16	0 100 100	43, 90, 136, 148	0
1	E	248/267 (92%)	-0.08	8 (3%) 48 25	18, 83, 140, 151	0
1	F	248/267 (92%)	-0.03	7 (2%) 53 29	44, 106, 146, 150	0
1	G	248/267 (92%)	0.07	7 (2%) 53 29	64, 114, 150, 151	0
1	H	245/267 (91%)	0.25	14 (5%) 24 11	77, 129, 151, 151	0
1	I	247/267 (92%)	-0.09	4 (1%) 72 51	65, 103, 147, 151	0
1	J	247/267 (92%)	0.19	14 (5%) 24 11	71, 119, 147, 151	0
1	K	248/267 (92%)	0.03	6 (2%) 59 37	57, 104, 137, 148	0
1	L	246/267 (92%)	-0.11	2 (0%) 86 71	27, 80, 126, 150	0
1	M	247/267 (92%)	-0.07	2 (0%) 86 71	25, 100, 139, 146	0
1	N	247/267 (92%)	-0.10	3 (1%) 79 61	67, 106, 138, 151	0
All	All	3456/3738 (92%)	-0.02	78 (2%) 61 39	18, 102, 148, 151	0

The worst 5 of 78 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	382	CYS	7.0
1	E	264	GLY	6.2
1	F	276	LEU	5.4
1	J	305	ILE	5.2
1	E	211	TYR	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ADP	A	6	27/27	0.93	0.20	-0.09	84,88,92,96	0
2	ADP	D	2	27/27	0.91	0.22	-0.21	71,82,91,93	0
2	ADP	H	14	27/27	0.92	0.25	-0.30	125,141,143,143	0
2	ADP	G	5	27/27	0.93	0.19	-0.33	52,65,75,80	0
2	ADP	I	8	27/27	0.96	0.17	-0.36	66,81,87,90	0
2	ADP	B	7	27/27	0.91	0.21	-0.50	96,111,114,114	0
2	ADP	L	11	27/27	0.94	0.20	-0.54	48,60,84,93	0
2	ADP	N	13	27/27	0.91	0.18	-0.55	100,106,110,111	0
2	ADP	M	12	27/27	0.93	0.18	-0.56	51,67,86,87	0
2	ADP	K	10	27/27	0.90	0.21	-0.61	99,103,108,110	0
2	ADP	C	1	27/27	0.96	0.19	-0.63	54,70,77,79	0
2	ADP	E	3	27/27	0.96	0.18	-0.71	51,64,70,79	0
2	ADP	J	9	27/27	0.92	0.16	-0.94	92,95,102,104	0
2	ADP	F	4	27/27	0.94	0.17	-0.97	62,66,84,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.