



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:38 pm GMT

PDB ID : 4O0P
Title : Crystal Structure of D. radiodurans Bacteriophytochrome Photosensory Core Module in its Dark Form
Authors : Takala, H.; Ihalainen, J.A.; Westenhoff, S.
Deposited on : 2013-12-14
Resolution : 3.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1019 (4.08-3.52)
Clashscore	112137	1030 (4.04-3.56)
Ramachandran outliers	110173	1011 (4.06-3.54)
Sidechain outliers	110143	1005 (4.06-3.54)
RSRZ outliers	101464	1032 (4.08-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	523	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>5%</div> <div>7%</div> </div> </div>
1	B	523	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>7%</div> <div>7%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriophytochrome.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	0	0
			3717	2368	658	680	11			
1	B	484	Total	C	N	O	S	0	0	0
			3699	2356	651	681	11			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-13	MET	-	EXPRESSION TAG	UNP Q9RZA4
A	-12	ALA	-	EXPRESSION TAG	UNP Q9RZA4
A	-11	SER	-	EXPRESSION TAG	UNP Q9RZA4
A	-10	MET	-	EXPRESSION TAG	UNP Q9RZA4
A	-9	THR	-	EXPRESSION TAG	UNP Q9RZA4
A	-8	GLY	-	EXPRESSION TAG	UNP Q9RZA4
A	-7	GLY	-	EXPRESSION TAG	UNP Q9RZA4
A	-6	GLN	-	EXPRESSION TAG	UNP Q9RZA4
A	-5	GLN	-	EXPRESSION TAG	UNP Q9RZA4
A	-4	MET	-	EXPRESSION TAG	UNP Q9RZA4
A	-3	GLY	-	EXPRESSION TAG	UNP Q9RZA4
A	-2	ARG	-	EXPRESSION TAG	UNP Q9RZA4
A	-1	GLY	-	EXPRESSION TAG	UNP Q9RZA4
A	0	SER	-	EXPRESSION TAG	UNP Q9RZA4
A	503	GLU	-	EXPRESSION TAG	UNP Q9RZA4
A	504	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	505	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	506	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	507	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	508	HIS	-	EXPRESSION TAG	UNP Q9RZA4
A	509	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	-13	MET	-	EXPRESSION TAG	UNP Q9RZA4
B	-12	ALA	-	EXPRESSION TAG	UNP Q9RZA4
B	-11	SER	-	EXPRESSION TAG	UNP Q9RZA4
B	-10	MET	-	EXPRESSION TAG	UNP Q9RZA4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	THR	-	EXPRESSION TAG	UNP Q9RZA4
B	-8	GLY	-	EXPRESSION TAG	UNP Q9RZA4
B	-7	GLY	-	EXPRESSION TAG	UNP Q9RZA4
B	-6	GLN	-	EXPRESSION TAG	UNP Q9RZA4
B	-5	GLN	-	EXPRESSION TAG	UNP Q9RZA4
B	-4	MET	-	EXPRESSION TAG	UNP Q9RZA4
B	-3	GLY	-	EXPRESSION TAG	UNP Q9RZA4
B	-2	ARG	-	EXPRESSION TAG	UNP Q9RZA4
B	-1	GLY	-	EXPRESSION TAG	UNP Q9RZA4
B	0	SER	-	EXPRESSION TAG	UNP Q9RZA4
B	503	GLU	-	EXPRESSION TAG	UNP Q9RZA4
B	504	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	505	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	506	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	507	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	508	HIS	-	EXPRESSION TAG	UNP Q9RZA4
B	509	HIS	-	EXPRESSION TAG	UNP Q9RZA4

- Molecule 2 is 3-[2-[(Z)-[3-(2-CARBOXYETHYL)-5-[(Z)-(4-ETHENYL-3-METHYL-5-OXIDANYLIDENE-PYRROL-2-YLIDENE)METHYL]-4-METHYL-PYRROL-1-IUM-2-YLIDENE]METHYL]-5-[(Z)-[(3E)-3-ETHYLIDENE-4-METHYL-5-OXIDANYLIDENE-PYRROLIDIN-2-YLIDENE]METHYL]-4-METHYL-1H-PYRROL-3-YL]PROPANOIC ACID (three-letter code: LBV) (formula: C₃₃H₃₇N₄O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			43	33	4	6		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			43	33	4	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bacteriophytochrome



• Molecule 1: Bacteriophytochrome



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	150.28Å 150.28Å 145.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.65 – 3.80 49.19 – 3.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.65-3.80) 99.7 (49.19-3.80)	Depositor EDS
R_{merge}	0.39	Depositor
R_{sym}	0.41	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.12 (at 3.77Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.253 , 0.266 0.231 , 0.249	Depositor DCC
R_{free} test set	928 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	167.6	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 164.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.116 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7502	wwPDB-VP
Average B, all atoms (Å ²)	210.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LBV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/3815	0.51	1/5220 (0.0%)
1	B	0.30	0/3795	0.52	2/5193 (0.0%)
All	All	0.31	0/7610	0.51	3/10413 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	457	ASP	CB-CG-OD2	9.48	126.83	118.30
1	A	457	ASP	CB-CG-OD2	-8.28	110.85	118.30
1	B	457	ASP	CB-CG-OD1	-6.83	112.15	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3717	0	3674	17	0
1	B	3699	0	3661	21	0
2	A	43	0	33	3	0
2	B	43	0	33	4	0
All	All	7502	0	7401	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:360:LEU:HD21	1:B:379:LEU:HD12	1.79	0.64
1:A:360:LEU:HD21	1:A:379:LEU:HD12	1.79	0.64
2:B:600:LBV:HMC1	2:B:600:LBV:N_D	2.20	0.57
1:B:449:VAL:HG11	1:B:451:TRP:CE2	2.40	0.56
1:A:449:VAL:HG11	1:A:451:TRP:CE2	2.39	0.56
2:B:600:LBV:HBA2	2:B:600:LBV:HMA1	1.87	0.56
1:A:384:PRO:HD2	1:A:418:LEU:HD21	1.91	0.52
1:B:384:PRO:HD2	1:B:418:LEU:HD21	1.92	0.52
2:A:600:LBV:C1D	2:A:600:LBV:HMC1	2.41	0.51
1:A:455:THR:N	1:A:456:PRO:CD	2.75	0.50
2:B:600:LBV:HMC1	2:B:600:LBV:C1D	2.43	0.49
1:B:455:THR:N	1:B:456:PRO:CD	2.76	0.49
1:B:209:PRO:HD3	2:B:600:LBV:C1B	2.44	0.47
1:B:379:LEU:HD13	1:B:380:GLY:N	2.30	0.46
1:A:379:LEU:HD13	1:A:380:GLY:N	2.30	0.46
1:A:113:LEU:HD13	1:A:126:PHE:CE1	2.51	0.46
1:A:202:ARG:O	1:A:451:TRP:HA	2.15	0.45
1:B:183:THR:OG1	1:B:202:ARG:NH1	2.49	0.45
1:B:434:TRP:HA	1:B:437:CYS:SG	2.57	0.45
1:B:202:ARG:HD3	1:B:449:VAL:HG13	1.99	0.45
1:B:315:GLN:O	1:B:319:LYS:HG2	2.16	0.45
1:A:434:TRP:HA	1:A:437:CYS:SG	2.57	0.44
1:B:449:VAL:HG12	1:B:450:ALA:N	2.32	0.44
1:A:419:ALA:N	1:A:420:PRO:CD	2.81	0.43
1:A:419:ALA:N	1:A:420:PRO:HD2	2.33	0.43
1:B:419:ALA:N	1:B:420:PRO:CD	2.81	0.43
1:B:419:ALA:N	1:B:420:PRO:HD2	2.33	0.43
2:A:600:LBV:HMA1	2:A:600:LBV:HBA2	1.99	0.43
1:B:113:LEU:HD13	1:B:126:PHE:CE1	2.53	0.43
1:A:449:VAL:HG12	1:A:450:ALA:N	2.34	0.43
1:A:368:LEU:HD12	1:A:368:LEU:C	2.38	0.43
1:A:202:ARG:HD3	1:A:449:VAL:HG13	2.01	0.43
1:B:154:ARG:NH1	1:B:187:ILE:O	2.52	0.42
1:B:368:LEU:C	1:B:368:LEU:HD12	2.39	0.42
1:B:360:LEU:HD22	1:B:360:LEU:N	2.35	0.41
1:A:360:LEU:HD22	1:A:360:LEU:N	2.34	0.41
1:A:428:ILE:HD12	1:A:491:ALA:HB1	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:PHE:O	1:B:199:LEU:C	2.59	0.41
1:B:202:ARG:O	1:B:451:TRP:HA	2.21	0.41
1:A:260:HIS:CE1	2:A:600:LBV:C1C	3.03	0.41
1:B:428:ILE:HD12	1:B:491:ALA:HB1	2.02	0.40
1:A:154:ARG:NH1	1:A:187:ILE:O	2.51	0.40
1:B:201:HIS:CE1	1:B:471:THR:HG23	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/523 (90%)	439 (93%)	32 (7%)	2 (0%)	38	77
1	B	474/523 (91%)	441 (93%)	31 (6%)	2 (0%)	38	77
All	All	947/1046 (90%)	880 (93%)	63 (7%)	4 (0%)	38	77

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	119	GLY
1	A	119	GLY
1	A	455	THR
1	B	455	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/410 (94%)	381 (99%)	4 (1%)	80	90
1	B	383/410 (93%)	378 (99%)	5 (1%)	73	88
All	All	768/820 (94%)	759 (99%)	9 (1%)	75	88

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	SER
1	A	222	ARG
1	A	289	CYS
1	A	354	ASP
1	B	206	SER
1	B	222	ARG
1	B	246	THR
1	B	289	CYS
1	B	354	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	201	HIS
1	A	290	HIS
1	B	64	GLN
1	B	201	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LBV	A	600	1	35,46,46	4.12	14 (40%)	38,67,67	3.11	15 (39%)
2	LBV	B	600	1	35,46,46	4.23	14 (40%)	38,67,67	2.92	18 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LBV	A	600	1	-	2/22/74/74	0/4/4/4
2	LBV	B	600	1	-	3/22/74/74	0/4/4/4

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	LBV	C1A-N_A	-3.00	1.33	1.37
2	B	600	LBV	C1A-N_A	-2.97	1.33	1.37
2	B	600	LBV	C4D-N_D	-2.20	1.33	1.37
2	B	600	LBV	O_D-C4D	2.24	1.27	1.23
2	B	600	LBV	C1C-C2C	2.28	1.49	1.45
2	A	600	LBV	CAA-C3A	2.33	1.36	1.33
2	A	600	LBV	O_D-C4D	2.33	1.28	1.23
2	A	600	LBV	C1C-C2C	2.53	1.49	1.45
2	B	600	LBV	C4A-C3A	2.82	1.50	1.45
2	B	600	LBV	C1D-C2D	2.90	1.50	1.45
2	A	600	LBV	C1B-CHB	2.98	1.51	1.40
2	A	600	LBV	C1D-C2D	3.03	1.50	1.45
2	B	600	LBV	C1B-CHB	3.11	1.52	1.40
2	A	600	LBV	C4B-CHC	3.16	1.52	1.40
2	A	600	LBV	C4A-C3A	3.32	1.51	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	600	LBV	C4B-CHC	3.47	1.53	1.40
2	B	600	LBV	C4A-N_A	4.04	1.44	1.37
2	A	600	LBV	C4A-N_A	4.23	1.45	1.37
2	A	600	LBV	CHD-C4C	4.38	1.50	1.40
2	B	600	LBV	CHD-C4C	4.80	1.51	1.40
2	A	600	LBV	CHD-C1D	5.33	1.50	1.37
2	B	600	LBV	C3D-C2D	5.46	1.47	1.37
2	A	600	LBV	C3D-C2D	5.50	1.47	1.37
2	B	600	LBV	CHD-C1D	5.80	1.51	1.37
2	B	600	LBV	CHB-C4A	8.16	1.52	1.34
2	A	600	LBV	CHB-C4A	8.70	1.53	1.34
2	A	600	LBV	CHC-C1C	18.48	1.51	1.35
2	B	600	LBV	CHC-C1C	19.45	1.52	1.35

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	LBV	CHB-C4A-C3A	-11.35	106.38	127.17
2	B	600	LBV	CHB-C4A-C3A	-9.63	109.51	127.17
2	A	600	LBV	C4B-CHC-C1C	-8.28	118.74	128.77
2	B	600	LBV	C4B-CHC-C1C	-7.73	119.40	128.77
2	A	600	LBV	C1D-CHD-C4C	-6.72	111.15	128.08
2	B	600	LBV	CHB-C4A-N_A	-5.57	111.26	130.29
2	B	600	LBV	C1D-CHD-C4C	-4.80	115.98	128.08
2	A	600	LBV	C4D-C3D-C2D	-3.86	102.86	108.08
2	A	600	LBV	CHB-C4A-N_A	-3.74	117.52	130.29
2	B	600	LBV	C4D-C3D-C2D	-3.73	103.04	108.08
2	B	600	LBV	C3D-C2D-C1D	-3.01	104.17	107.91
2	A	600	LBV	C3D-C2D-C1D	-3.00	104.19	107.91
2	B	600	LBV	CAB-CBB-CGB	-2.92	107.68	112.66
2	B	600	LBV	CAA-C3A-C4A	-2.90	123.18	126.46
2	A	600	LBV	CAA-C3A-C4A	-2.90	123.18	126.46
2	A	600	LBV	CBB-CAB-C3B	-2.64	107.44	112.48
2	A	600	LBV	O_D-C4D-C3D	-2.62	123.71	129.77
2	A	600	LBV	C1D-N_D-C4D	-2.51	107.36	110.70
2	B	600	LBV	CHD-C4C-N_C	-2.37	120.41	124.97
2	B	600	LBV	C1D-N_D-C4D	-2.18	107.80	110.70
2	B	600	LBV	CHD-C1D-C2D	-2.09	122.64	126.83
2	B	600	LBV	O_D-C4D-C3D	-2.06	125.00	129.77
2	A	600	LBV	C1C-N_C-C4C	2.21	110.89	106.52
2	A	600	LBV	CMB-C2B-C3B	2.32	129.31	124.94
2	B	600	LBV	CMC-C3C-C4C	2.36	128.72	125.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	LBV	CAD-C3D-C4D	2.52	131.01	123.40
2	B	600	LBV	CHD-C4C-C3C	2.58	131.01	124.81
2	B	600	LBV	C1C-N_C-C4C	2.60	111.66	106.52
2	B	600	LBV	CAD-C3D-C4D	2.61	131.27	123.40
2	B	600	LBV	CMD-C2D-C1D	2.69	127.68	124.17
2	B	600	LBV	C2D-C1D-N_D	2.70	111.06	107.04
2	A	600	LBV	C2D-C1D-N_D	2.90	111.36	107.04
2	A	600	LBV	CMD-C2D-C1D	3.02	128.12	124.17

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	600	LBV	C1B-CHB-C4A-N_A
2	A	600	LBV	C1D-CHD-C4C-C3C
2	A	600	LBV	C1D-CHD-C4C-N_C
2	B	600	LBV	C1D-CHD-C4C-C3C
2	B	600	LBV	C1D-CHD-C4C-N_C

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	LBV	3	0
2	B	600	LBV	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	485/523 (92%)	0.03	11 (2%) 61 51	94, 208, 275, 324	0
1	B	484/523 (92%)	0.03	14 (2%) 52 42	118, 206, 281, 329	0
All	All	969/1046 (92%)	0.03	25 (2%) 56 46	94, 207, 278, 329	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	ASP	6.0
1	B	362	LEU	4.5
1	B	404	GLN	4.1
1	B	403	VAL	3.6
1	B	503	GLU	3.3
1	B	438	LEU	3.1
1	A	44	ALA	3.1
1	B	40	ALA	3.0
1	A	62	LEU	2.9
1	B	435	SER	2.8
1	A	438	LEU	2.8
1	A	331	LEU	2.6
1	B	126	PHE	2.5
1	B	4	ASP	2.4
1	A	106	PRO	2.4
1	A	14	LEU	2.3
1	B	370	LEU	2.3
1	B	391	LEU	2.3
1	A	61	PHE	2.2
1	B	490	GLU	2.2
1	A	13	TYR	2.2
1	B	487	GLU	2.1
1	A	203	PHE	2.1
1	A	17	PRO	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	415	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	LBV	B	600	43/43	0.93	0.38	1.55	131,174,209,237	0
2	LBV	A	600	43/43	0.95	0.39	0.85	116,159,192,236	0

6.5 Other polymers [i](#)

There are no such residues in this entry.