



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 08:52 pm GMT

PDB ID : 2O35
Title : Protein of Unknown Function (DUF1244) from Sinorhizobium meliloti
Authors : Kim, Y.; Joachimiak, A.; Evdokimova, E.; Kudritska, M.; Edwards, A.; Savchenko, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-11-30
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4988 (2.14-2.10)
Clashscore	112137	5557 (2.14-2.10)
Ramachandran outliers	110173	5504 (2.14-2.10)
Sidechain outliers	110143	5505 (2.14-2.10)
RSRZ outliers	101464	5021 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	105	<div> <div>22%</div> <div> <div></div> <div>54%</div> <div>18%</div> <div>•</div> <div>25%</div> </div> </div>
1	B	105	<div> <div>25%</div> <div> <div></div> <div>58%</div> <div>25%</div> <div>7%</div> <div>•</div> <div>10%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein DUF1244.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	79	Total	C	N	O	S	Se	0	5	0
			693	428	123	136	2	4			
1	B	95	Total	C	N	O	S	Se	0	9	0
			855	521	159	170	2	3			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP Q92M60
A	0	HIS	-	CLONING ARTIFACT	UNP Q92M60
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
A	16	VAL	ALA	SEE REMARK 999	UNP Q92M60
A	36	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
A	61	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
A	73	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
A	102	GLY	-	CLONING ARTIFACT	UNP Q92M60
A	103	SER	-	CLONING ARTIFACT	UNP Q92M60
B	-1	GLY	-	CLONING ARTIFACT	UNP Q92M60
B	0	HIS	-	CLONING ARTIFACT	UNP Q92M60
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
B	16	VAL	ALA	SEE REMARK 999	UNP Q92M60
B	36	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
B	61	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
B	73	MSE	MET	MODIFIED RESIDUE	UNP Q92M60
B	102	GLY	-	CLONING ARTIFACT	UNP Q92M60
B	103	SER	-	CLONING ARTIFACT	UNP Q92M60

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	46	Total	O	0	0
			46	46		
3	B	71	Total	O	0	0
			71	71		

4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.99Å 80.99Å 64.71Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.07 – 2.12 35.07 – 2.12	Depositor EDS
% Data completeness (in resolution range)	98.1 (35.07-2.12) 98.1 (35.07-2.12)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.209 , 0.263 0.284 , 0.322	Depositor DCC
R_{free} test set	1382 reflections (9.93%)	DCC
Wilson B-factor (Å ²)	39.9	Xtriage
Anisotropy	0.539	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.042 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	1667	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/703 (0.1%)	0.83	1/940 (0.1%)
1	B	1.00	5/866 (0.6%)	0.88	1/1158 (0.1%)
All	All	0.94	6/1569 (0.4%)	0.86	2/2098 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	64[A]	GLU	CG-CD	5.98	1.60	1.51
1	B	64[B]	GLU	CG-CD	5.98	1.60	1.51
1	B	64[A]	GLU	CD-OE2	5.55	1.31	1.25
1	B	64[B]	GLU	CD-OE2	5.55	1.31	1.25
1	B	16	VAL	CB-CG2	-5.53	1.41	1.52
1	A	80	THR	CB-OG1	5.01	1.53	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	51	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	B	19	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	693	0	649	17	0
1	B	855	0	791	39	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	46	0	0	1	0
3	B	71	0	0	4	0
All	All	1667	0	1440	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:MSE:HE3	1:B:66:SER:HB3	1.13	1.08
1:B:51:ARG:HA	1:B:61:MSE:HE2	1.36	1.07
1:B:43[A]:ARG:HH21	1:B:43[A]:ARG:HG3	1.27	0.97
1:B:87:SER:H	1:B:90[B]:GLN:HE21	0.94	0.91
1:B:61:MSE:HE3	1:B:66:SER:CB	2.00	0.91
1:B:12:PHE:O	1:B:16:VAL:HG23	1.70	0.89
1:B:87:SER:H	1:B:90[B]:GLN:NE2	1.70	0.89
1:B:51:ARG:HA	1:B:61:MSE:CE	2.05	0.86
1:A:12:PHE:O	1:A:16:VAL:HG23	1.78	0.82
1:B:87:SER:N	1:B:90[B]:GLN:HE21	1.78	0.78
1:A:51:ARG:HD2	1:A:61:MSE:O	1.85	0.77
1:A:79:ARG:HH11	1:A:79:ARG:CG	1.98	0.76
1:B:43[A]:ARG:HH21	1:B:43[A]:ARG:CG	2.04	0.70
1:A:79:ARG:HG2	1:A:79:ARG:HH11	1.57	0.69
1:B:4[B]:ILE:HG22	1:B:9[B]:ARG:HD2	1.74	0.68
1:A:52:GLU:OE2	1:B:19:ARG:NH1	2.28	0.67
1:A:76[B]:GLU:CD	1:A:76[B]:GLU:H	1.96	0.67
1:B:44[B]:ASN:O	1:B:48[B]:ASN:ND2	2.29	0.65
1:B:87:SER:OG	1:B:90[B]:GLN:HG3	1.97	0.64
1:B:43[B]:ARG:NH2	3:B:218:HOH:O	2.31	0.62
1:B:51:ARG:O	1:B:55:GLU:HG3	2.00	0.62
1:B:51:ARG:CA	1:B:61:MSE:HE2	2.22	0.61
1:B:4[B]:ILE:CG2	1:B:9[B]:ARG:HD2	2.31	0.61
3:A:204:HOH:O	1:B:19:ARG:HG3	2.01	0.60
1:B:51:ARG:CA	1:B:61:MSE:CE	2.79	0.60
1:A:79:ARG:HG2	1:A:79:ARG:NH1	2.15	0.60
1:B:22[A]:GLU:HG2	1:B:23:HIS:N	2.10	0.59
1:B:43[A]:ARG:HG3	1:B:43[A]:ARG:NH2	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:TYR:HD2	1:B:61:MSE:HE1	1.71	0.54
1:B:28:SER:HB3	3:B:240:HOH:O	2.07	0.53
1:A:38:LEU:HD11	1:B:34:ASP:HB3	1.89	0.53
1:A:17:PHE:HB2	1:B:17:PHE:HB2	1.90	0.52
1:B:32:ASN:OD1	1:B:43[B]:ARG:HG3	2.10	0.52
1:B:97:ASN:HB2	3:B:233:HOH:O	2.11	0.51
1:B:32:ASN:OD1	1:B:43[A]:ARG:HG3	2.10	0.51
1:A:22[B]:GLU:O	1:A:26[B]:GLU:HG3	2.11	0.51
1:A:36[B]:MSE:HE2	1:A:37:ASN:OD1	2.14	0.48
1:A:2:SER:N	1:A:9:ARG:HH21	2.11	0.47
1:A:50:TYR:HA	1:B:16:VAL:CG2	2.44	0.47
1:A:51:ARG:CZ	1:A:63[A]:LYS:HG3	2.45	0.47
1:A:32:ASN:HD21	1:A:43:ARG:HB2	1.81	0.46
1:B:4[B]:ILE:HG23	1:B:8:GLN:HB2	1.98	0.46
1:B:26[A]:GLU:HG3	3:B:264:HOH:O	2.16	0.45
1:B:79:ARG:O	1:B:82:ASN:HB2	2.18	0.44
1:B:96:ARG:HG3	1:B:97:ASN:N	2.33	0.43
1:B:48[B]:ASN:O	1:B:52:GLU:HG3	2.19	0.43
1:B:2:SER:OG	1:B:3:GLU:N	2.52	0.42
1:A:16:VAL:CG2	1:B:50:TYR:HA	2.49	0.42
1:B:80:THR:C	1:B:82:ASN:H	2.23	0.41
1:A:32:ASN:ND2	1:A:43:ARG:HH11	2.19	0.40
1:B:22[B]:GLU:OE2	1:B:25:ARG:NH1	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	82/105 (78%)	79 (96%)	3 (4%)	0	100	100
1	B	100/105 (95%)	94 (94%)	6 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	182/210 (87%)	173 (95%)	9 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	74/85 (87%)	69 (93%)	5 (7%)	18	15
1	B	90/85 (106%)	80 (89%)	10 (11%)	7	3
All	All	164/170 (96%)	149 (91%)	15 (9%)	14	7

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	21	LEU
1	A	22[A]	GLU
1	A	22[B]	GLU
1	A	79	ARG
1	B	2	SER
1	B	19	ARG
1	B	22[A]	GLU
1	B	22[B]	GLU
1	B	28	SER
1	B	35	LEU
1	B	43[A]	ARG
1	B	43[B]	ARG
1	B	79	ARG
1	B	82	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	76/105 (72%)	1.43	23 (30%) 0 1	45, 55, 63, 76	0
1	B	92/105 (87%)	1.35	26 (28%) 1 1	20, 50, 66, 81	0
All	All	168/210 (80%)	1.39	49 (29%) 1 1	20, 53, 66, 81	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	37	ASN	5.4
1	B	80	THR	4.8
1	B	38	LEU	4.8
1	B	84	GLY	4.6
1	B	64[A]	GLU	4.5
1	A	49	TRP	4.0
1	B	33	ILE	3.9
1	A	11	ALA	3.8
1	A	31	GLN	3.8
1	B	96	ARG	3.8
1	B	76	GLU	3.8
1	A	54	ALA	3.7
1	A	15	ALA	3.5
1	A	56	ALA	3.4
1	A	53	ALA	3.3
1	A	50	TYR	3.3
1	B	63	LYS	3.3
1	B	75	TYR	3.2
1	B	39	ALA	3.2
1	A	14	ALA	3.0
1	A	29	ASP	2.9
1	B	9[A]	ARG	2.8
1	B	15	ALA	2.7
1	B	74	PRO	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	79	ARG	2.7
1	B	10	THR	2.7
1	B	41	PHE	2.7
1	A	16	VAL	2.6
1	B	11	ALA	2.6
1	A	10	THR	2.6
1	A	57	SER	2.6
1	A	27	ARG	2.6
1	A	46	LEU	2.6
1	B	72	GLY	2.5
1	A	34	ASP	2.5
1	A	79	ARG	2.5
1	B	2	SER	2.5
1	B	14	ALA	2.4
1	B	16	VAL	2.3
1	B	34	ASP	2.3
1	A	76[A]	GLU	2.2
1	B	5	SER	2.2
1	A	77	GLU	2.2
1	B	97	ASN	2.1
1	B	3	GLU	2.1
1	A	30	VAL	2.1
1	A	2	SER	2.0
1	A	45	CYS	2.1
1	A	63[A]	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	B	201	1/1	0.76	0.12	-1.79	76,76,76,76	0
2	MG	A	201	1/1	0.78	0.10	-	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.