



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 09:29 pm GMT

PDB ID : 3O43  
Title : Complex of an alpha/beta-peptide based on the gp41 CHR domain bound to gp41-5  
Authors : Horne, W.S.; Johnson, L.M.; Gellman, S.H.  
Deposited on : 2010-07-26  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	<div> <div>8%</div> <div>76%</div> <div>22%</div> <div>•</div> </div>
2	B	40	<div> <div>3%</div> <div>55%</div> <div>28%</div> <div>•</div> <div>15%</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3835 atoms, of which 1924 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called gp41-5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	198	Total	C	H	N	O	S	507	5	0
			3142	976	1563	293	308	2			

- Molecule 2 is a protein called alpha/beta-peptide derived from gp41 CHR domain sequence.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	34	Total	C	H	N	O	66	0	0
			554	183	273	46	52			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	2	0
			14	3	8	3		
3	A	1	Total	C	H	O	5	0
			10	3	6	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	2	0
			14	3	8	3		
3	B	1	Total	C	H	O	2	0
			14	3	8	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	24	Total	H	O	48	0
			72	48	24		
4	B	5	Total	H	O	10	0
			15	10	5		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: gp41-5



- Molecule 2: alpha/beta-peptide derived from gp41 CHR domain sequence



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.81Å 57.62Å 101.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.80 24.95 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (25.00-2.80) 99.4 (24.95-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.17 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, $R_{free}$	0.193 , 0.285 0.193 , 0.279	Depositor DCC
$R_{free}$ test set	355 reflections (5.12%)	DCC
Wilson B-factor (Å <sup>2</sup> )	47.4	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 66.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3835	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: XPC, GOL, XCP, B3E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/1599	0.67	0/2156
2	B	0.60	0/202	0.71	0/260
All	All	0.64	0/1801	0.67	0/2416

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	9

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	11	ALA	Peptide
2	B	14	ALA	Peptide
2	B	18	GLU	Peptide
2	B	21	ILE	Peptide
2	B	25	GLN	Peptide
2	B	28	GLN	Peptide
2	B	32	GLU	Peptide
2	B	4	GLU	Peptide
2	B	7	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1579	1563	1553	37	1
2	B	281	273	261	5	0
3	A	16	22	18	2	0
3	B	6	8	8	0	0
4	A	24	48	0	2	0
4	B	5	10	0	1	0
All	All	1911	1924	1840	38	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120[B]:GLY:O	1:A:121[B]:SER:HB2	1.71	0.91
1:A:166:VAL:HG21	2:B:35:LEU:HD13	1.59	0.84
1:A:29:LYS:NZ	1:A:47:ASP:OD1	2.15	0.79
1:A:120[B]:GLY:O	1:A:121[B]:SER:CB	2.31	0.79
1:A:17:GLN:HE21	1:A:180:GLN:HG2	1.54	0.73
1:A:102:LEU:HD21	1:A:183:LEU:HD13	1.71	0.71
1:A:1:SER:O	1:A:1:SER:OG	2.10	0.69
1:A:18:GLN:NE2	1:A:57:ILE:HG21	2.08	0.69
1:A:20:LEU:HD13	2:B:13:TYR:HB2	1.75	0.66
1:A:57:ILE:HD13	1:A:98:GLN:HG3	1.84	0.60
1:A:35:ILE:HD11	1:A:197:ILE:HD11	1.85	0.58
1:A:14:ILE:HG23	1:A:98:GLN:OE1	2.04	0.58
1:A:194:GLN:HG2	1:A:198:LEU:HD12	1.84	0.58
1:A:154:GLN:O	1:A:157:LEU:HD23	2.05	0.56
1:A:99:GLN:HA	1:A:99:GLN:OE1	2.05	0.56
1:A:157:LEU:HD12	1:A:158:GLY:N	2.21	0.56
1:A:54:THR:O	1:A:57:ILE:HG22	2.05	0.56
1:A:117:LEU:HD22	3:A:199:GOL:H31	1.88	0.55
2:B:8:XPC:HE	4:B:45:HOH:O	2.07	0.53
1:A:110:LYS:NZ	1:A:128:ASP:OD1	2.46	0.48
1:A:57:ILE:CD1	1:A:98:GLN:HG3	2.44	0.48

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:76:LEU:HD23	1:A:76:LEU:N	2.29	0.48
1:A:82:SER:HA	1:A:156:LEU:HD21	1.95	0.48
1:A:134:TYR:HB3	1:A:182:LEU:HD13	1.96	0.47
1:A:163:SER:HA	2:B:35:LEU:HD22	1.96	0.47
1:A:102:LEU:HD13	1:A:182:LEU:HD23	1.98	0.46
1:A:34:ARG:HB3	4:A:208:HOH:O	2.16	0.46
1:A:103:GLN:NE2	1:A:139:HIS:NE2	2.65	0.45
1:A:163:SER:CA	2:B:35:LEU:HD22	2.47	0.44
1:A:116:ILE:HD11	1:A:193:LEU:CD2	2.47	0.44
1:A:34:ARG:HG3	1:A:198:LEU:HD21	1.99	0.44
1:A:116:ILE:HD11	1:A:193:LEU:HD23	1.99	0.44
1:A:1:SER:C	1:A:3:ILE:N	2.71	0.43
1:A:76:LEU:HG	3:A:201:GOL:H32	2.00	0.43
1:A:134:TYR:CB	1:A:182:LEU:HD13	2.50	0.42
1:A:57:ILE:HD12	1:A:97:ALA:HB3	2.03	0.41
1:A:18:GLN:HE21	1:A:57:ILE:HG12	1.86	0.41
1:A:70:LYS:HE3	4:A:217:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:SER:H	1:A:140:SER:HG[3_555]	1.10	0.50

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	201/198 (102%)	195 (97%)	4 (2%)	2 (1%)	18	50
2	B	23/40 (58%)	19 (83%)	4 (17%)	0	100	100
All	All	224/238 (94%)	214 (96%)	8 (4%)	2 (1%)	32	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	121[A]	SER
1	A	121[B]	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/167 (99%)	159 (96%)	6 (4%)	40	74
2	B	19/21 (90%)	18 (95%)	1 (5%)	26	59
All	All	184/188 (98%)	177 (96%)	7 (4%)	38	72

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LEU
1	A	40	SER
1	A	55	SER
1	A	57	ILE
1	A	144	GLU
1	A	194	GLN
2	B	18	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	18	GLN
1	A	19	HIS
1	A	22	GLN
1	A	103	GLN
1	A	147	ASN
1	A	194	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	B3E	B	12	2	5,5,10	0.57	0	4,5,12	1.91	1 (25%)
2	XCP	B	15	2	8,8,9	0.75	0	6,10,12	1.50	2 (33%)
2	XCP	B	19	2	8,8,9	1.38	1 (12%)	6,10,12	0.90	0
2	XPC	B	22	2	4,8,9	0.85	0	4,10,12	1.17	1 (25%)
2	B3E	B	26	2	6,9,10	0.48	0	4,10,12	1.72	1 (25%)
2	B3E	B	29	2	6,9,10	0.40	0	4,10,12	1.42	0
2	XCP	B	33	2	8,8,9	0.74	0	6,10,12	1.34	1 (16%)
2	XCP	B	5	2	8,8,9	0.67	0	6,10,12	1.20	0
2	XPC	B	8	2	4,8,9	0.99	0	4,10,12	2.16	2 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	B3E	B	12	2	-	0/3/3/9	0/0/0/0
2	XCP	B	15	2	-	0/1/12/14	0/1/1/1
2	XCP	B	19	2	-	0/1/12/14	0/1/1/1
2	XPC	B	22	2	-	0/1/12/14	0/1/1/1
2	B3E	B	26	2	-	0/6/8/9	0/0/0/0
2	B3E	B	29	2	-	0/6/8/9	0/0/0/0
2	XCP	B	33	2	-	0/1/12/14	0/1/1/1
2	XCP	B	5	2	-	0/1/12/14	0/1/1/1
2	XPC	B	8	2	-	0/1/12/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	19	XCP	CA-C	3.59	1.55	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	12	B3E	CG-CB-CA	-3.39	108.10	111.97
2	B	33	XCP	O-C-CA	-2.53	119.25	125.29
2	B	8	XPC	O-C-CA	-2.51	119.30	125.29
2	B	22	XPC	O-C-CA	-2.08	120.33	125.29
2	B	15	XCP	O-C-CA	-2.01	120.50	125.29
2	B	15	XCP	CE-CA-CB	2.24	105.39	100.96
2	B	26	B3E	CB-CA-C	2.31	115.75	112.25
2	B	8	XPC	CG-CB-CA	2.99	106.78	104.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	8	XPC	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	199	-	5,5,5	0.55	0	5,5,5	0.53	0
3	GOL	A	200	-	3,3,5	0.55	0	3,3,5	0.33	0
3	GOL	A	201	-	5,5,5	0.35	0	5,5,5	0.20	0
3	GOL	B	40	-	5,5,5	0.55	0	5,5,5	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	199	-	-	0/4/4/4	0/0/0/0
3	GOL	A	200	-	-	0/0/0/4	0/0/0/0
3	GOL	A	201	-	-	0/4/4/4	0/0/0/0
3	GOL	B	40	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	199	GOL	1	0
3	A	201	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	198/198 (100%)	-0.13	15 (7%) 15 8	21, 39, 81, 97	0
2	B	25/40 (62%)	0.01	1 (4%) 39 28	33, 50, 84, 102	0
All	All	223/238 (93%)	-0.11	16 (7%) 16 9	21, 39, 82, 102	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	160	SER	7.5
1	A	79	SER	6.4
1	A	77	GLY	5.0
2	B	35	LEU	4.9
1	A	161	GLY	4.7
1	A	198	LEU	4.4
1	A	159	GLY	4.1
1	A	39	GLY	3.2
1	A	78	GLY	3.0
1	A	80	GLY	2.8
1	A	38	GLY	2.8
1	A	157	LEU	2.8
1	A	1	SER	2.4
1	A	37	SER	2.3
1	A	162	GLY	2.1
1	A	155	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	B3E	B	26	10/11	0.87	0.36	-	51,56,74,78	0
2	B3E	B	29	10/11	0.86	0.36	-	54,63,85,88	0
2	XCP	B	5	8/9	0.89	0.21	-	53,61,67,68	0
2	XPC	B	8	8/9	0.96	0.12	-	43,50,56,57	0
2	XPC	B	22	8/9	0.94	0.14	-	41,44,48,49	0
2	B3E	B	12	6/11	0.97	0.14	-	39,43,48,49	2
2	XCP	B	19	8/9	0.94	0.17	-	40,45,47,47	0
2	XCP	B	15	8/9	0.97	0.16	-	34,38,42,43	0
2	XCP	B	33	8/9	0.83	0.47	-	81,87,91,91	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	A	199	6/6	0.83	0.24	1.96	56,62,69,70	2
3	GOL	B	40	6/6	0.86	0.26	0.81	35,39,46,50	2
3	GOL	A	200	4/6	0.93	0.20	0.59	2,44,49,50	5
3	GOL	A	201	6/6	0.90	0.18	-0.35	74,77,80,81	2

### 6.5 Other polymers [i](#)

There are no such residues in this entry.