



Full wwPDB X-ray Structure Validation Report i

Feb 12, 2017 – 07:18 pm GMT

PDB ID : 4O41
Title : Amide linked RNA
Authors : Pallan, P.S.; Egli, M.
Deposited on : 2013-12-18
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1131 (1.24-1.16)
Clashscore	112137	1201 (1.24-1.16)
RSRZ outliers	101464	1132 (1.24-1.16)
RNA backbone	2435	1043 (2.60-0.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 1365 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called AMIDE LINKED RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			
1	B	12	Total	C	N	O	P	0	3	0
			320	145	55	106	14			
1	C	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			
1	D	11	Total	C	N	O	P	0	0	0
			253	115	44	83	11			

- Molecule 2 is STRONTIUM ION (three-letter code: SR) (formula: Sr).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Sr 2 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	71	Total O 71 71	0	0
3	B	76	Total O 76 76	0	0
3	C	71	Total O 71 71	0	0
3	D	66	Total O 66 66	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AMIDE LINKED RNA



- Molecule 1: AMIDE LINKED RNA



- Molecule 1: AMIDE LINKED RNA



- Molecule 1: AMIDE LINKED RNA



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	22.36 Å 34.33 Å 43.30 Å 110.03° 94.24° 95.83°	Depositor
Resolution (Å)	28.00 – 1.20 20.69 – 1.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (28.00-1.20) 84.0 (20.69-1.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	5.81 (at 1.20 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R , R_{free}	0.147 , 0.212 0.144 , 0.205	Depositor DCC
R_{free} test set	1797 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	11.5	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 66.3	EDS
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1365	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: URU, SR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.14	0/237	1.67	4/365 (1.1%)
1	B	1.64	2/303 (0.7%)	2.14	12/466 (2.6%)
1	C	1.07	1/237 (0.4%)	1.78	5/365 (1.4%)
1	D	1.21	1/237 (0.4%)	1.98	13/365 (3.6%)
All	All	1.31	4/1014 (0.4%)	1.92	34/1561 (2.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	204[A]	G	C2'-O2'	-15.67	1.21	1.41
1	B	204[B]	G	C2'-O2'	-15.67	1.21	1.41
1	D	404	G	C2'-O2'	-5.16	1.34	1.41
1	C	311	C	C2'-O2'	5.02	1.48	1.41

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	201	U	C5-C4-O4	-9.37	120.28	125.90
1	B	204[A]	G	C3'-C2'-C1'	-9.20	94.14	101.50
1	B	204[B]	G	C3'-C2'-C1'	-9.20	94.14	101.50
1	D	409	G	C2-N3-C4	8.39	116.09	111.90
1	B	209	G	C2-N3-C4	7.55	115.68	111.90
1	D	409	G	C6-N1-C2	-7.48	120.61	125.10
1	D	409	G	C5-C6-N1	7.27	115.13	111.50
1	B	201	U	N3-C4-O4	6.82	124.17	119.40
1	B	201	U	C2-N3-C4	-6.71	122.97	127.00
1	D	409	G	C5-N7-C8	6.70	107.65	104.30
1	C	302	G	N1-C6-O6	-6.47	116.02	119.90
1	D	412	U	C2-N3-C4	6.25	130.75	127.00
1	B	201	U	C3'-C2'-C1'	-5.91	96.77	101.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	A	C2-N3-C4	5.79	113.50	110.60
1	D	402	G	C5-C6-O6	5.71	132.03	128.60
1	C	313	C	N3-C2-O2	5.67	125.87	121.90
1	D	413	C	C6-N1-C2	-5.66	118.04	120.30
1	A	111	C	N3-C2-O2	5.61	125.83	121.90
1	B	209	G	C5-C6-O6	-5.56	125.26	128.60
1	B	204[A]	G	P-O3'-C3'	5.47	126.26	119.70
1	B	204[B]	G	P-O3'-C3'	5.47	126.26	119.70
1	A	110	G	C2-N3-C4	5.43	114.62	111.90
1	D	404	G	C5-C6-O6	-5.43	125.34	128.60
1	A	113	C	C5-C4-N4	5.27	123.89	120.20
1	D	409	G	N3-C4-C5	-5.26	125.97	128.60
1	D	409	G	N7-C8-N9	-5.23	110.48	113.10
1	C	312	U	N3-C4-C5	5.23	117.74	114.60
1	B	202	G	N7-C8-N9	-5.18	110.51	113.10
1	C	313	C	N1-C2-O2	-5.18	115.79	118.90
1	B	201	U	N1-C2-N3	5.17	118.00	114.90
1	D	404	G	C5-C6-N1	5.15	114.08	111.50
1	C	308	C	N3-C4-N4	-5.13	114.41	118.00
1	D	402	G	C5-C6-N1	-5.10	108.95	111.50
1	D	413	C	C4-C5-C6	5.05	119.93	117.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	253	0	131	0	0
1	B	320	0	155	2	0
1	C	253	0	133	0	0
1	D	253	0	133	0	0
2	A	2	0	0	0	0
3	A	71	0	0	0	0
3	B	76	0	0	0	0
3	C	71	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	66	0	0	0	0
All	All	1365	0	552	2	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (2) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:204[B]:G:N9	1:B:204[B]:G:C1'	2.00	1.25
1:B:204[B]:G:N9	1:B:204[B]:G:O4'	2.46	0.48

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [\(i\)](#)

There are no protein molecules in this entry.

5.3.3 RNA [\(i\)](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	8/12 (66%)	0	0
1	B	7/12 (58%)	0	0
1	C	8/12 (66%)	0	0
1	D	8/12 (66%)	0	0
All	All	31/48 (64%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	URU	A	106	1	31,42,43	1.22	4 (12%)	29,61,64	4.17	7 (24%)
1	URU	B	206[A]	1	31,42,43	1.44	6 (19%)	29,61,64	4.31	10 (34%)
1	URU	B	206[B]	1	31,42,43	1.42	6 (19%)	29,61,64	4.22	9 (31%)
1	URU	C	306	1	31,42,43	1.26	3 (9%)	29,61,64	4.36	10 (34%)
1	URU	D	406	1	31,42,43	1.61	5 (16%)	29,61,64	4.44	9 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	URU	A	106	1	-	0/12/54/55	0/4/4/4
1	URU	B	206[A]	1	-	0/12/54/55	0/4/4/4
1	URU	B	206[B]	1	-	0/12/54/55	0/4/4/4
1	URU	C	306	1	-	0/12/54/55	0/4/4/4
1	URU	D	406	1	-	0/12/54/55	0/4/4/4

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	206[A]	URU	O2'-C2'	-2.60	1.37	1.43
1	B	206[B]	URU	O2'-C2'	-2.60	1.37	1.43
1	D	406	URU	C16-C15	-2.40	1.32	1.38
1	B	206[A]	URU	C6-C5	-2.06	1.33	1.38
1	B	206[B]	URU	C6-C5	-2.06	1.33	1.38
1	A	106	URU	C16-N11	2.00	1.38	1.35
1	B	206[A]	URU	C16-N11	2.14	1.38	1.35
1	B	206[B]	URU	C16-N11	2.14	1.38	1.35
1	D	406	URU	C4-N3	2.35	1.37	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	306	URU	C4-N3	2.44	1.37	1.33
1	B	206[A]	URU	C4-N3	2.57	1.37	1.33
1	B	206[B]	URU	C4-N3	2.57	1.37	1.33
1	A	106	URU	C4-N3	2.62	1.37	1.33
1	B	206[A]	URU	C14-N13	3.06	1.38	1.33
1	B	206[B]	URU	C14-N13	3.06	1.38	1.33
1	D	406	URU	C-NZ'	3.07	1.40	1.33
1	A	106	URU	C-NZ'	3.07	1.40	1.33
1	A	106	URU	C14-N13	3.09	1.38	1.33
1	D	406	URU	C14-N13	3.48	1.39	1.33
1	C	306	URU	C-NZ'	3.89	1.42	1.33
1	C	306	URU	C14-N13	4.02	1.40	1.33
1	B	206[B]	URU	C-NZ'	4.12	1.43	1.33
1	B	206[A]	URU	C-NZ'	4.50	1.44	1.33
1	D	406	URU	OW'-CW'	5.76	1.56	1.43

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206[A]	URU	C3'-C6'-C	-5.35	104.29	113.22
1	D	406	URU	O3-C-NZ'	-4.15	115.06	122.97
1	B	206[B]	URU	CY'-CZ'-NZ'	-4.04	104.64	112.68
1	A	106	URU	C5-C4-N3	-3.87	113.87	123.12
1	C	306	URU	C15-C14-N13	-3.77	114.11	123.12
1	B	206[A]	URU	C15-C14-N13	-3.76	114.13	123.12
1	B	206[B]	URU	C15-C14-N13	-3.76	114.13	123.12
1	D	406	URU	C15-C14-N13	-3.73	114.20	123.12
1	D	406	URU	CZ'-NZ'-C	-3.59	114.44	122.73
1	A	106	URU	C15-C14-N13	-3.28	115.28	123.12
1	D	406	URU	C5-C4-N3	-3.16	115.58	123.12
1	A	106	URU	CY'-CZ'-NZ'	-3.11	106.49	112.68
1	C	306	URU	CY'-CZ'-NZ'	-3.02	106.67	112.68
1	B	206[A]	URU	C5-C4-N3	-2.94	116.10	123.12
1	B	206[B]	URU	C5-C4-N3	-2.94	116.10	123.12
1	C	306	URU	OW'-CW'-CX'	-2.89	102.58	111.83
1	C	306	URU	C5-C4-N3	-2.87	116.27	123.12
1	B	206[A]	URU	O3-C-NZ'	-2.86	117.52	122.97
1	D	406	URU	OW'-CW'-CX'	-2.85	102.70	111.83
1	D	406	URU	C3'-C6'-C	-2.64	108.81	113.22
1	C	306	URU	CZ'-NZ'-C	-2.55	116.85	122.73
1	B	206[B]	URU	CZ'-NZ'-C	-2.49	116.98	122.73
1	B	206[A]	URU	CZ'-NZ'-C	-2.38	117.25	122.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	206[A]	URU	CW'-CX'-CY'	-2.19	98.36	102.62
1	B	206[B]	URU	CW'-CX'-CY'	-2.19	98.36	102.62
1	A	106	URU	O3-C-C6'	2.03	124.47	121.42
1	B	206[B]	URU	O3-C-C6'	2.06	124.52	121.42
1	A	106	URU	OX'-CX'-CW'	2.11	118.60	111.83
1	C	306	URU	O3-C-C6'	2.12	124.62	121.42
1	C	306	URU	OX'-CX'-CW'	2.14	118.68	111.83
1	B	206[A]	URU	O3-C-C6'	3.36	126.48	121.42
1	B	206[A]	URU	CY'-OY'-CV'	3.68	113.69	109.77
1	B	206[B]	URU	CY'-OY'-CV'	3.68	113.69	109.77
1	C	306	URU	CY'-OY'-CV'	3.83	113.84	109.77
1	D	406	URU	O3-C-C6'	6.73	131.56	121.42
1	D	406	URU	C14-N13-C12	12.55	124.91	114.13
1	B	206[A]	URU	C4-N3-C2	13.25	125.51	114.13
1	B	206[B]	URU	C4-N3-C2	13.25	125.51	114.13
1	A	106	URU	C14-N13-C12	13.53	125.75	114.13
1	C	306	URU	C4-N3-C2	14.62	126.69	114.13
1	C	306	URU	C14-N13-C12	15.98	127.86	114.13
1	B	206[A]	URU	C14-N13-C12	15.99	127.87	114.13
1	B	206[B]	URU	C14-N13-C12	15.99	127.87	114.13
1	A	106	URU	C4-N3-C2	16.17	128.02	114.13
1	D	406	URU	C4-N3-C2	16.48	128.29	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	10/12 (83%)	-0.49	0 [100] [100]	12, 15, 16, 18	0
1	B	11/12 (91%)	-0.11	1 (9%) [10] [10]	12, 15, 18, 19	1 (9%)
1	C	10/12 (83%)	-0.43	0 [100] [100]	12, 15, 19, 19	0
1	D	10/12 (83%)	-0.40	0 [100] [100]	12, 17, 22, 23	0
All	All	41/48 (85%)	-0.35	1 (2%) [59] [55]	12, 15, 21, 23	1 (2%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	201	U	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	URU	B	206[A]	39/40	0.97	0.07	-	10,13,18,20	7
1	URU	B	206[B]	39/40	0.97	0.07	-	10,13,18,27	7
1	URU	C	306	39/40	0.98	0.06	-	11,13,18,19	0
1	URU	A	106	39/40	0.98	0.05	-	9,11,15,17	0
1	URU	D	406	39/40	0.97	0.07	-	11,14,18,22	0

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SR	A	701	1/1	1.00	0.02	-	13,13,13,13	1
2	SR	A	702	1/1	1.00	0.02	-	13,13,13,13	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.