



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:33 am GMT

PDB ID : 3O6M  
Title : Anti-Tat HIV 11H6H1 Fab' complexed with a 9-mer Tat peptide  
Authors : Serriere, J.; Gouet, P.; Guillon, C.  
Deposited on : 2010-07-29  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	219	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 68%, green 27%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>%</span> <span>68%</span> <span>27%</span> <span>..</span> </div> </div>
2	H	219	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 5%, orange 1%, yellow 74%, green 21%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>5%</span> <span>74%</span> <span>21%</span> <span>..</span> </div> </div>
3	C	9	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 11%, orange 1%, yellow 78%, green 22%, grey 0%);"></div> <div style="display: flex; justify-content: space-between; margin-top: 2px;"> <span>11%</span> <span>78%</span> <span>22%</span> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3507 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 11H6H1 Fab' light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	216	Total	C	N	O	S	0	0	0
			1678	1054	284	334	6			

- Molecule 2 is a protein called 11H6H1 Fab' heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	0	0	0
			1648	1041	269	330	8			

- Molecule 3 is a protein called Protein Tat 9-mer peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			80	55	14	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	L	40	Total	O	0	0
			40	40		
4	H	58	Total	O	0	0
			58	58		
4	C	3	Total	O	0	0
			3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 11H6H1 Fab' light chain



- Molecule 2: 11H6H1 Fab' heavy chain



- Molecule 3: Protein Tat 9-mer peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	123.84Å 69.72Å 62.16Å 90.00° 109.94° 90.00°	Depositor
Resolution (Å)	19.97 – 2.40 19.97 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.97-2.40) 99.8 (19.97-2.40)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.07 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.211 , 0.290 0.211 , 0.288	Depositor DCC
$R_{free}$ test set	1002 reflections (5.41%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.5	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 38.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3507	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.90	0/1716	0.92	3/2329 (0.1%)
2	H	0.93	1/1692 (0.1%)	0.90	2/2312 (0.1%)
3	C	0.93	1/85 (1.2%)	0.62	0/115
All	All	0.92	2/3493 (0.1%)	0.91	5/4756 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
2	H	0	1
All	All	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	144	CYS	CB-SG	-5.72	1.72	1.81
3	C	11	TRP	CE3-CZ3	5.47	1.47	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	181	LEU	CA-CB-CG	7.54	132.65	115.30
1	L	41	LEU	CA-CB-CG	6.82	130.97	115.30
1	L	65	ASP	CB-CG-OD2	6.49	124.14	118.30
2	H	132	SER	N-CA-C	5.36	125.47	111.00
1	L	33	ASP	CB-CG-OD1	5.14	122.93	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	H	193	PRO	Peptide
1	L	206	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1678	0	1642	50	0
2	H	1648	0	1591	37	0
3	C	80	0	81	2	0
4	C	3	0	0	0	0
4	H	58	0	0	0	0
4	L	40	0	0	0	0
All	All	3507	0	3314	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:95:GLN:NE2	1:L:98:HIS:H	1.41	1.17
1:L:95:GLN:HE22	1:L:98:HIS:H	1.02	0.95
1:L:10:THR:HG22	1:L:108:LYS:HB3	1.49	0.94
1:L:95:GLN:HE22	1:L:98:HIS:N	1.71	0.88
1:L:95:GLN:NE2	1:L:98:HIS:N	2.23	0.87
2:H:152:GLU:HG3	2:H:153:PRO:HA	1.58	0.85
2:H:197:VAL:H	2:H:213:LYS:HZ3	1.24	0.84
2:H:159:ASN:ND2	2:H:198:THR:H	1.77	0.82
2:H:159:ASN:HD21	2:H:198:THR:H	1.29	0.81
2:H:130:PRO:O	2:H:217:ARG:HD2	1.85	0.77
1:L:12:SER:HB3	1:L:110:GLU:OE1	1.84	0.76
1:L:216:ARG:HH11	1:L:216:ARG:HG3	1.50	0.76
1:L:168:TRP:CD1	1:L:168:TRP:N	2.54	0.75
1:L:1:ASP:OD1	1:L:1:ASP:N	2.19	0.74
1:L:29:LEU:HD22	1:L:95:GLN:HG3	1.69	0.74
1:L:29:LEU:CD2	1:L:95:GLN:HG3	2.17	0.73

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:129:ALA:O	2:H:217:ARG:NH1	2.23	0.72
2:H:136:THR:O	2:H:138:SER:N	2.24	0.71
1:L:154:LYS:HD2	1:L:200:GLU:OE2	1.92	0.70
1:L:212:LYS:NZ	2:H:135:GLN:HG3	2.07	0.68
1:L:216:ARG:NH1	1:L:216:ARG:HG3	2.09	0.67
2:H:151:PRO:O	2:H:203:HIS:HE1	1.80	0.65
2:H:152:GLU:HG3	2:H:153:PRO:CA	2.27	0.64
2:H:176:SER:O	2:H:177:ASP:HB2	1.99	0.62
1:L:150:ASN:HD22	1:L:202:THR:HB	1.64	0.62
1:L:141:LEU:N	1:L:141:LEU:HD12	2.13	0.62
1:L:110:GLU:HG3	1:L:178:TYR:OH	2.00	0.62
1:L:31:TYR:CZ	3:C:10:PRO:HB3	2.36	0.61
2:H:1:GLU:O	2:H:1:GLU:HG2	2.00	0.60
1:L:31:TYR:CE2	3:C:10:PRO:HB3	2.36	0.60
2:H:203:HIS:HD2	2:H:206:SER:OG	1.83	0.60
2:H:159:ASN:HD21	2:H:198:THR:N	1.99	0.59
2:H:175:GLN:HG3	2:H:175:GLN:O	2.04	0.57
1:L:212:LYS:HZ1	2:H:135:GLN:HG3	1.69	0.56
1:L:154:LYS:HB2	1:L:198:THR:HB	1.87	0.56
1:L:215:ASN:O	1:L:216:ARG:HB2	2.05	0.56
1:L:155:ILE:HD12	1:L:160:ARG:HD2	1.87	0.56
2:H:40:THR:HB	2:H:41:PRO:HD2	1.89	0.54
1:L:29:LEU:HD21	1:L:95:GLN:HG3	1.90	0.54
2:H:177:ASP:O	2:H:178:LEU:HD23	2.07	0.53
1:L:31:TYR:O	1:L:33:ASP:N	2.35	0.53
1:L:95:GLN:HE21	1:L:98:HIS:H	1.49	0.53
1:L:216:ARG:HH11	1:L:216:ARG:CG	2.21	0.53
1:L:29:LEU:HD22	1:L:95:GLN:CG	2.36	0.53
2:H:134:ALA:HB3	2:H:137:ASN:HA	1.91	0.53
1:L:13:VAL:HG22	1:L:14:THR:H	1.74	0.52
1:L:95:GLN:HE21	1:L:97:THR:N	2.07	0.52
2:H:125:VAL:O	2:H:212:LYS:HE3	2.09	0.52
1:L:11:LEU:HD23	1:L:109:LEU:HD11	1.91	0.52
1:L:212:LYS:HZ3	2:H:135:GLN:HG3	1.76	0.50
1:L:187:THR:OG1	1:L:190:GLU:HG3	2.12	0.49
2:H:137:ASN:HD22	2:H:137:ASN:H	1.61	0.49
2:H:191:THR:O	2:H:195:GLU:HG2	2.13	0.48
1:L:164:VAL:HG12	1:L:165:LEU:N	2.27	0.48
1:L:13:VAL:HG22	1:L:14:THR:N	2.29	0.47
1:L:13:VAL:CG2	1:L:14:THR:H	2.27	0.47
2:H:136:THR:C	2:H:138:SER:N	2.68	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:136:SER:HA	1:L:184:LEU:O	2.14	0.46
1:L:95:GLN:NE2	1:L:97:THR:N	2.64	0.45
2:H:6:GLU:HA	2:H:21:SER:O	2.17	0.45
1:L:109:LEU:HD12	1:L:109:LEU:HA	1.69	0.45
2:H:136:THR:C	2:H:138:SER:H	2.20	0.44
1:L:215:ASN:O	1:L:216:ARG:CB	2.66	0.44
1:L:42:LEU:HD13	1:L:91:TYR:CZ	2.52	0.44
2:H:203:HIS:CD2	2:H:206:SER:OG	2.69	0.44
2:H:51:ILE:CD1	2:H:71:SER:HA	2.48	0.43
1:L:125:PRO:HG2	1:L:135:ALA:HB1	1.99	0.43
1:L:112:LYS:HA	1:L:145:TYR:OH	2.18	0.43
1:L:193:ARG:HB2	1:L:193:ARG:HE	1.69	0.43
1:L:208:SER:HA	1:L:209:PRO:HD3	1.92	0.43
2:H:197:VAL:H	2:H:213:LYS:NZ	2.06	0.42
1:L:31:TYR:C	1:L:33:ASP:H	2.18	0.42
1:L:37:TYR:CD2	1:L:37:TYR:N	2.87	0.42
2:H:152:GLU:CG	2:H:153:PRO:HA	2.41	0.42
2:H:196:THR:HA	2:H:213:LYS:NZ	2.35	0.42
2:H:42:GLU:HG3	2:H:44:ARG:HH12	1.85	0.41
2:H:196:THR:HA	2:H:213:LYS:HZ1	1.84	0.41
1:L:127:SER:O	1:L:131:THR:HG23	2.21	0.41
2:H:137:ASN:HD22	2:H:137:ASN:N	2.18	0.41
2:H:175:GLN:CG	2:H:175:GLN:O	2.65	0.41
1:L:14:THR:O	1:L:15:ILE:C	2.58	0.41
2:H:158:TRP:CZ3	2:H:199:CYS:HB3	2.57	0.40
2:H:22:CYS:HB3	2:H:79:LEU:HB3	2.02	0.40
1:L:21:ILE:HG12	1:L:107:THR:HG21	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	214/219 (98%)	198 (92%)	12 (6%)	4 (2%)	9	11
2	H	216/219 (99%)	200 (93%)	11 (5%)	5 (2%)	7	8
3	C	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	437/447 (98%)	404 (92%)	24 (6%)	9 (2%)	8	9

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	207	THR
2	H	132	SER
2	H	137	ASN
1	L	32	SER
2	H	195	GLU
1	L	66	ARG
2	H	130	PRO
2	H	134	ALA
1	L	158	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	192/195 (98%)	175 (91%)	17 (9%)	11	17
2	H	187/188 (100%)	169 (90%)	18 (10%)	10	14
3	C	9/9 (100%)	9 (100%)	0	100	100
All	All	388/392 (99%)	353 (91%)	35 (9%)	11	16

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	1	ASP
1	L	27	GLN
1	L	30	LEU
1	L	41	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	L	45	PRO
1	L	82	ARG
1	L	108	LYS
1	L	121	SER
1	L	128	GLU
1	L	147	LYS
1	L	148	ASP
1	L	152	LYS
1	L	158	SER
1	L	168	TRP
1	L	186	LEU
1	L	204	LYS
1	L	216	ARG
2	H	1	GLU
2	H	12	VAL
2	H	31	SER
2	H	109	GLN
2	H	119	LYS
2	H	132	SER
2	H	136	THR
2	H	137	ASN
2	H	138	SER
2	H	141	THR
2	H	163	LEU
2	H	173	VAL
2	H	174	LEU
2	H	181	LEU
2	H	189	SER
2	H	194	SER
2	H	209	LYS
2	H	218	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	47	GLN
1	L	95	GLN
1	L	150	ASN
2	H	137	ASN
2	H	159	ASN
2	H	203	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	216/219 (98%)	-0.07	3 (1%) 75 74	16, 34, 47, 58	0
2	H	218/219 (99%)	-0.02	11 (5%) 30 28	14, 27, 52, 79	0
3	C	9/9 (100%)	0.17	1 (11%) 6 5	27, 35, 45, 51	0
All	All	443/447 (99%)	-0.04	15 (3%) 46 44	14, 31, 48, 79	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	136	THR	9.9
2	H	134	ALA	7.8
2	H	133	ALA	6.9
2	H	135	GLN	6.2
2	H	218	ASP	5.5
2	H	132	SER	4.3
2	H	131	GLY	3.9
2	H	75	ALA	3.1
2	H	137	ASN	2.4
1	L	193	ARG	2.2
1	L	162	ASN	2.2
3	C	6	PRO	2.2
1	L	207	THR	2.1
2	H	54	GLY	2.1
2	H	193	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.