



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2018 – 07:01 AM EST

PDB ID : 5O8H
Title : Crystal structure of R. ruber ADH-A, mutant Y294F, W295A, F43H, H39Y
Authors : Dobritsch, D.; Maurer, D.; Hamnevik, E.; Reddy Enugala, T.; Widersten, M.
Deposited on : 2017-06-13
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	<div> <div>2%</div> <div>97%</div> <div>..</div> </div>
1	B	352	<div> <div>2%</div> <div>96%</div> <div>..</div> </div>
1	C	352	<div> <div>4%</div> <div>96%</div> <div>..</div> </div>
1	D	352	<div> <div>15%</div> <div>95%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	K	C	504	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	S	0	2	0
			2489	1568	439	470	12			
1	B	346	Total	C	N	O	S	0	3	0
			2493	1571	439	471	12			
1	C	346	Total	C	N	O	S	0	2	0
			2482	1564	436	470	12			
1	D	345	Total	C	N	O	S	0	1	0
			2467	1555	432	467	13			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	TYR	HIS	engineered mutation	UNP A0A1Q8I6M1
A	43	HIS	PHE	engineered mutation	UNP A0A1Q8I6M1
A	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
A	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
A	346	THR	-	expression tag	UNP A0A1Q8I6M1
A	347	SER	-	expression tag	UNP A0A1Q8I6M1
A	348	HIS	-	expression tag	UNP A0A1Q8I6M1
A	349	HIS	-	expression tag	UNP A0A1Q8I6M1
A	350	HIS	-	expression tag	UNP A0A1Q8I6M1
A	351	HIS	-	expression tag	UNP A0A1Q8I6M1
A	352	HIS	-	expression tag	UNP A0A1Q8I6M1
B	39	TYR	HIS	engineered mutation	UNP A0A1Q8I6M1
B	43	HIS	PHE	engineered mutation	UNP A0A1Q8I6M1
B	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
B	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
B	346	THR	-	expression tag	UNP A0A1Q8I6M1
B	347	SER	-	expression tag	UNP A0A1Q8I6M1
B	348	HIS	-	expression tag	UNP A0A1Q8I6M1
B	349	HIS	-	expression tag	UNP A0A1Q8I6M1
B	350	HIS	-	expression tag	UNP A0A1Q8I6M1
B	351	HIS	-	expression tag	UNP A0A1Q8I6M1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	352	HIS	-	expression tag	UNP A0A1Q8I6M1
C	39	TYR	HIS	engineered mutation	UNP A0A1Q8I6M1
C	43	HIS	PHE	engineered mutation	UNP A0A1Q8I6M1
C	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
C	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
C	346	THR	-	expression tag	UNP A0A1Q8I6M1
C	347	SER	-	expression tag	UNP A0A1Q8I6M1
C	348	HIS	-	expression tag	UNP A0A1Q8I6M1
C	349	HIS	-	expression tag	UNP A0A1Q8I6M1
C	350	HIS	-	expression tag	UNP A0A1Q8I6M1
C	351	HIS	-	expression tag	UNP A0A1Q8I6M1
C	352	HIS	-	expression tag	UNP A0A1Q8I6M1
D	39	TYR	HIS	engineered mutation	UNP A0A1Q8I6M1
D	43	HIS	PHE	engineered mutation	UNP A0A1Q8I6M1
D	294	PHE	TYR	engineered mutation	UNP A0A1Q8I6M1
D	295	ALA	TRP	engineered mutation	UNP A0A1Q8I6M1
D	346	THR	-	expression tag	UNP A0A1Q8I6M1
D	347	SER	-	expression tag	UNP A0A1Q8I6M1
D	348	HIS	-	expression tag	UNP A0A1Q8I6M1
D	349	HIS	-	expression tag	UNP A0A1Q8I6M1
D	350	HIS	-	expression tag	UNP A0A1Q8I6M1
D	351	HIS	-	expression tag	UNP A0A1Q8I6M1
D	352	HIS	-	expression tag	UNP A0A1Q8I6M1

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Zn 2 2	0	0
2	A	2	Total Zn 2 2	0	0
2	D	2	Total Zn 2 2	0	0
2	C	2	Total Zn 2 2	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total 1 K	0	0
4	C	1	Total 1 K	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	329	Total	O	0	1
			330	330		
6	B	325	Total	O	0	1
			326	326		
6	C	250	Total	O	0	2
			252	252		
6	D	218	Total	O	0	1
			219	219		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol dehydrogenase




- Molecule 1: Alcohol dehydrogenase



- Molecule 1: Alcohol dehydrogenase



- Molecule 1: Alcohol dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.70 Å 105.68 Å 109.62 Å 90.00° 91.51° 90.00°	Depositor
Resolution (Å)	29.63 – 1.96 29.63 – 1.96	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.63-1.96) 99.9 (29.63-1.96)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.96 Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.176 , 0.204 0.184 , 0.210	Depositor DCC
R_{free} test set	5190 reflections (5.09%)	DCC
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.004 for -h,l,k 0.019 for -h,-l,-k 0.028 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11250	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, K, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2542	0.72	3/3470 (0.1%)
1	B	0.50	0/2549	0.74	4/3481 (0.1%)
1	C	0.50	0/2537	0.73	3/3463 (0.1%)
1	D	0.50	0/2519	0.71	2/3439 (0.1%)
All	All	0.50	0/10147	0.72	12/13853 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	205	ASP	CB-CG-OD2	-6.42	112.52	118.30
1	B	330	ARG	NE-CZ-NH2	6.33	123.47	120.30
1	A	314	ASP	CB-CG-OD1	5.81	123.53	118.30
1	B	153	ASP	CB-CG-OD1	5.57	123.31	118.30
1	C	232	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	B	312	ARG	NE-CZ-NH1	-5.32	117.64	120.30
1	C	153	ASP	CB-CG-OD1	5.24	123.01	118.30
1	D	143	LEU	CA-CB-CG	5.22	127.31	115.30
1	B	164	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	A	153	ASP	CB-CG-OD1	5.11	122.90	118.30
1	A	164	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	D	232	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2489	0	2486	2	0
1	B	2493	0	2493	2	0
1	C	2482	0	2485	2	0
1	D	2467	0	2465	5	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	44	0	26	0	0
3	B	44	0	26	0	0
3	C	44	0	26	0	0
3	D	44	0	26	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	6	0	8	0	0
6	A	330	0	0	1	0
6	B	326	0	0	1	0
6	C	252	0	0	0	0
6	D	219	0	0	0	0
All	All	11250	0	10041	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:ASP:OD2	6:A:601:HOH:O	2.13	0.67
1:B:312:ARG:NH2	6:B:602:HOH:O	2.42	0.48
1:A:179:GLY:O	1:A:184:GLY:HA3	2.17	0.45
1:D:34:ALA:HB3	1:D:65:VAL:CG2	2.48	0.43
1:C:179:GLY:O	1:C:184:GLY:HA3	2.17	0.43
1:D:179:GLY:O	1:D:184:GLY:HA3	2.18	0.43
1:B:179:GLY:O	1:B:184:GLY:HA3	2.18	0.42
1:D:65:VAL:HG22	1:D:149:ALA:HB2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:165:VAL:HG11	1:C:265:HIS:CG	2.56	0.41
1:D:71:LEU:HD23	1:D:75:VAL:O	2.21	0.40
1:D:165:VAL:HG11	1:D:265:HIS:CG	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/352 (98%)	340 (98%)	6 (2%)	0	100	100
1	B	347/352 (99%)	341 (98%)	6 (2%)	0	100	100
1	C	346/352 (98%)	340 (98%)	6 (2%)	0	100	100
1	D	344/352 (98%)	338 (98%)	6 (2%)	0	100	100
All	All	1383/1408 (98%)	1359 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/254 (98%)	249 (100%)	1 (0%)	93	92
1	B	251/254 (99%)	250 (100%)	1 (0%)	93	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	250/254 (98%)	249 (100%)	1 (0%)	93	92
1	D	248/254 (98%)	247 (100%)	1 (0%)	93	92
All	All	999/1016 (98%)	995 (100%)	4 (0%)	93	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	272	HIS
1	B	272	HIS
1	C	222	LYS
1	D	272	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 10 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAD	A	503	-	41,48,48	0.85	1 (2%)	43,73,73	1.53	6 (13%)
5	GOL	A	505	-	5,5,5	0.40	0	5,5,5	0.69	0
3	NAD	B	503	-	41,48,48	0.88	2 (4%)	43,73,73	1.47	8 (18%)
3	NAD	C	503	-	41,48,48	0.95	2 (4%)	43,73,73	1.52	5 (11%)
3	NAD	D	503	-	41,48,48	0.86	1 (2%)	43,73,73	1.29	4 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	503	-	-	0/22/62/62	0/5/5/5
5	GOL	A	505	-	-	0/4/4/4	0/0/0/0
3	NAD	B	503	-	-	0/22/62/62	0/5/5/5
3	NAD	C	503	-	-	0/22/62/62	0/5/5/5
3	NAD	D	503	-	-	0/22/62/62	0/5/5/5

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	503	NAD	C5A-C4A	2.41	1.45	1.40
3	A	503	NAD	C5A-C4A	2.62	1.46	1.40
3	C	503	NAD	C2A-N3A	2.67	1.36	1.32
3	C	503	NAD	C5A-C4A	2.83	1.46	1.40
3	B	503	NAD	O4D-C1D	3.06	1.45	1.41
3	D	503	NAD	C5A-C4A	3.37	1.48	1.40

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	503	NAD	N3A-C2A-N1A	-7.15	122.63	128.86
3	A	503	NAD	N3A-C2A-N1A	-6.62	123.09	128.86
3	B	503	NAD	N3A-C2A-N1A	-5.49	124.08	128.86
3	D	503	NAD	N3A-C2A-N1A	-4.37	125.05	128.86
3	D	503	NAD	C4A-C5A-N7A	-3.45	106.07	109.41
3	A	503	NAD	C1B-N9A-C4A	-3.05	121.37	126.64
3	B	503	NAD	O7N-C7N-N7N	-2.63	118.84	122.58
3	A	503	NAD	C4D-O4D-C1D	-2.38	107.23	109.77
3	D	503	NAD	O7N-C7N-N7N	-2.30	119.30	122.58
3	C	503	NAD	O7N-C7N-N7N	-2.29	119.33	122.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	503	NAD	C4A-C5A-N7A	-2.19	107.29	109.41
3	A	503	NAD	C4A-C5A-N7A	-2.18	107.31	109.41
3	B	503	NAD	C4D-O4D-C1D	-2.09	107.54	109.77
3	B	503	NAD	C4B-O4B-C1B	-2.08	107.55	109.77
3	B	503	NAD	C1B-N9A-C4A	-2.04	123.12	126.64
3	A	503	NAD	O2N-PN-O1N	2.03	122.79	112.28
3	C	503	NAD	C3N-C7N-N7N	2.06	120.12	117.77
3	D	503	NAD	C3N-C7N-N7N	2.10	120.17	117.77
3	B	503	NAD	O2A-PA-O1A	2.15	123.40	112.28
3	C	503	NAD	C3N-C2N-N1N	2.28	122.72	120.43
3	B	503	NAD	O2N-PN-O1N	2.36	124.51	112.28
3	B	503	NAD	C3N-C7N-N7N	2.80	120.97	117.77
3	A	503	NAD	C3N-C7N-N7N	2.89	121.08	117.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/352 (98%)	-0.07	8 (2%) 61 71	18, 26, 44, 64	0
1	B	346/352 (98%)	-0.07	7 (2%) 65 75	17, 26, 45, 62	0
1	C	346/352 (98%)	0.27	14 (4%) 39 49	18, 32, 52, 72	0
1	D	345/352 (98%)	0.74	53 (15%) 2 3	19, 34, 68, 85	0
All	All	1383/1408 (98%)	0.22	82 (5%) 23 32	17, 28, 58, 85	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	75	VAL	7.2
1	C	76	THR	6.3
1	D	80	VAL	5.9
1	D	51	GLN	5.2
1	D	50	ALA	5.2
1	A	282	PHE	4.6
1	D	48	PRO	4.5
1	D	11	SER	4.4
1	D	61	GLY	4.2
1	D	76	THR	4.0
1	D	58	LEU	4.0
1	D	53	ALA	3.9
1	D	335	GLY	3.9
1	B	282	PHE	3.8
1	D	9	ILE	3.7
1	D	18	ILE	3.6
1	D	128	TYR	3.6
1	D	336	SER	3.5
1	C	346	THR	3.5
1	D	69	ALA	3.5
1	D	25	PRO	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	47	MET	3.4
1	D	38[A]	CYS	3.3
1	D	321	THR	3.1
1	D	125	MET	3.0
1	C	110	ALA	3.0
1	D	71	LEU	2.9
1	D	81	GLY	2.9
1	D	62	HIS	2.9
1	D	334	GLU	2.9
1	D	42	ILE	2.9
1	D	57	PRO	2.9
1	D	37	LEU	2.9
1	A	50	ALA	2.8
1	D	36	GLY	2.8
1	D	10	GLY	2.8
1	D	344	VAL	2.8
1	D	54	TYR	2.8
1	C	48	PRO	2.7
1	D	56	LEU	2.7
1	C	113	GLY	2.7
1	A	346	THR	2.7
1	D	339	GLY	2.6
1	D	243	VAL	2.6
1	C	50	ALA	2.6
1	D	55	GLY	2.6
1	D	20	THR	2.6
1	C	49	ALA	2.6
1	A	53	ALA	2.5
1	B	183	LEU	2.5
1	D	124	SER	2.5
1	B	62[A]	HIS	2.5
1	D	4	VAL	2.5
1	C	107	THR	2.4
1	C	80	VAL	2.4
1	D	23	PRO	2.4
1	D	24	GLY	2.4
1	D	338	ARG	2.3
1	D	17	ASP	2.3
1	D	126	ALA	2.3
1	D	332	LEU	2.3
1	D	73	GLU	2.3
1	B	273	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	76	THR	2.2
1	D	110	ALA	2.2
1	A	113	GLY	2.2
1	D	22	THR	2.2
1	C	47	MET	2.2
1	A	110	ALA	2.2
1	C	17	ASP	2.2
1	B	51	GLN	2.2
1	D	46	ASP	2.2
1	D	337	ILE	2.1
1	A	183	LEU	2.1
1	A	51	GLN	2.1
1	C	51	GLN	2.1
1	C	9	ILE	2.1
1	B	50	ALA	2.1
1	C	21	PRO	2.1
1	D	59	THR	2.0
1	D	127	GLU	2.0
1	D	320	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	K	C	504	1/1	0.75	0.19	3.21	77,77,77,77	0
4	K	A	504	1/1	0.94	0.10	0.02	63,63,63,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAD	B	503	44/44	0.98	0.09	-0.82	20,22,23,23	0
3	NAD	A	503	44/44	0.98	0.09	-0.89	19,21,22,23	0
3	NAD	D	503	44/44	0.97	0.09	-0.98	23,26,28,31	0
2	ZN	C	501	1/1	1.00	0.03	-1.05	23,23,23,23	0
2	ZN	A	501	1/1	1.00	0.03	-1.19	25,25,25,25	0
3	NAD	C	503	44/44	0.98	0.08	-1.28	22,24,26,26	0
2	ZN	B	501	1/1	1.00	0.02	-1.30	24,24,24,24	0
2	ZN	D	501	1/1	1.00	0.02	-1.36	24,24,24,24	0
2	ZN	B	502	1/1	0.98	0.05	-2.22	57,57,57,57	0
2	ZN	A	502	1/1	0.97	0.05	-2.32	52,52,52,52	0
2	ZN	D	502	1/1	0.94	0.06	-2.86	81,81,81,81	0
2	ZN	C	502	1/1	0.98	0.08	-	61,61,61,61	0
5	GOL	A	505	6/6	0.67	0.26	-	54,57,58,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.