



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 11:21 am GMT

PDB ID : 2OB2  
Title : ppm1 in the absence of 1,8-ANS (cf 1JD)  
Authors : Groves, M.R.; Mueller, I.B.; Kreplin, X.; Mueller-Dieckmann, J.  
Deposited on : 2006-12-18  
Resolution : 1.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6276 (1.94-1.90)
Clashscore	112137	7025 (1.94-1.90)
Ramachandran outliers	110173	6947 (1.94-1.90)
Sidechain outliers	110143	6948 (1.94-1.90)
RSRZ outliers	101464	6332 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	<div> <div>78%</div> <div>20%</div> <div>•</div> </div>
1	B	327	<div> <div>79%</div> <div>20%</div> <div>•</div> </div>
1	C	327	<div> <div>2%</div> <div>65%</div> <div>31%</div> <div>•</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8641 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Leucine carboxyl methyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	327	Total	C	N	O	S	0	0	0
			2636	1676	448	494	18			
1	B	327	Total	C	N	O	S	0	2	0
			2650	1684	451	497	18			
1	C	327	Total	C	N	O	S	0	0	0
			2636	1676	448	494	18			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	202	SMC	CYS	MODIFIED RESIDUE	UNP Q04081
B	202	SMC	CYS	MODIFIED RESIDUE	UNP Q04081
C	202	SMC	CYS	MODIFIED RESIDUE	UNP Q04081

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 26	C 14	N 6	O 5	S 1	0	0
3	B	1	Total 26	C 14	N 6	O 5	S 1	0	0
3	C	1	Total 26	C 14	N 6	O 5	S 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	246	Total	O	0	0
			246	246		
5	B	232	Total	O	0	0
			232	232		
5	C	146	Total	O	0	0
			146	146		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Leucine carboxyl methyltransferase 1



- Molecule 1: Leucine carboxyl methyltransferase 1



- Molecule 1: Leucine carboxyl methyltransferase 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.62Å 110.62Å 161.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.06 – 1.92 33.05 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.06-1.92) 99.7 (33.05-1.92)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.32 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, $R_{free}$	0.206 , 0.268 0.207 , 0.269	Depositor DCC
$R_{free}$ test set	4269 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.4	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 21.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.336 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8641	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4, SAH, SMC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.95	0/2677	1.01	6/3613 (0.2%)
1	B	0.93	0/2697	1.01	6/3639 (0.2%)
1	C	0.87	0/2677	0.96	8/3613 (0.2%)
All	All	0.92	0/8051	0.99	20/10865 (0.2%)

There are no bond length outliers.

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	ASP	CB-CG-OD2	6.78	124.41	118.30
1	C	232	ASP	CB-CG-OD2	6.63	124.27	118.30
1	A	112	MET	CG-SD-CE	6.63	110.81	100.20
1	B	111	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	B	64	ASP	CB-CG-OD2	6.11	123.80	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2636	0	2666	60	0
1	B	2650	0	2682	59	0
1	C	2636	0	2666	79	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	5	0	0	1	0
3	A	26	0	19	0	0
3	B	26	0	19	0	0
3	C	26	0	19	0	0
4	A	12	0	16	1	0
5	A	246	0	0	10	0
5	B	232	0	0	16	0
5	C	146	0	0	10	0
All	All	8641	0	8087	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317:MET:HE3	1:A:318:GLN:NE2	1.66	1.10
1:B:46[A]:ARG:HH21	1:B:46[A]:ARG:HB3	1.06	1.10
1:A:317:MET:HE3	1:A:318:GLN:HE21	0.98	1.09
1:C:317:MET:HE3	1:C:318:GLN:NE2	1.82	0.94
1:A:317:MET:CE	1:A:318:GLN:HE21	1.79	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/327 (99%)	310 (96%)	13 (4%)	1 (0%)	44	33
1	B	326/327 (100%)	318 (98%)	7 (2%)	1 (0%)	44	33
1	C	324/327 (99%)	312 (96%)	11 (3%)	1 (0%)	44	33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	974/981 (99%)	940 (96%)	31 (3%)	3 (0%)	44 33

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	156	ASP
1	C	97	GLU
1	A	273	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	295/295 (100%)	289 (98%)	6 (2%)	60 54
1	B	297/295 (101%)	289 (97%)	8 (3%)	50 41
1	C	295/295 (100%)	283 (96%)	12 (4%)	35 23
All	All	887/885 (100%)	861 (97%)	26 (3%)	49 37

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	210	ASN
1	C	36	ASP
1	C	263	MET
1	B	249	SER
1	B	300	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	121	HIS
1	B	209	ASN
1	C	209	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	117	GLN
1	C	217	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	SMC	A	202	1	6,6,7	0.79	0	3,6,8	2.64	2 (66%)
1	SMC	B	202	1	6,6,7	0.66	0	3,6,8	3.02	2 (66%)
1	SMC	C	202	1	6,6,7	1.34	1 (16%)	3,6,8	1.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SMC	A	202	1	-	0/3/5/7	0/0/0/0
1	SMC	B	202	1	-	0/3/5/7	0/0/0/0
1	SMC	C	202	1	-	0/3/5/7	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	202	SMC	CB-SG	-2.29	1.78	1.80

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	202	SMC	CA-CB-SG	-3.10	109.95	114.22
1	A	202	SMC	CA-CB-SG	-2.28	111.08	114.22
1	A	202	SMC	CS-SG-CB	3.81	107.31	101.19
1	B	202	SMC	CS-SG-CB	3.92	107.48	101.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	A	1001	-	4,4,4	0.85	0	6,6,6	0.51	0
3	SAH	A	1801	-	20,28,28	1.21	3 (15%)	20,40,40	2.43	2 (10%)
4	GOL	A	2002	-	5,5,5	0.57	0	5,5,5	0.64	0
4	GOL	A	2003	-	5,5,5	0.41	0	5,5,5	1.09	0
3	SAH	B	1802	-	20,28,28	1.16	3 (15%)	20,40,40	2.83	6 (30%)
3	SAH	C	1803	-	20,28,28	1.43	3 (15%)	20,40,40	2.77	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1001	-	-	0/0/0/0	0/0/0/0
3	SAH	A	1801	-	-	0/7/31/31	0/3/3/3
4	GOL	A	2002	-	-	0/4/4/4	0/0/0/0
4	GOL	A	2003	-	-	0/4/4/4	0/0/0/0
3	SAH	B	1802	-	-	0/7/31/31	0/3/3/3
3	SAH	C	1803	-	-	0/7/31/31	0/3/3/3

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1801	SAH	O4'-C1'	-2.21	1.38	1.41
3	B	1802	SAH	C2-N1	2.16	1.38	1.33
3	A	1801	SAH	C2-N1	2.42	1.38	1.33
3	B	1802	SAH	C4-N3	2.54	1.39	1.35
3	C	1803	SAH	C4-N3	2.85	1.39	1.35

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1803	SAH	N3-C2-N1	-10.79	119.46	128.86
3	B	1802	SAH	N3-C2-N1	-10.23	119.95	128.86
3	A	1801	SAH	N3-C2-N1	-8.89	121.12	128.86
3	C	1803	SAH	C5'-SD-CG	-4.64	88.22	102.29
3	A	1801	SAH	C5'-SD-CG	-4.04	90.05	102.29

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	PO4	1	0
4	A	2003	GOL	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/327 (99%)	-0.47	0	100 100	7, 14, 28, 41	0
1	B	326/327 (99%)	-0.45	0	100 100	7, 14, 30, 43	0
1	C	326/327 (99%)	-0.10	8 (2%)	58 62	5, 15, 34, 41	0
All	All	978/981 (99%)	-0.34	8 (0%)	86 88	5, 14, 32, 43	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	257	LEU	2.9
1	C	242	ARG	2.9
1	C	255	ARG	2.9
1	C	249	SER	2.8
1	C	239	PRO	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	SMC	B	202	7/8	0.97	0.08	-	6,10,12,16	0
1	SMC	C	202	7/8	0.98	0.07	-	7,9,11,15	0
1	SMC	A	202	7/8	0.97	0.08	-	9,11,12,17	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	2003	6/6	0.92	0.09	1.06	22,27,28,30	0
4	GOL	A	2002	6/6	0.91	0.09	0.01	9,14,15,20	0
3	SAH	C	1803	26/26	0.97	0.07	-1.23	10,17,22,23	0
3	SAH	A	1801	26/26	0.98	0.06	-1.26	2,4,6,8	0
3	SAH	B	1802	26/26	0.99	0.05	-1.95	2,4,6,8	0
2	PO4	A	1001	5/5	0.92	0.17	-	47,48,51,51	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.