



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 10:00 pm GMT

PDB ID : 4OBZ  
Title : Structure of Cathepsin D with inhibitor 2-(3,4-dimethoxyphenyl)-N-[N-(4-methylbenzyl)carbamimidoyl]acetamide  
Authors : Graedler, U.; Czodrowski, P.; Tsaklakidis, C.; Klein, M.; Maskos, K.; Leuthner, B.  
Deposited on : 2014-01-08  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	103	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 85%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>85%</span> <span>9%</span> <span>6%</span> </div> </div>
1	C	103	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 88%, yellow 11%, red 1%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>88%</span> <span>11%</span> </div> </div>
2	B	243	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 85%, yellow 12%, orange 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>85%</span> <span>12%</span> </div> </div>
2	D	243	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 84%, yellow 13%, orange 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>84%</span> <span>13%</span> </div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cathepsin D light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	97	Total	C	N	O	S	0	0	0
			753	481	120	147	5			
1	C	103	Total	C	N	O	S	0	0	0
			792	505	126	156	5			

- Molecule 2 is a protein called Cathepsin D heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	240	Total	C	N	O	S	0	0	0
			1840	1182	301	346	11			
2	D	237	Total	C	N	O	S	0	0	0
			1817	1168	296	342	11			

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is 2-(3,4-DIMETHOXYPHENYL)-N-[N-(4-METHYLBENZYL)CARBAMIMID OYL]ACETAMIDE (three-letter code: 2S4) (formula: C<sub>19</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			25	19	3	3		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	8	Total	O	0	0
			8	8		
6	B	6	Total	O	0	0
			6	6		
6	C	5	Total	O	0	0
			5	5		
6	D	19	Total	O	0	0
			19	19		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Cathepsin D light chain





- Molecule 1: Cathepsin D light chain



- Molecule 2: Cathepsin D heavy chain



- Molecule 2: Cathepsin D heavy chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	230.34Å 42.45Å 73.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.26 – 2.90 41.75 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.26-2.90) 98.9 (41.75-2.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 2.90Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, $R_{free}$	0.177 , 0.233 0.191 , 0.252	Depositor DCC
$R_{free}$ test set	678 reflections (4.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	54.0	Xtriage
Anisotropy	0.466	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 64.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5307	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2S4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/776	0.68	0/1058
1	C	0.54	0/816	0.77	1/1114 (0.1%)
2	B	0.48	0/1879	0.72	0/2546
2	D	0.51	0/1855	0.75	0/2513
All	All	0.50	0/5326	0.73	1/7231 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	70	ASN	CB-CG-ND2	6.52	132.36	116.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	753	0	708	4	0
1	C	792	0	747	3	0
2	B	1840	0	1847	17	0
2	D	1817	0	1820	17	0
3	A	14	0	13	0	0
4	B	25	0	22	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	28	0	25	1	0
6	A	8	0	0	0	0
6	B	6	0	0	0	0
6	C	5	0	0	0	0
6	D	19	0	0	0	0
All	All	5307	0	5182	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:HIS:CE1	2:D:174:ASP:HB2	2.29	0.67
1:C:43:SER:HB2	2:D:117:GLU:HB3	1.79	0.63
2:D:195:LEU:HD13	2:D:343:ALA:HB2	1.81	0.61
2:D:248:GLN:HG2	2:D:253:ALA:HB3	1.83	0.59
2:B:284:LYS:H	2:B:284:LYS:HD3	1.68	0.58
2:B:248:GLN:HG2	2:B:253:ALA:HB3	1.85	0.57
1:A:77:HIS:HE1	2:D:174:ASP:HB2	1.73	0.54
2:B:195:LEU:HD13	2:B:343:ALA:HB2	1.90	0.53
2:D:171:ARG:HD3	2:D:327:GLY:HA3	1.89	0.53
2:D:256:LEU:HD12	2:D:257:ILE:H	1.74	0.53
2:D:124:ILE:HA	2:D:127:ILE:HD12	1.91	0.51
2:B:124:ILE:HA	2:B:127:ILE:HD12	1.93	0.51
2:B:171:ARG:HD2	2:B:327:GLY:HA3	1.93	0.50
2:B:231:ASP:OD2	4:B:401:2S4:N17	2.44	0.50
1:A:43:SER:HB2	2:B:117:GLU:HB3	1.94	0.49
1:C:70:ASN:OD1	1:C:87:LEU:HB2	2.12	0.48
2:B:164:ILE:HG22	2:B:334:ASP:HA	1.96	0.47
2:D:256:LEU:CD1	2:D:257:ILE:H	2.28	0.46
2:B:236:LEU:HD22	2:B:305:GLY:HA2	1.99	0.45
2:B:253:ALA:HB2	2:B:263:ILE:HG12	1.99	0.45
2:B:167:PHE:CZ	2:B:181:LEU:HD13	2.52	0.44
2:D:164:ILE:HG22	2:D:334:ASP:HA	2.00	0.44
2:D:253:ALA:HB2	2:D:263:ILE:HG12	1.99	0.44
2:D:264:PRO:HG2	2:D:267:LYS:HE2	1.99	0.44
2:B:248:GLN:HE21	2:B:248:GLN:HB2	1.66	0.43
1:C:2:PRO:HB2	2:D:182:MET:SD	2.59	0.43
2:B:222:CYS:O	2:B:222:CYS:SG	2.76	0.43
2:D:142:ILE:HG23	2:D:204:ALA:HB1	2.01	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:223:LYS:H	2:B:223:LYS:HG2	1.60	0.42
2:B:231:ASP:OD2	4:B:401:2S4:H18	2.20	0.42
5:C:401:NAG:H83	2:D:145:ASN:O	2.19	0.42
2:D:167:PHE:CZ	2:D:181:LEU:HD13	2.55	0.42
2:D:251:ILE:HB	2:D:271:LEU:HD22	2.02	0.42
2:B:232:THR:HG22	2:B:326:ILE:HD13	2.03	0.41
1:A:40:TRP:HA	2:B:116:GLY:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/103 (92%)	93 (98%)	2 (2%)	0	100	100
1	C	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
2	B	238/243 (98%)	228 (96%)	10 (4%)	0	100	100
2	D	233/243 (96%)	223 (96%)	10 (4%)	0	100	100
All	All	667/692 (96%)	644 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/91 (96%)	81 (93%)	6 (7%)	18	46
1	C	91/91 (100%)	82 (90%)	9 (10%)	9	28
2	B	199/201 (99%)	184 (92%)	15 (8%)	16	42
2	D	197/201 (98%)	185 (94%)	12 (6%)	22	53
All	All	574/584 (98%)	532 (93%)	42 (7%)	16	43

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	LYS
1	A	53	CYS
1	A	75	ASP
1	A	82	SER
1	A	83	LEU
1	A	93	SER
2	B	125	THR
2	B	153	ASN
2	B	158	LYS
2	B	171	ARG
2	B	198	LEU
2	B	223	LYS
2	B	224	GLU
2	B	242	ASP
2	B	248	GLN
2	B	256	LEU
2	B	257	ILE
2	B	269	SER
2	B	284	LYS
2	B	328	ARG
2	B	339	ARG
1	C	11	MET
1	C	48	LEU
1	C	53	CYS
1	C	69	LYS
1	C	70	ASN
1	C	76	ILE
1	C	83	LEU
1	C	84	SER
1	C	93	SER
2	D	125	THR
2	D	153	ASN
2	D	174	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	198	LEU
2	D	223	LYS
2	D	242	ASP
2	D	248	GLN
2	D	256	LEU
2	D	269	SER
2	D	292	LEU
2	D	300	THR
2	D	339	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	162	GLN
2	D	162	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	C	400	1,5	14,14,15	0.49	0	15,19,21	1.53	1 (6%)
5	NAG	C	401	5	14,14,15	0.36	0	15,19,21	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	C	400	1,5	-	0/6/23/26	0/1/1/1
5	NAG	C	401	5	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	400	NAG	C1-O5-C5	-4.87	105.45	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	401	NAG	1	0

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	400	1	14,14,15	0.34	0	15,19,21	0.86	1 (6%)
4	2S4	B	401	-	25,26,26	0.65	0	34,34,34	2.33	12 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	400	1	-	0/6/23/26	0/1/1/1
4	2S4	B	401	-	-	0/17/17/17	0/2/2/2

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	2S4	O7-C3-C1	-3.60	118.10	124.17
4	B	401	2S4	O9-C5-C6	-3.36	118.73	124.37
4	B	401	2S4	N13-C14-N17	-2.75	112.18	121.69
4	B	401	2S4	C11-C2-C4	-2.49	117.30	120.89
4	B	401	2S4	C18-C19-C22	-2.12	116.37	120.89
4	B	401	2S4	C11-C12-N13	2.04	116.98	114.34
3	A	400	NAG	C1-O5-C5	2.63	115.79	112.17
4	B	401	2S4	C19-C18-N16	2.76	119.04	112.96
4	B	401	2S4	C8-O7-C3	3.18	122.11	117.54
4	B	401	2S4	N13-C14-N16	3.22	126.35	118.08
4	B	401	2S4	O7-C3-C5	3.75	120.56	115.41
4	B	401	2S4	O9-C5-C3	5.31	122.70	115.41
4	B	401	2S4	C10-O9-C5	5.95	126.10	117.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	401	2S4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	97/103 (94%)	-0.14	1 (1%) 82 81	31, 51, 79, 93	0
1	C	103/103 (100%)	-0.24	0 100 100	30, 51, 71, 78	0
2	B	240/243 (98%)	-0.13	0 100 100	30, 61, 85, 105	0
2	D	237/243 (97%)	-0.40	0 100 100	23, 49, 72, 96	0
All	All	677/692 (97%)	-0.24	1 (0%) 95 95	23, 54, 79, 105	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	SER	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	C	400	14/15	0.80	0.23	0.63	82,90,91,92	0
5	NAG	C	401	14/15	0.87	0.21	-	85,95,98,101	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	2S4	B	401	25/25	0.94	0.23	0.83	39,45,68,71	0
3	NAG	A	400	14/15	0.82	0.20	0.58	73,82,84,84	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.