



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 09:37 AM EDT

PDB ID : 2OER  
Title : Probable Transcriptional Regulator from *Pseudomonas aeruginosa*  
Authors : Kim, Y.; Skarina, T.; Kagan, O.; Edwards, A.; Savchenko, A.; Joachimiak, A.;  
Midwest Center for Structural Genomics (MCSG)  
Deposited on : unknown  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>13%</div> <div> <div></div> <div>63%</div> <div>19%</div> <div>•</div> <div>16%</div> </div> </div>
1	B	214	<div> <div>13%</div> <div> <div></div> <div>70%</div> <div>14%</div> <div>•</div> <div>12%</div> </div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3272 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable transcriptional regulator.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	179	Total	C	N	O	S	Se	0	10	0
			1496	932	276	285	2	1			
1	B	189	Total	C	N	O	S	Se	0	11	0
			1576	977	294	302	2	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	CLONING ARTIFACT	UNP Q9I3U1
A	0	HIS	-	CLONING ARTIFACT	UNP Q9I3U1
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q9I3U1
A	199	MSE	MET	MODIFIED RESIDUE	UNP Q9I3U1
A	211	GLY	-	CLONING ARTIFACT	UNP Q9I3U1
A	212	SER	-	CLONING ARTIFACT	UNP Q9I3U1
B	-1	GLY	-	CLONING ARTIFACT	UNP Q9I3U1
B	0	HIS	-	CLONING ARTIFACT	UNP Q9I3U1
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q9I3U1
B	199	MSE	MET	MODIFIED RESIDUE	UNP Q9I3U1
B	211	GLY	-	CLONING ARTIFACT	UNP Q9I3U1
B	212	SER	-	CLONING ARTIFACT	UNP Q9I3U1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	121	Total	O	0	0
			121	121		
2	B	79	Total	O	0	0
			79	79		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Probable transcriptional regulator



#### • Molecule 1: Probable transcriptional regulator



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.26Å 46.39Å 56.98Å 100.62° 101.49° 109.84°	Depositor
Resolution (Å)	27.62 – 2.00 27.62 – 1.98	Depositor EDS
% Data completeness (in resolution range)	95.1 (27.62-2.00) 86.1 (27.62-1.98)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.2.0000	Depositor
R, $R_{free}$	0.212 , 0.293 0.258 , 0.324	Depositor DCC
$R_{free}$ test set	2468 reflections (11.12%)	DCC
Wilson B-factor (Å <sup>2</sup> )	34.0	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 52.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3272	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	2/1511 (0.1%)	0.84	3/2034 (0.1%)
1	B	0.81	0/1593	0.82	1/2146 (0.0%)
All	All	0.86	2/3104 (0.1%)	0.83	4/4180 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55	SER	CB-OG	7.06	1.51	1.42
1	A	108	VAL	CB-CG2	5.15	1.63	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ARG	NE-CZ-NH2	-6.88	116.86	120.30
1	A	167	LEU	CB-CG-CD2	-6.07	100.67	111.00
1	B	172	LEU	CB-CG-CD2	5.53	120.40	111.00
1	A	99	ARG	NE-CZ-NH1	5.46	123.03	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1496	0	1497	34	1
1	B	1576	0	1571	21	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	121	0	0	8	0
2	B	79	0	0	2	0
All	All	3272	0	3068	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94[B]:ARG:NH1	1:A:102[B]:ARG:HE	1.43	1.17
1:A:94[B]:ARG:NH1	1:A:102[B]:ARG:NE	2.14	0.94
1:A:94[B]:ARG:NH2	1:A:102[B]:ARG:HH21	1.65	0.93
1:A:32:VAL:HG23	1:A:73:LEU:HD22	1.49	0.92
1:B:56:ILE:H	1:B:56:ILE:HD13	1.32	0.91
1:A:94[B]:ARG:CZ	1:A:102[B]:ARG:HE	1.90	0.84
1:A:118[B]:ARG:NH1	2:A:290:HOH:O	2.04	0.83
1:A:168:LEU:HD11	2:A:245:HOH:O	1.77	0.83
1:B:20:SER:N	1:B:62:TYR:HH	1.77	0.82
1:B:47:ARG:O	1:B:51[B]:ARG:HG2	1.80	0.81
1:A:191[B]:ARG:NH1	2:A:333:HOH:O	2.04	0.79
1:B:56:ILE:H	1:B:56:ILE:CD1	2.03	0.72
1:A:94[B]:ARG:CZ	1:A:102[B]:ARG:HH21	2.04	0.70
1:B:35:LEU:HD13	1:B:43:PHE:CD1	2.31	0.65
1:A:160:GLU:HG3	2:A:304:HOH:O	1.97	0.64
1:B:99:ARG:HH22	1:B:151:GLU:CD	2.02	0.63
1:A:94[B]:ARG:HH11	1:A:102[B]:ARG:HE	1.43	0.59
1:A:94[B]:ARG:HH22	1:A:102[B]:ARG:HH21	1.50	0.58
1:B:56:ILE:N	1:B:56:ILE:HD13	2.12	0.56
1:A:80[B]:ARG:HG3	2:A:262:HOH:O	2.04	0.56
1:A:23:LEU:HD11	1:A:54:VAL:HG11	1.91	0.52
1:B:61[A]:GLN:HG2	1:B:62:TYR:CE2	2.44	0.52
1:A:24:VAL:HG12	1:A:28:LEU:HD22	1.92	0.52
1:A:105:LEU:O	1:A:109:ARG:HG3	2.11	0.51
1:A:80[B]:ARG:NH1	2:A:312:HOH:O	2.41	0.51
1:B:99:ARG:NH2	1:B:151:GLU:OE2	2.40	0.51
1:B:48:VAL:HG22	1:B:59:LEU:HD11	1.93	0.50
1:A:115:ALA:O	1:A:119:VAL:HG23	2.12	0.50
1:B:52:ALA:HB3	1:B:54:VAL:HG23	1.94	0.50
1:A:92:THR:HG22	1:A:92:THR:O	2.12	0.50
1:A:48:VAL:HG13	1:A:59:LEU:CD2	2.42	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:LEU:O	1:A:59:LEU:HD12	2.13	0.49
1:A:48:VAL:HG13	1:A:59:LEU:HD21	1.95	0.49
1:A:66:LYS:HD2	1:A:124:ALA:HB2	1.94	0.48
1:A:94[B]:ARG:HD3	1:A:98:GLU:HB3	1.95	0.48
1:A:81:THR:O	1:A:85:LEU:HG	2.14	0.47
1:B:138:LYS:HB3	2:B:247:HOH:O	2.14	0.47
1:B:61[A]:GLN:HG2	1:B:62:TYR:CD2	2.51	0.46
1:A:191[A]:ARG:HG3	2:A:236:HOH:O	2.15	0.46
2:A:261:HOH:O	1:B:170:THR:HG21	2.15	0.46
1:B:25:ALA:O	1:B:29:GLU:HG3	2.16	0.46
1:A:134:ALA:O	1:A:138:LYS:HG3	2.17	0.45
1:B:23:LEU:HD11	1:B:54:VAL:HG11	1.98	0.45
1:A:94[B]:ARG:CZ	1:A:102[B]:ARG:NH2	2.76	0.45
1:B:20:SER:HB2	1:B:21:SER:H	1.65	0.45
1:A:30:ALA:O	1:A:34:VAL:HG23	2.16	0.44
1:A:121:LEU:HA	1:A:121:LEU:HD23	1.88	0.42
1:A:94[B]:ARG:CZ	1:A:102[B]:ARG:NE	2.68	0.41
1:A:94[B]:ARG:NH1	1:A:102[B]:ARG:CZ	2.80	0.41
1:B:146:GLN:OE1	1:B:146:GLN:HA	2.20	0.41
1:A:135:ARG:HG2	1:A:138:LYS:HE3	2.02	0.41
1:B:20:SER:O	1:B:24:VAL:HG23	2.20	0.41
1:B:83:ARG:HG2	2:B:223:HOH:O	2.20	0.41
1:A:60:TYR:HA	1:A:60:TYR:HD1	1.69	0.41
1:B:85:LEU:HD21	1:B:141:GLY:HA2	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83[B]:ARG:NH2	1:B:50:GLU:OE2[1_666]	2.08	0.12

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/214 (85%)	180 (99%)	2 (1%)	0	100	100
1	B	196/214 (92%)	194 (99%)	2 (1%)	0	100	100
All	All	378/428 (88%)	374 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/167 (90%)	136 (90%)	15 (10%)	9	5
1	B	158/167 (95%)	134 (85%)	24 (15%)	3	1
All	All	309/334 (92%)	270 (87%)	39 (13%)	7	3

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	SER
1	A	28	LEU
1	A	50	GLU
1	A	58	SER
1	A	59	LEU
1	A	66	LYS
1	A	84	LEU
1	A	118[A]	ARG
1	A	118[B]	ARG
1	A	155[A]	GLU
1	A	155[B]	GLU
1	A	166[A]	ASP
1	A	166[B]	ASP
1	A	209[A]	GLU
1	A	209[B]	GLU
1	B	20	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	22	GLU
1	B	26	SER
1	B	35	LEU
1	B	48	VAL
1	B	50	GLU
1	B	54	VAL
1	B	56	ILE
1	B	61[A]	GLN
1	B	61[B]	GLN
1	B	114	GLU
1	B	118[A]	ARG
1	B	118[B]	ARG
1	B	127	LEU
1	B	129	ARG
1	B	130	ASP
1	B	136	GLU
1	B	160[A]	GLU
1	B	160[B]	GLU
1	B	162	SER
1	B	172	LEU
1	B	186	GLU
1	B	191[A]	ARG
1	B	191[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/214 (83%)	0.68	27 (15%) 2 3	8, 18, 33, 47	0
1	B	188/214 (87%)	0.85	28 (14%) 3 3	9, 23, 49, 64	0
All	All	366/428 (85%)	0.77	55 (15%) 3 3	8, 20, 43, 64	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	131	ALA	7.3
1	B	135	ARG	6.6
1	B	133	GLU	4.8
1	B	134	ALA	4.7
1	B	20	SER	4.2
1	A	135	ARG	4.0
1	A	134	ALA	3.8
1	A	196	LEU	3.8
1	B	136	GLU	3.7
1	B	21	SER	3.7
1	A	172	LEU	3.6
1	A	200	LEU	3.6
1	B	132	ASP	3.6
1	A	21	SER	3.5
1	A	42	ARG	3.5
1	B	200	LEU	3.4
1	A	136	GLU	3.2
1	A	137	VAL	3.2
1	B	92	THR	3.1
1	A	138	LYS	3.0
1	B	129	ARG	2.9
1	A	171	THR	2.8
1	B	41	GLN	2.8
1	A	209[A]	GLU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	183	PRO	2.7
1	A	168	LEU	2.6
1	B	210	ARG	2.6
1	B	171	THR	2.6
1	B	39	GLY	2.6
1	B	196	LEU	2.6
1	B	130	ASP	2.5
1	B	42	ARG	2.5
1	A	133	GLU	2.5
1	A	175	VAL	2.4
1	B	104	VAL	2.4
1	A	167	LEU	2.4
1	B	140	GLU	2.4
1	A	50	GLU	2.4
1	B	172	LEU	2.3
1	B	137	VAL	2.3
1	A	195	ALA	2.3
1	A	41	GLN	2.3
1	A	192	TYR	2.3
1	A	197	ALA	2.3
1	B	100	LEU	2.2
1	A	108	VAL	2.2
1	B	35	LEU	2.2
1	A	170	THR	2.1
1	A	104	VAL	2.1
1	A	128	TYR	2.1
1	A	92	THR	2.1
1	A	173	GLY	2.1
1	B	187	ALA	2.1
1	B	40	ALA	2.1
1	B	167	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.