



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 25, 2018 – 08:24 AM EST

PDB ID : 5OEN
Title : Crystal Structure of STAT2 in complex with IRF9
Authors : Rengachari, S.; Panne, D.
Deposited on : 2017-07-09
Resolution : 2.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1813 (2.94-2.90)
Clashscore	112137	2045 (2.94-2.90)
Ramachandran outliers	110173	1997 (2.94-2.90)
Sidechain outliers	110143	1999 (2.94-2.90)
RSRZ outliers	101464	1825 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	171	<div> <div>19%</div> <div>94%</div> <div>• •</div> </div>
2	B	175	<div> <div>17%</div> <div>97%</div> <div>• •</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5436 atoms, of which 2719 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon regulatory factor 9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	165	Total	C	H	N	O	S	0	0	0
			2595	837	1289	225	238	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	347	ALA	GLU	engineered mutation	UNP Q61179
A	348	ALA	GLU	engineered mutation	UNP Q61179

- Molecule 2 is a protein called Signal transducer and activator of transcription.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	171	Total	C	H	N	O	S	0	0	0
			2822	884	1430	249	253	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	242	ALA	GLN	engineered mutation	UNP Q3UDU1
B	243	ALA	LYS	engineered mutation	UNP Q3UDU1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total	O	0	0
			12	12		
3	B	7	Total	O	0	0
			7	7		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interferon regulatory factor 9



- Molecule 2: Signal transducer and activator of transcription



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	30.60Å 124.04Å 51.03Å 90.00° 92.10° 90.00°	Depositor
Resolution (Å)	32.12 – 2.92 47.16 – 2.88	Depositor EDS
% Data completeness (in resolution range)	88.3 (32.12-2.92) 87.1 (47.16-2.88)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.255 , 0.307 0.247 , 0.302	Depositor DCC
R_{free} test set	748 reflections (10.14%)	DCC
Wilson B-factor (Å ²)	38.5	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.068 for h,-k,-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	5436	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1342	0.41	0/1827
2	B	0.23	0/1412	0.34	0/1898
All	All	0.24	0/2754	0.37	0/3725

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1306	1289	1291	3	0
2	B	1392	1430	1432	1	0
3	A	12	0	0	0	0
3	B	7	0	0	1	0
All	All	2717	2719	2723	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:160:ARG:NH1	3:B:401:HOH:O	1.94	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:PRO:O	1:A:259:LEU:HB3	2.21	0.41
1:A:236:ARG:HD3	1:A:274:LEU:HD21	2.03	0.41
1:A:258:PRO:O	1:A:259:LEU:CB	2.69	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/171 (94%)	152 (94%)	9 (6%)	0	100	100
2	B	167/175 (95%)	163 (98%)	4 (2%)	0	100	100
All	All	328/346 (95%)	315 (96%)	13 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/151 (97%)	146 (100%)	0	100	100
2	B	154/158 (98%)	153 (99%)	1 (1%)	89	97
All	All	300/309 (97%)	299 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	233	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/171 (96%)	1.28	33 (20%) 1 1	14, 31, 76, 112	0
2	B	171/175 (97%)	1.15	29 (16%) 2 1	20, 37, 66, 82	0
All	All	336/346 (97%)	1.21	62 (18%) 1 1	14, 35, 68, 112	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	374	GLU	7.1
1	A	375	LYS	6.0
1	A	333	GLN	5.3
1	A	371	HIS	5.1
1	A	303	PRO	5.1
1	A	301	PRO	5.0
2	B	290	GLY	4.9
2	B	288	TYR	4.9
1	A	245	GLU	4.8
2	B	286	LEU	4.6
2	B	289	LYS	4.4
2	B	234	LEU	4.2
1	A	310	SER	4.1
2	B	287	ARG	4.1
2	B	291	ASP	4.1
1	A	313	CYS	4.0
1	A	208	LEU	4.0
1	A	207	PRO	4.0
1	A	257	ASP	3.8
1	A	264	HIS	3.8
2	B	314	ARG	3.7
2	B	189	PRO	3.6
1	A	376	ILE	3.5
1	A	255	LYS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	304	GLY	3.4
1	A	311	ASN	3.3
2	B	154	ASP	3.2
1	A	356	GLN	3.0
2	B	313	GLN	3.0
1	A	260	GLU	3.0
2	B	301	ASN	2.9
1	A	209	ASN	2.9
1	A	330	PHE	2.8
2	B	267	LYS	2.8
1	A	329	TYR	2.8
1	A	302	GLY	2.7
1	A	258	PRO	2.6
2	B	292	MET	2.6
1	A	373	LEU	2.5
1	A	309	PRO	2.5
2	B	252	GLU	2.4
2	B	223	LEU	2.4
2	B	315	SER	2.4
2	B	146	ASN	2.4
1	A	265	LEU	2.4
2	B	253	LEU	2.4
2	B	257	GLN	2.4
2	B	278	GLN	2.4
2	B	229	LEU	2.3
1	A	216	LEU	2.3
2	B	147	ARG	2.2
2	B	190	HIS	2.2
2	B	187	SER	2.2
2	B	254	GLN	2.2
1	A	206	LEU	2.1
1	A	334	GLY	2.1
2	B	248	ALA	2.1
1	A	331	GLN	2.0
2	B	272	LEU	2.0
1	A	308	LEU	2.0
1	A	306	HIS	2.0
2	B	262	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.