



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:44 am GMT

PDB ID : 3OFU  
Title : Crystal Structure of Cytochrome P450 CYP101C1  
Authors : Zhou, W.; Ma, M.; Bell, S.G.; Yang, W.; Hao, Y.; Rees, N.H.; Bartlam, M.;  
Wong, L.-L.; Rao, Z.  
Deposited on : 2010-08-16  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	<div> <div style="width: 61%; background-color: green;"></div> <div style="width: 28%; background-color: yellow;"></div> <div style="width: 10%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>61% 28% 10% .</div>
1	B	396	<div> <div style="width: 57%; background-color: green;"></div> <div style="width: 34%; background-color: yellow;"></div> <div style="width: 7%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>57% 34% 7% .</div>
1	C	396	<div> <div style="width: 58%; background-color: green;"></div> <div style="width: 35%; background-color: yellow;"></div> <div style="width: 6%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>58% 35% 6% .</div>
1	D	396	<div> <div style="width: 61%; background-color: green;"></div> <div style="width: 34%; background-color: yellow;"></div> <div style="width: 1%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>61% 34% . .</div>
1	E	396	<div> <div style="width: 50%; background-color: green;"></div> <div style="width: 38%; background-color: yellow;"></div> <div style="width: 9%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>% 50% 38% 9% . .</div>
1	F	396	<div> <div style="width: 50%; background-color: green;"></div> <div style="width: 38%; background-color: yellow;"></div> <div style="width: 9%; background-color: orange;"></div> <div style="width: 1%; background-color: red;"></div> <div style="width: 1%; background-color: grey;"></div> </div> <div>% 50% 38% 9% . .</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ID3	A	397	-	-	-	X
3	ID3	B	397	-	-	-	X
3	ID3	C	397	-	-	-	X

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 18917 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	396	Total	C	N	O	S	0	0	0
			3068	1950	552	547	19			
1	B	396	Total	C	N	O	S	0	0	0
			3068	1950	552	547	19			
1	C	396	Total	C	N	O	S	0	0	0
			3068	1950	552	547	19			
1	D	394	Total	C	N	O	S	0	0	0
			3053	1941	550	543	19			
1	E	390	Total	C	N	O	S	0	0	0
			3025	1926	543	537	19			
1	F	390	Total	C	N	O	S	0	0	0
			3025	1926	543	537	19			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	E	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	F	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is (3E)-4-(2,6,6-TRIMETHYLCYCLOHEX-1-EN-1-YL)BUT-3-EN-2-ONE (three-letter code: ID3) (formula: C<sub>13</sub>H<sub>20</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			14	13	1		
3	B	1	Total	C	O	0	0
			14	13	1		
3	C	1	Total	C	O	0	0
			14	13	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	51	Total 51	O 51	0	0
4	B	62	Total 62	O 62	0	0
4	C	64	Total 64	O 64	0	0
4	D	52	Total 52	O 52	0	0
4	E	41	Total 41	O 41	0	0
4	F	40	Total 40	O 40	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cytochrome P450




#### • Molecule 1: Cytochrome P450




#### • Molecule 1: Cytochrome P450






• Molecule 1: Cytochrome P450

Chain D: 61% 34%



• Molecule 1: Cytochrome P450


Chain E: 50% 38% 9%







● Molecule 1: Cytochrome P450



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.36Å 113.42Å 150.23Å 90.00° 89.90° 90.00°	Depositor
Resolution (Å)	50.00 – 2.80 46.21 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-2.80) 98.5 (46.21-2.80)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.80 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.193 , 0.303 0.195 , 0.303	Depositor DCC
$R_{free}$ test set	2827 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtriage
Anisotropy	0.339	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 3.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.458 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	18917	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.44 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.7644e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ID3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.74	39/3140 (1.2%)	1.38	32/4269 (0.7%)
1	B	1.70	37/3140 (1.2%)	1.30	25/4269 (0.6%)
1	C	1.73	45/3140 (1.4%)	1.37	35/4269 (0.8%)
1	D	1.66	24/3123 (0.8%)	1.28	23/4243 (0.5%)
1	E	1.61	26/3095 (0.8%)	1.31	26/4206 (0.6%)
1	F	1.64	33/3095 (1.1%)	1.30	29/4206 (0.7%)
All	All	1.68	204/18733 (1.1%)	1.32	170/25462 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
1	F	0	1
All	All	0	4

The worst 5 of 204 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	262	GLU	CG-CD	12.96	1.71	1.51
1	C	114	GLU	CG-CD	-10.01	1.36	1.51
1	A	17	PHE	CE1-CZ	9.69	1.55	1.37
1	C	319	GLU	CG-CD	9.58	1.66	1.51
1	C	203	ASP	CB-CG	9.41	1.71	1.51

The worst 5 of 170 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	10	ARG	NE-CZ-NH1	-13.45	113.58	120.30
1	A	66	ARG	NE-CZ-NH1	-11.52	114.54	120.30
1	A	157	LEU	CA-CB-CG	10.49	139.44	115.30
1	A	66	ARG	NE-CZ-NH2	10.36	125.48	120.30
1	C	10	ARG	NE-CZ-NH1	-10.07	115.27	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	172	ARG	Peptide
1	C	171	THR	Peptide
1	C	172	ARG	Peptide
1	F	197	ARG	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3068	0	3088	100	0
1	B	3068	0	3088	119	1
1	C	3068	0	3088	101	0
1	D	3053	0	3076	97	1
1	E	3025	0	3048	133	0
1	F	3025	0	3048	123	0
2	A	43	0	30	4	0
2	B	43	0	30	1	0
2	C	43	0	30	2	0
2	D	43	0	30	5	0
2	E	43	0	30	3	0
2	F	43	0	30	5	0
3	A	14	0	20	2	0
3	B	14	0	20	5	0
3	C	14	0	20	2	0
4	A	51	0	0	0	0
4	B	62	0	0	6	0
4	C	64	0	0	5	0
4	D	52	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	41	0	0	4	0
4	F	40	0	0	3	0
All	All	18917	0	18676	672	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 672 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:279:PRO:CG	1:B:279:PRO:CB	1.76	1.32
1:C:172:ARG:HH22	1:C:237:ASP:HB3	1.03	1.12
1:C:257:GLN:HE21	1:C:261:ARG:NH2	1.46	1.11
1:A:172:ARG:NH1	1:A:237:ASP:HB3	1.66	1.09
1:B:172:ARG:NH2	1:B:237:ASP:HB3	1.65	1.09

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:LYS:NZ	1:D:200:GLN:OE1[1_455]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/396 (100%)	355 (90%)	34 (9%)	5 (1%)	14	41
1	B	394/396 (100%)	365 (93%)	25 (6%)	4 (1%)	18	50
1	C	394/396 (100%)	360 (91%)	30 (8%)	4 (1%)	18	50
1	D	390/396 (98%)	351 (90%)	38 (10%)	1 (0%)	44	77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	386/396 (98%)	331 (86%)	42 (11%)	13 (3%)	4	15
1	F	386/396 (98%)	321 (83%)	52 (14%)	13 (3%)	4	15
All	All	2344/2376 (99%)	2083 (89%)	221 (9%)	40 (2%)	11	34

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	170	LEU
1	A	172	ARG
1	B	172	ARG
1	B	173	PRO
1	B	174	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/321 (100%)	287 (89%)	34 (11%)	8	23
1	B	321/321 (100%)	290 (90%)	31 (10%)	9	27
1	C	321/321 (100%)	295 (92%)	26 (8%)	14	37
1	D	319/321 (99%)	296 (93%)	23 (7%)	17	43
1	E	316/321 (98%)	279 (88%)	37 (12%)	6	19
1	F	316/321 (98%)	278 (88%)	38 (12%)	6	18
All	All	1914/1926 (99%)	1725 (90%)	189 (10%)	9	26

5 of 189 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	236	LEU
1	D	183	LYS
1	F	169	GLN
1	C	303	LEU
1	D	75	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	48	HIS
1	D	230	ASN
1	F	165	GLN
1	D	88	GLN
1	D	169	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ID3	A	397	-	14,14,14	1.85	7 (50%)	20,20,20	2.18	6 (30%)
2	HEM	A	417	1	28,50,50	2.44	10 (35%)	17,82,82	2.57	6 (35%)
3	ID3	B	397	-	14,14,14	1.75	6 (42%)	20,20,20	3.01	7 (35%)
2	HEM	B	417	1	28,50,50	2.56	12 (42%)	17,82,82	1.81	4 (23%)
3	ID3	C	397	-	14,14,14	2.10	7 (50%)	20,20,20	2.69	8 (40%)
2	HEM	C	417	1	28,50,50	2.54	9 (32%)	17,82,82	2.15	5 (29%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	D	417	1	28,50,50	2.34	9 (32%)	17,82,82	2.28	6 (35%)
2	HEM	E	417	1	28,50,50	2.59	9 (32%)	17,82,82	2.41	7 (41%)
2	HEM	F	417	1	28,50,50	2.64	11 (39%)	17,82,82	2.48	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ID3	A	397	-	-	0/5/22/22	0/1/1/1
2	HEM	A	417	1	-	0/6/54/54	0/0/8/8
3	ID3	B	397	-	-	2/5/22/22	0/1/1/1
2	HEM	B	417	1	-	0/6/54/54	0/0/8/8
3	ID3	C	397	-	-	0/5/22/22	0/1/1/1
2	HEM	C	417	1	-	0/6/54/54	0/0/8/8
2	HEM	D	417	1	-	0/6/54/54	0/0/8/8
2	HEM	E	417	1	-	0/6/54/54	0/0/8/8
2	HEM	F	417	1	-	0/6/54/54	0/0/8/8

The worst 5 of 80 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	417	HEM	C3B-C2B	-6.62	1.31	1.40
2	B	417	HEM	C3B-C2B	-6.35	1.32	1.40
2	D	417	HEM	C3B-C2B	-5.95	1.32	1.40
2	E	417	HEM	C3C-C2C	-4.99	1.33	1.40
2	E	417	HEM	C3B-C2B	-4.96	1.33	1.40

The worst 5 of 54 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	397	ID3	C18-C5-C6	-7.57	116.03	124.51
3	A	397	ID3	C1-C6-C5	-6.44	113.55	122.59
3	C	397	ID3	C1-C6-C5	-5.69	114.59	122.59
3	C	397	ID3	C18-C5-C6	-5.58	118.27	124.51
2	E	417	HEM	C1D-C2D-C3D	-5.47	103.19	107.00

There are no chirality outliers.

All (2) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	B	397	ID3	O1-C9-C8-C7
3	B	397	ID3	C7-C8-C9-C19

There are no ring outliers.

9 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	397	ID3	2	0
2	A	417	HEM	4	0
3	B	397	ID3	5	0
2	B	417	HEM	1	0
3	C	397	ID3	2	0
2	C	417	HEM	2	0
2	D	417	HEM	5	0
2	E	417	HEM	3	0
2	F	417	HEM	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	396/396 (100%)	-0.73	0	100   100	5, 17, 39, 75	0
1	B	396/396 (100%)	-0.68	0	100   100	6, 19, 44, 71	0
1	C	396/396 (100%)	-0.69	1 (0%)	93   92	5, 18, 37, 80	0
1	D	394/396 (99%)	-0.65	0	100   100	7, 19, 43, 64	0
1	E	390/396 (98%)	-0.12	2 (0%)	90   88	19, 38, 70, 78	0
1	F	390/396 (98%)	-0.15	3 (0%)	86   81	22, 38, 67, 77	0
All	All	2362/2376 (99%)	-0.50	6 (0%)	93   92	5, 23, 60, 80	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	202	GLY	3.4
1	F	203	ASP	2.7
1	E	179	VAL	2.7
1	F	213	PRO	2.3
1	E	154	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	ID3	B	397	14/14	0.85	0.27	6.31	28,44,50,52	0
3	ID3	C	397	14/14	0.76	0.60	6.04	54,59,63,65	14
3	ID3	A	397	14/14	0.91	0.18	2.49	31,41,43,43	0
2	HEM	C	417	43/43	0.97	0.14	0.72	2,10,16,21	0
2	HEM	A	417	43/43	0.98	0.14	0.53	2,7,14,17	0
2	HEM	D	417	43/43	0.98	0.13	0.34	3,12,18,27	0
2	HEM	B	417	43/43	0.98	0.13	0.30	3,10,16,22	0
2	HEM	F	417	43/43	0.97	0.16	0.08	13,22,27,32	0
2	HEM	E	417	43/43	0.97	0.15	-0.34	17,24,29,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.