



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2017 – 05:53 PM EDT

PDB ID : 3OKT  
Title : Mouse Plexin A2, extracellular domains 1-4  
Authors : Janssen, B.J.C.; Robinson, R.A.; Bell, C.H.; Siebold, C.; Jones, E.Y.  
Deposited on : unknown  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	681	<div> <div>3%</div> <div>75%</div> <div>21%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1655	-	-	-	X
5	GOL	A	2004	-	-	X	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	2005	-	-	-	X
5	GOL	A	2006	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Plexin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	656	5138	3255	876	972	35	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	GLU	-	expression tag	UNP P70207
A	33	THR	-	expression tag	UNP P70207
A	34	GLY	-	expression tag	UNP P70207
A	704	GLY	-	expression tag	UNP P70207
A	705	THR	-	expression tag	UNP P70207
A	706	LYS	-	expression tag	UNP P70207
A	707	HIS	-	expression tag	UNP P70207
A	708	HIS	-	expression tag	UNP P70207
A	709	HIS	-	expression tag	UNP P70207
A	710	HIS	-	expression tag	UNP P70207
A	711	HIS	-	expression tag	UNP P70207
A	712	HIS	-	expression tag	UNP P70207

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	182	Total	O	0	0
			182	182		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Plexin-A2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.86 Å 92.90 Å 62.90 Å 90.00° 102.87° 90.00°	Depositor
Resolution (Å)	44.91 – 2.30 44.91 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.3 (44.91-2.30) 97.3 (44.91-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.85 (at 2.29 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6 _289)	Depositor
R, $R_{free}$	0.196 , 0.251 0.186 , 0.240	Depositor DCC
$R_{free}$ test set	1939 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.1	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5417	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, NAG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/5261	0.48	0/7141

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5138	0	4990	94	0
2	A	70	0	65	7	0
3	A	2	0	0	0	0
4	A	1	0	0	0	0
5	A	24	0	32	7	0
6	A	182	0	0	0	0
All	All	5417	0	5087	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:VAL:HA	5:A:2004:GOL:H12	1.55	0.88
1:A:237:SER:HA	1:A:239:PHE:H	1.39	0.87
1:A:76:ASN:H	2:A:1076:NAG:H82	1.40	0.87
1:A:697:VAL:HG12	1:A:699:GLU:H	1.50	0.76
1:A:524:GLY:HA2	5:A:2007:GOL:H2	1.68	0.75
1:A:532:ALA:HB1	1:A:560:MET:CE	2.17	0.74
1:A:564:VAL:HG22	1:A:580:LEU:HD13	1.74	0.69
1:A:274:LEU:HD12	1:A:274:LEU:H	1.58	0.68
1:A:532:ALA:HB1	1:A:560:MET:HE3	1.75	0.68
1:A:666:VAL:HG11	1:A:698:SER:N	2.09	0.68
1:A:118:ILE:CG2	5:A:2004:GOL:H31	2.25	0.67
1:A:461:ASP:O	1:A:466:GLY:O	2.14	0.65
1:A:425:TYR:OH	1:A:456:LYS:HE2	1.97	0.64
1:A:544:GLN:HG2	1:A:545:ARG:HG2	1.79	0.64
1:A:175:SER:OG	1:A:178:GLU:HG2	1.97	0.64
1:A:237:SER:HA	1:A:239:PHE:N	2.13	0.63
1:A:76:ASN:H	2:A:1076:NAG:C8	2.12	0.60
1:A:172:ILE:HB	5:A:2004:GOL:H32	1.83	0.59
1:A:97:PRO:HG2	1:A:158:TYR:CE1	2.37	0.59
1:A:407:LEU:O	1:A:411:GLN:HG2	2.04	0.58
1:A:574:HIS:ND1	1:A:618:PRO:HD3	2.19	0.57
1:A:191:LYS:HB3	1:A:194:TYR:HB2	1.87	0.57
1:A:118:ILE:HA	5:A:2005:GOL:H32	1.85	0.57
1:A:79:ILE:HD11	1:A:82:ALA:HB2	1.85	0.57
1:A:663:LEU:HD23	1:A:698:SER:HB2	1.88	0.55
1:A:163:ASN:HD22	2:A:1163:NAG:H83	1.72	0.54
1:A:529:GLY:HA3	1:A:552:PHE:CZ	2.43	0.54
1:A:118:ILE:CG2	5:A:2004:GOL:H11	2.38	0.54
1:A:274:LEU:N	1:A:274:LEU:HD12	2.25	0.52
1:A:510:SER:HB3	1:A:513:GLN:HE21	1.74	0.52
1:A:235:LEU:HG	1:A:236:VAL:HG23	1.90	0.52
1:A:57:ARG:HG2	1:A:121:TYR:CE2	2.44	0.52
1:A:351:ASP:HA	1:A:431:ARG:HB2	1.92	0.52
1:A:660:GLN:C	1:A:661:LEU:HD12	2.30	0.52
1:A:308:ARG:HB2	1:A:343:GLN:HG2	1.92	0.51
1:A:95:TYR:HA	1:A:96:PRO:C	2.31	0.51
1:A:137:CYS:SG	1:A:159:LEU:HD11	2.52	0.50
1:A:638:ARG:HB2	1:A:645:ILE:HD13	1.94	0.50
1:A:118:ILE:HG21	5:A:2004:GOL:H11	1.94	0.50
1:A:235:LEU:HD21	1:A:263:PRO:HG2	1.94	0.50
1:A:50:PHE:CE2	1:A:503:VAL:HG23	2.47	0.50
1:A:191:LYS:HD2	1:A:191:LYS:N	2.26	0.50

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:GLN:NE2	1:A:37:GLN:HA	2.26	0.50
1:A:56:HIS:HB3	1:A:59:THR:OG1	2.12	0.50
1:A:176:GLU:HG3	1:A:177:GLY:N	2.27	0.49
1:A:112:ASN:HB2	1:A:132:LEU:HD22	1.95	0.49
1:A:555:SER:HB3	1:A:558:GLN:OE1	2.12	0.49
1:A:327:ASN:O	1:A:327:ASN:CG	2.52	0.48
1:A:663:LEU:CD2	1:A:698:SER:HB2	2.43	0.48
1:A:141:ARG:HD2	1:A:143:ASP:OD1	2.15	0.47
1:A:162:VAL:CG2	1:A:189:ASP:HB2	2.45	0.47
1:A:574:HIS:CE1	1:A:618:PRO:HD3	2.50	0.47
1:A:672:CYS:HB3	1:A:681:CYS:SG	2.53	0.47
1:A:630:TRP:HB3	1:A:670:PHE:CE2	2.50	0.47
1:A:88:GLU:HB2	1:A:132:LEU:HD21	1.97	0.47
1:A:238:HIS:CD2	1:A:238:HIS:N	2.82	0.47
1:A:564:VAL:HG23	1:A:649:THR:HG21	1.97	0.46
1:A:73:LEU:HD23	1:A:79:ILE:HA	1.98	0.46
1:A:192:GLN:HG3	1:A:228:ILE:O	2.16	0.46
1:A:594:CYS:O	1:A:601:GLU:HA	2.15	0.46
1:A:695:ILE:HG23	1:A:700:ASP:OD1	2.16	0.45
1:A:571:VAL:HG22	1:A:654:TYR:O	2.16	0.45
1:A:381:LEU:HD21	1:A:410:ASN:HA	1.99	0.44
1:A:661:LEU:HD12	1:A:661:LEU:N	2.33	0.44
1:A:328:ILE:HB	1:A:332:GLU:OE1	2.17	0.44
1:A:257:THR:O	1:A:278:SER:HA	2.18	0.43
1:A:509:GLU:HG3	1:A:511:CYS:SG	2.58	0.43
1:A:359:PRO:HG2	1:A:362:ALA:CB	2.48	0.43
1:A:533:LEU:HB3	1:A:642:THR:HG21	2.01	0.43
1:A:565:HIS:HA	1:A:566:PRO:HA	1.86	0.43
1:A:359:PRO:HG2	1:A:362:ALA:HB3	2.01	0.43
1:A:695:ILE:HD13	1:A:700:ASP:O	2.18	0.43
1:A:175:SER:HG	1:A:178:GLU:HG2	1.83	0.43
1:A:315:LEU:HD11	1:A:333:ASP:HB3	2.01	0.42
1:A:525:ASP:HA	1:A:526:PRO:HD3	1.87	0.42
1:A:539:ARG:HB2	1:A:542:LYS:HD3	2.01	0.42
1:A:417:THR:HA	1:A:418:PRO:HD3	1.72	0.42
1:A:432:LEU:HD22	1:A:448:VAL:HG11	2.02	0.42
2:A:1655:NAG:H2	2:A:1655:NAG:H82	1.86	0.42
1:A:117:LEU:HD11	1:A:126:LEU:HD21	2.02	0.42
1:A:279:ARG:HD3	1:A:295:SER:OG	2.20	0.42
1:A:75:GLY:N	2:A:1076:NAG:H81	2.34	0.42
1:A:46:ARG:HG2	1:A:46:ARG:O	2.19	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:VAL:HG22	1:A:150:GLU:N	2.34	0.41
1:A:673:HIS:CD2	1:A:685:PRO:HG3	2.55	0.41
1:A:482:ILE:HG23	1:A:497:VAL:HG13	2.02	0.41
1:A:76:ASN:N	2:A:1076:NAG:C8	2.83	0.41
1:A:598:ASN:ND2	2:A:1598:NAG:H83	2.36	0.41
1:A:619:LYS:HD3	1:A:619:LYS:HA	1.87	0.41
1:A:311:GLN:O	1:A:484:ARG:NH1	2.52	0.40
1:A:533:LEU:HD22	1:A:639:SER:HB2	2.03	0.40
1:A:113:VAL:O	1:A:130:GLY:HA2	2.21	0.40
1:A:218:HIS:HA	1:A:223:SER:OG	2.21	0.40
1:A:411:GLN:HA	1:A:412:PRO:C	2.42	0.40
1:A:582:VAL:CG1	1:A:585:ALA:HB2	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	650/681 (95%)	635 (98%)	13 (2%)	2 (0%)	44 55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	SER
1	A	175	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	578/597 (97%)	562 (97%)	16 (3%)	49 65

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	ILE
1	A	192	GLN
1	A	200	SER
1	A	230	SER
1	A	237	SER
1	A	261	GLU
1	A	274	LEU
1	A	298	PHE
1	A	438	TYR
1	A	473	VAL
1	A	474	SER
1	A	509	GLU
1	A	542	LYS
1	A	580	LEU
1	A	626	LEU
1	A	680	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	101	GLN
1	A	238	HIS
1	A	513	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	1076	1	14,14,15	0.38	0	15,19,21	1.89	2 (13%)
2	NAG	A	1163	1	14,14,15	0.42	0	15,19,21	1.26	2 (13%)
2	NAG	A	1327	1	14,14,15	0.45	0	15,19,21	1.76	2 (13%)
2	NAG	A	1598	1	14,14,15	0.50	0	15,19,21	0.99	1 (6%)
2	NAG	A	1655	1	14,14,15	0.40	0	15,19,21	1.43	4 (26%)
5	GOL	A	2004	-	5,5,5	0.36	0	5,5,5	0.28	0
5	GOL	A	2005	-	5,5,5	0.37	0	5,5,5	0.31	0
5	GOL	A	2006	-	5,5,5	0.35	0	5,5,5	0.21	0
5	GOL	A	2007	-	5,5,5	0.28	0	5,5,5	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1076	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1163	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1327	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1598	1	-	0/6/23/26	0/1/1/1
2	NAG	A	1655	1	-	0/6/23/26	0/1/1/1
5	GOL	A	2004	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2005	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2006	-	-	0/4/4/4	0/0/0/0
5	GOL	A	2007	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1655	NAG	C4-C3-C2	-2.45	107.43	111.02
2	A	1163	NAG	O5-C1-C2	-2.34	108.22	111.47
2	A	1655	NAG	O5-C1-C2	-2.24	108.36	111.47
2	A	1655	NAG	C6-C5-C4	-2.03	108.24	113.00
2	A	1076	NAG	O5-C1-C2	2.41	114.83	111.47
2	A	1327	NAG	O5-C1-C2	3.08	115.76	111.47
2	A	1598	NAG	C1-O5-C5	3.12	116.47	112.17
2	A	1655	NAG	C1-O5-C5	3.14	116.49	112.17
2	A	1163	NAG	C1-O5-C5	3.44	116.90	112.17
2	A	1327	NAG	C1-O5-C5	5.49	119.74	112.17
2	A	1076	NAG	C1-O5-C5	6.24	120.76	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1076	NAG	4	0
2	A	1163	NAG	1	0
2	A	1598	NAG	1	0
2	A	1655	NAG	1	0
5	A	2004	GOL	5	0
5	A	2005	GOL	1	0
5	A	2007	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	656/681 (96%)	0.32	23 (3%)	44 51	19, 37, 69, 105	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	36	PRO	4.8
1	A	176	GLU	3.8
1	A	631	PHE	3.3
1	A	629	ASP	3.2
1	A	376	HIS	3.0
1	A	465	HIS	2.9
1	A	388	GLY	2.9
1	A	661	LEU	2.8
1	A	235	LEU	2.8
1	A	598	ASN	2.7
1	A	93	ALA	2.6
1	A	236	VAL	2.6
1	A	399	PRO	2.6
1	A	327	ASN	2.4
1	A	401	ASP	2.3
1	A	623	VAL	2.3
1	A	263	PRO	2.2
1	A	625	PRO	2.2
1	A	571	VAL	2.2
1	A	104	SER	2.1
1	A	391	VAL	2.1
1	A	400	ILE	2.1
1	A	312	ALA	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GOL	A	2005	6/6	0.73	0.31	8.80	46,52,53,54	0
5	GOL	A	2004	6/6	0.87	0.26	4.26	37,46,48,55	0
2	NAG	A	1655	14/15	0.89	0.35	3.28	81,86,89,90	0
5	GOL	A	2006	6/6	0.68	0.26	2.59	74,76,80,81	0
5	GOL	A	2007	6/6	0.90	0.17	1.44	54,56,58,60	0
2	NAG	A	1163	14/15	0.85	0.15	0.20	56,59,78,84	0
3	CL	A	2002	1/1	0.93	0.11	-0.49	66,66,66,66	0
3	CL	A	2001	1/1	0.97	0.13	-1.13	41,41,41,41	0
4	NA	A	2003	1/1	0.81	0.41	-	52,52,52,52	0
2	NAG	A	1327	14/15	0.70	0.45	-	107,119,122,123	0
2	NAG	A	1076	14/15	0.89	0.17	-	58,68,72,73	0
2	NAG	A	1598	14/15	0.67	0.54	-	112,118,123,123	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.