



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2018 – 10:52 PM EST

PDB ID : 1OLZ
Title : The ligand-binding face of the semaphorins revealed by the high resolution crystal structure of SEMA4D
Authors : Love, C.A.; Harlos, K.; Mavaddat, N.; Davis, S.J.; Stuart, D.I.; Jones, E.Y.; Esnouf, R.M.
Deposited on : 2003-08-19
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	663	<div> <div>5%</div> <div> <div></div> <div>57%</div> <div>28%</div> <div>8%</div> <div>6%</div> </div> </div>
1	B	663	<div> <div>6%</div> <div> <div></div> <div>60%</div> <div>26%</div> <div>8%</div> <div>6%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SEMAPHORIN 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	622	Total	C	N	O	S	0	0	1
			4914	3128	850	913	23			
1	B	622	Total	C	N	O	S	0	0	1
			4914	3128	850	913	23			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	398	Total	O	0	0
			398	398		
2	B	443	Total	O	0	0
			443	443		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SEMAPHORIN 4D



• Molecule 1: SEMAPHORIN 4D



SER	K574	M471	P346	P225
VAL	A575	S472	D351	R226
VAL	E576	S473	G352	I229
GLN	S577	G473	S353	P230
THR	P578	V474	A354	R231
GLU	K579	V475	R355	R234
GLY	Y580	L479	A356	R257
SER	G581	K484	A357	Q240
ARG	L582	E489	N358	R263
ILE	M583	R495	T369	D264
ALA	G584	P503	L370	S265
THR	K585	P504	G371	L272
LYS	K586	L511	L370	L278
VAL	N587	H512	D387	R279
LEU	L588	Q513	N388	S280
ALA	L590	M525	R389	P281
SER	F591	S526	R391	G282
THR	N592	GE27	L392	L283
THR	L593	C532	T393	K284
LYS	L594	K394	K394	V285
HIS	S594	P533	K395	L291
HIS	E595	D534	D396	N297
HIS	G596	K535	V397	L301
HIS	D597	S536	N398	N308
HIS	S598	GE27	Y399	L309
HIS	G599	K537	Y403	K320
	Y600	G538	V403	Y321
	Y601	R541	Q408	N322
	L604	Q542	A409	Q323
	E607	H543	L410	S324
	R608	F544	L410	T325
	V609	F545	V414	T326
	K610	T550	M418	V327
	N611	A551	F419	E328
	K612	E552	E434	Q329
	T613	C555	H435	S330
	V617	S556	F446	H331
	V618	Q557	Q447	T332
	A619	A562	D448	K333
	K620	R563	Q453	W334
	H621	V564	F565	V335
	H622	F566	K567	R336
	V623	W566	F568	P342
	E624	K569	K460	K343
	L623	N570	K461	P344
	V625	G571	N463	R345
	V626	L573	R464	
	V627			
	V628			
	PRO			
	LYS			
	PRO			
	VAL			
	ALA			
	PRO			
	THR			
	LEU			

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	73.32Å 76.76Å 89.41Å 77.41° 73.35° 63.57°	Depositor
Resolution (Å)	20.00 – 2.00 19.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.7 (20.00-2.00) 93.6 (19.51-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.38 (at 2.01Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.206 , 0.270 0.206 , 0.267	Depositor DCC
R_{free} test set	5440 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 72.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10669	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/5034	0.64	1/6825 (0.0%)
1	B	0.39	0/5034	0.68	2/6825 (0.0%)
All	All	0.39	0/10068	0.66	3/13650 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	199	SER	C-N-CD	-7.49	104.11	120.60
1	B	199	SER	C-N-CA	6.12	147.71	122.00
1	A	457	LEU	CA-CB-CG	5.20	127.25	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4914	0	4841	223	0
1	B	4914	0	4841	198	0
2	A	398	0	0	24	0
2	B	443	0	0	39	0
All	All	10669	0	9682	414	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 414 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:ARG:HG3	1:A:495:ARG:HH11	1.06	1.12
1:A:608:ARG:HD2	1:A:608:ARG:H	1.13	1.12
1:B:200:PRO:HD3	1:B:205:GLY:HA2	1.35	1.08
1:B:343:LYS:HB3	1:B:344:PRO:HD3	1.33	1.07
1:B:199:SER:HB3	1:B:200:PRO:C	1.82	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	618/663 (93%)	565 (91%)	35 (6%)	18 (3%)	5	1
1	B	618/663 (93%)	567 (92%)	38 (6%)	13 (2%)	8	3
All	All	1236/1326 (93%)	1132 (92%)	73 (6%)	31 (2%)	6	2

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	434	GLU
1	A	511	LEU
1	A	516	SER
1	A	610	LYS
1	B	199	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	545/582 (94%)	471 (86%)	74 (14%)	4	2
1	B	545/582 (94%)	465 (85%)	80 (15%)	3	1
All	All	1090/1164 (94%)	936 (86%)	154 (14%)	4	2

5 of 154 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	606	GLU
1	B	82	SER
1	B	592	ASN
1	A	608	ARG
1	B	38	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	308	ASN
1	B	338	ASN
1	B	463	ASN
1	B	13	HIS
1	B	471	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	622/663 (93%)	-0.02	32 (5%) 29 29	13, 36, 81, 100	0
1	B	622/663 (93%)	-0.07	42 (6%) 18 18	8, 29, 85, 100	0
All	All	1244/1326 (93%)	-0.05	74 (5%) 23 23	8, 33, 84, 100	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	200	PRO	8.4
1	A	327	VAL	7.8
1	B	577	SER	7.8
1	A	628	VAL	7.4
1	A	199	SER	7.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.