



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 15, 2017 – 05:46 AM EDT

PDB ID : 4OM0  
Title : Crystal structure of antibody VRC07-G54Y in complex with clade A/E  
93TH057 HIV-1 gp120 core  
Authors : Kwon, Y.D.; Kwong, P.D.  
Deposited on : unknown  
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5609 (2.30-2.26)
Clashscore	112137	6364 (2.30-2.26)
Ramachandran outliers	110173	6281 (2.30-2.26)
Sidechain outliers	110143	6281 (2.30-2.26)
RSRZ outliers	101464	5639 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	353	<div> <div>5%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
2	H	228	<div> <div>4%</div> <div>89%</div> <div>10%</div> <div>.</div> </div>
3	L	210	<div> <div>91%</div> <div>8%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	G	504	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6440 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein gp160.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	342	Total	C	N	O	S	0	0	0
			2677	1680	466	509	22			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	124	GLY	-	LINKER	UNP Q0ED31
G	198	GLY	-	LINKER	UNP Q0ED31
G	318	GLY	-	LINKER	UNP Q0ED31
G	319	GLY	-	LINKER	UNP Q0ED31
G	320	SER	-	LINKER	UNP Q0ED31
G	321	GLY	-	LINKER	UNP Q0ED31
G	322	SER	-	LINKER	UNP Q0ED31
G	323	GLY	-	LINKER	UNP Q0ED31

- Molecule 2 is a protein called Antigen binding fragment of heavy chain: Antibody VRC01.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	228	Total	C	N	O	S	0	0	0
			1755	1104	304	334	13			

- Molecule 3 is a protein called Antigen binding fragment of light chain: Antibody VRC01.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	208	Total	C	N	O	S	0	0	0
			1607	1007	273	322	5			

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		
4	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	G	70	Total	O	0	0
			70	70		
5	H	99	Total	O	0	0
			99	99		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	92	Total	O	0	0
			92	92		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Envelope glycoprotein gp160



- Molecule 2: Antigen binding fragment of heavy chain: Antibody VRC01



- Molecule 3: Antigen binding fragment of light chain: Antibody VRC01



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.90Å 75.53Å 198.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.10 – 2.29 37.10 – 2.29	Depositor EDS
% Data completeness (in resolution range)	90.5 (37.10-2.29) 84.3 (37.10-2.29)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 2.29Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, $R_{free}$	0.191 , 0.247 0.193 , 0.242	Depositor DCC
$R_{free}$ test set	1867 reflections (4.66%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.8	Xtriage
Anisotropy	0.453	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 38.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6440	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	G	0.25	0/2733	0.42	0/3709
2	H	0.29	0/1802	0.48	0/2447
3	L	0.26	0/1644	0.43	0/2232
All	All	0.27	0/6179	0.44	0/8388

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2677	0	2611	19	0
2	H	1755	0	1713	20	0
3	L	1607	0	1542	12	0
4	G	140	0	130	2	0
5	G	70	0	0	3	0
5	H	99	0	0	10	0
5	L	92	0	0	3	0
All	All	6440	0	5996	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:1:GLN:N	5:H:385:HOH:O	2.15	0.80
3:L:47:VAL:HG12	3:L:48:ILE:HG12	1.67	0.77
2:H:23:ARG:NH2	5:H:382:HOH:O	2.17	0.76
1:G:461:ASN:HD21	2:H:61:ARG:HH21	1.34	0.74
1:G:123:THR:HG21	1:G:430:THR:HG22	1.72	0.71
3:L:107:ASP:OD1	5:L:307:HOH:O	2.09	0.70
1:G:336:THR:OG1	1:G:411:ASN:OD1	2.11	0.67
1:G:65:VAL:HG21	1:G:208:ILE:HD12	1.78	0.66
3:L:110:ARG:HD2	3:L:173:SER:HB2	1.80	0.64
2:H:143:LYS:NZ	2:H:171:GLN:OE1	2.32	0.63
2:H:20:ILE:HD11	2:H:80:LEU:HD23	1.83	0.60
1:G:461:ASN:ND2	2:H:61:ARG:HH21	1.99	0.59
3:L:199:THR:HG22	3:L:206:PRO:HB3	1.83	0.59
3:L:30:SER:HB2	3:L:50:SER:HA	1.87	0.56
1:G:88:ASN:ND2	5:G:660:HOH:O	2.41	0.53
3:L:110:ARG:NH1	3:L:111:THR:O	2.41	0.53
2:H:193:THR:HG23	2:H:210:LYS:HE3	1.92	0.51
2:H:96:LYS:HB2	2:H:101:GLU:HB2	1.93	0.51
1:G:461:ASN:ND2	2:H:61:ARG:HD2	2.25	0.51
1:G:72:HIS:ND1	5:G:608:HOH:O	2.33	0.50
1:G:389:GLN:HG2	4:G:509:NAG:H81	1.93	0.50
2:H:94:ARG:NH2	5:H:337:HOH:O	2.37	0.49
2:H:76:GLU:OE2	5:H:345:HOH:O	2.20	0.49
1:G:267:GLU:HG2	5:G:648:HOH:O	2.13	0.48
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.94	0.48
2:H:192:GLN:OE1	5:H:370:HOH:O	2.20	0.47
3:L:157:GLN:HG3	3:L:160:ASN:HD21	1.80	0.47
3:L:191:HIS:O	3:L:213:ARG:NH1	2.47	0.46
3:L:157:GLN:NE2	5:L:385:HOH:O	2.49	0.46
2:H:148:GLU:OE2	5:H:380:HOH:O	2.21	0.46
1:G:369:LEU:HD12	1:G:421:LYS:HE3	1.99	0.45
2:H:154:TRP:CH2	2:H:196:CYS:HB3	2.53	0.43
1:G:54:CYS:SG	1:G:215:ILE:HG23	2.58	0.43
1:G:44:VAL:HG13	1:G:492:GLU:HG2	2.00	0.43
2:H:216:CYS:SG	5:H:376:HOH:O	2.62	0.42
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.55	0.42
3:L:199:THR:HG23	5:L:311:HOH:O	2.18	0.42
2:H:59:TYR:O	5:H:302:HOH:O	2.22	0.42
1:G:378:CYS:HB3	1:G:383:PHE:CE1	2.55	0.41
1:G:446:VAL:HG21	4:G:506:NAG:H82	2.02	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:12:SER:HB3	3:L:107:ASP:HB2	2.02	0.41
1:G:298:ARG:HB3	1:G:443:ILE:HB	2.02	0.41
1:G:416:LEU:HA	1:G:417:PRO:HD3	1.93	0.41
1:G:343:LYS:NZ	1:G:347:GLU:OE2	2.54	0.40
2:H:216:CYS:HB3	5:H:367:HOH:O	2.21	0.40
2:H:126:PRO:HB2	2:H:127:SER:H	1.61	0.40
2:H:192:GLN:HB2	5:H:370:HOH:O	2.21	0.40
3:L:189:GLU:HA	3:L:213:ARG:NE	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	336/353 (95%)	322 (96%)	14 (4%)	0	100	100
2	H	226/228 (99%)	220 (97%)	4 (2%)	2 (1%)	20	21
3	L	206/210 (98%)	203 (98%)	3 (2%)	0	100	100
All	All	768/791 (97%)	745 (97%)	21 (3%)	2 (0%)	44	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	126	PRO
2	H	133	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	305/311 (98%)	303 (99%)	2 (1%)	87	93
2	H	196/196 (100%)	194 (99%)	2 (1%)	80	89
3	L	179/182 (98%)	179 (100%)	0	100	100
All	All	680/689 (99%)	676 (99%)	4 (1%)	89	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	276	ASN
1	G	430	THR
2	H	71	ARG
2	H	94	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	461	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	G	501	1	14,14,15	0.31	0	15,19,21	0.37	0
4	NAG	G	502	1	14,14,15	1.10	2 (14%)	15,19,21	1.59	1 (6%)
4	NAG	G	503	1	14,14,15	0.24	0	15,19,21	0.71	1 (6%)
4	NAG	G	504	1	14,14,15	0.41	0	15,19,21	0.79	1 (6%)
4	NAG	G	505	1	14,14,15	0.48	0	15,19,21	0.51	0
4	NAG	G	506	1	14,14,15	0.26	0	15,19,21	0.65	1 (6%)
4	NAG	G	507	1	14,14,15	0.42	0	15,19,21	0.59	0
4	NAG	G	508	1	14,14,15	0.64	1 (7%)	15,19,21	0.60	0
4	NAG	G	509	1	14,14,15	0.50	0	15,19,21	0.62	0
4	NAG	G	510	1	14,14,15	0.39	0	15,19,21	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	501	1	-	0/6/23/26	0/1/1/1
4	NAG	G	502	1	-	0/6/23/26	0/1/1/1
4	NAG	G	503	1	-	0/6/23/26	0/1/1/1
4	NAG	G	504	1	-	0/6/23/26	0/1/1/1
4	NAG	G	505	1	-	0/6/23/26	0/1/1/1
4	NAG	G	506	1	-	0/6/23/26	0/1/1/1
4	NAG	G	507	1	-	0/6/23/26	0/1/1/1
4	NAG	G	508	1	-	0/6/23/26	0/1/1/1
4	NAG	G	509	1	-	0/6/23/26	0/1/1/1
4	NAG	G	510	1	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	508	NAG	O5-C1	-2.20	1.40	1.43
4	G	502	NAG	C1-C2	2.31	1.55	1.52
4	G	502	NAG	O5-C1	3.32	1.49	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	506	NAG	C1-O5-C5	2.07	115.01	112.17
4	G	503	NAG	C1-O5-C5	2.23	115.24	112.17
4	G	504	NAG	C1-O5-C5	2.74	115.95	112.17
4	G	502	NAG	C1-O5-C5	5.84	120.22	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	506	NAG	1	0
4	G	509	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	G	342/353 (96%)	0.27	19 (5%)	25 30	34, 58, 92, 117	0
2	H	228/228 (100%)	-0.01	10 (4%)	35 41	29, 43, 84, 162	0
3	L	208/210 (99%)	-0.38	1 (0%)	90 93	26, 42, 70, 144	0
All	All	778/791 (98%)	0.01	30 (3%)	40 46	26, 49, 87, 162	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	132	SER	14.3
2	H	133	GLY	11.7
2	H	128	SER	8.2
2	H	131	THR	7.1
2	H	129	LYS	7.0
1	G	410	CYS	6.5
2	H	130	SER	6.5
1	G	124	GLY	6.3
2	H	216	CYS	6.0
1	G	44	VAL	5.7
1	G	462	ASN	5.5
1	G	198	GLY	4.9
1	G	123	THR	4.5
1	G	301	ASN	4.5
1	G	461	ASN	4.2
1	G	492	GLU	3.9
3	L	216	CYS	3.8
2	H	215	SER	3.7
1	G	411	ASN	3.4
1	G	199	SER	3.3
1	G	200	VAL	3.2
2	H	74	TYR	3.2
1	G	441	GLY	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	G	328	LYS	2.9
1	G	201	ILE	2.8
1	G	45	TRP	2.6
1	G	114	GLN	2.5
1	G	326	ILE	2.3
2	H	214	LYS	2.2
1	G	87	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	G	504	14/15	0.80	0.22	2.59	71,89,106,107	0
4	NAG	G	509	14/15	0.90	0.30	0.93	84,99,116,123	0
4	NAG	G	506	14/15	0.91	0.17	0.50	47,64,79,82	0
4	NAG	G	501	14/15	0.93	0.13	-0.26	44,57,69,69	0
4	NAG	G	508	14/15	0.89	0.15	-0.35	66,84,91,100	0
4	NAG	G	503	14/15	0.92	0.12	-0.40	43,53,59,61	0
4	NAG	G	505	14/15	0.92	0.12	-0.63	53,66,76,85	0
4	NAG	G	510	14/15	0.84	0.32	-	94,110,122,122	0
4	NAG	G	507	14/15	0.90	0.29	-	76,95,100,101	0
4	NAG	G	502	14/15	0.63	0.51	-	115,145,162,163	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.