



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 14, 2017 – 08:09 AM EST

PDB ID : 5ON4  
Title : Crystal structure of NikA in complex with Fe-L1 (N-(2-hydroxybenzyl)-N'-(2-thiomethylbenzyl)-N,N'-ethylenediamine diacetic acid)  
Authors : Cavazza, C.; Menage, S.  
Deposited on : unknown  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	<div> <div>2%</div> <div>90%</div> <div>9%</div> <div>.</div> </div>
1	B	502	<div> <div>11%</div> <div>78%</div> <div>20%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	9YK	A	602	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nickel-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	0	9	0
			3993	2560	672	751	10			
1	B	498	Total	C	N	O	S	0	11	0
			4014	2573	676	755	10			

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Fe	0	0
			1	1		
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is 2-[2-[[2,3-bis(oxidanyl)phenyl]methyl-(2-hydroxy-2-oxoethyl)amino]ethyl-[(2-methylsulfonylphenyl)methyl]amino]ethanoic acid (three-letter code: 9YK) (formula: C<sub>21</sub>H<sub>26</sub>N<sub>2</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			29	21	2	5	1		
3	B	1	Total	C	N	O	S	0	0
			29	21	2	5	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	53	Total	O	0	0
			53	53		
5	B	37	Total	O	0	0
			37	37		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Nickel-binding periplasmic protein



#### • Molecule 1: Nickel-binding periplasmic protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.69 Å 94.12 Å 129.54 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.44 – 2.30 45.44 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.0 (45.44-2.30) 97.0 (45.44-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 2.29 Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.221 , 0.267 0.221 , 0.267	Depositor DCC
$R_{free}$ test set	2317 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.5	Xtriage
Anisotropy	0.461	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8181	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FE, 9YK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.22	0/4118	0.41	0/5609
1	B	0.27	0/4143	0.49	0/5645
All	All	0.24	0/8261	0.45	0/11254

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	57	SER	Peptide
1	B	58	GLU	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3993	0	3953	27	0
1	B	4014	0	3974	82	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	29	0	0	2	0
3	B	29	0	0	0	0
4	A	12	0	15	1	0
4	B	12	0	16	0	0
5	A	53	0	0	4	0
5	B	37	0	0	3	0
All	All	8181	0	7958	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:LYS:HD3	1:B:68:ARG:HD3	1.53	0.90
1:B:58:GLU:HG2	1:B:59:ASP:H	1.37	0.88
1:A:369:ALA:O	5:A:701:HOH:O	1.91	0.87
1:B:57:SER:N	1:B:58:GLU:OE1	2.07	0.87
1:B:57:SER:HB2	1:B:58:GLU:HB3	1.57	0.86
1:B:58:GLU:CG	1:B:59:ASP:H	1.88	0.86
1:B:370:ASP:OD2	5:B:701:HOH:O	1.99	0.79
1:B:82:GLU:HA	1:B:85:ALA:H	1.50	0.77
1:B:52:LYS:HG2	1:B:68:ARG:HG2	1.71	0.72
1:B:58:GLU:HG2	1:B:59:ASP:N	2.06	0.71
1:B:58:GLU:OE2	1:B:60:GLY:N	2.19	0.71
1:B:114:SER:H	1:B:117:GLU:HB2	1.55	0.70
1:A:433:LYS:NZ	1:A:437:GLU:OE2	2.25	0.70
1:B:241:GLN:HB2	1:B:474:VAL:HB	1.72	0.70
1:B:110:VAL:HG22	1:B:120:ILE:HG12	1.72	0.70
1:A:221:GLU:OE2	1:A:396:ARG:NH2	2.26	0.69
1:B:104:ALA:C	1:B:105:ASN:HD22	1.95	0.69
1:B:83:ALA:O	1:B:87:ASN:ND2	2.21	0.68
1:B:105:ASN:HD22	1:B:105:ASN:N	1.91	0.68
1:B:110:VAL:C	1:B:111:LYS:HD3	2.15	0.67
1:B:252:ALA:HB3	1:B:393:ILE:HG22	1.76	0.67
1:B:61:LYS:HE2	1:B:123:LYS:C	2.16	0.66
1:B:40:TYR:HA	1:B:46:VAL:HG12	1.78	0.65
1:B:58:GLU:CG	1:B:59:ASP:N	2.59	0.65

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:ASP:HB3	1:B:284:TYR:CZ	2.31	0.64
1:A:100:TRP:HA	1:A:413:VAL:HG21	1.83	0.61
1:B:19:PRO:HG3	1:B:142:ILE:HB	1.84	0.60
1:B:276:LYS:HG2	1:B:289:VAL:HG21	1.84	0.60
1:A:241:GLN:HB2	1:A:474:VAL:HB	1.84	0.60
1:B:75:ASN:ND2	1:B:77:GLU:OE1	2.34	0.58
1:A:227:ASP:HB3	1:A:284:TYR:CZ	2.40	0.57
1:B:251:LEU:HD11	1:B:360:ILE:HD13	1.87	0.56
1:A:132:GLU:HA	1:A:135:LEU:HG	1.88	0.56
1:B:81:ALA:HB3	1:B:115:LYS:HD2	1.87	0.55
1:B:105:ASN:ND2	1:B:105:ASN:N	2.54	0.55
1:B:128:PRO:O	1:B:132:GLU:HG3	2.07	0.55
1:B:58:GLU:HG2	1:B:60:GLY:N	2.22	0.55
1:B:82:GLU:H	1:B:115:LYS:NZ	2.04	0.55
1:B:109:ASP:OD1	1:B:121:THR:HB	2.06	0.54
1:B:446:GLN:HA	1:B:449:ALA:HB3	1.90	0.54
1:B:111:LYS:HD3	1:B:111:LYS:N	2.21	0.54
1:B:80:ASP:HB2	1:B:115:LYS:HG3	1.90	0.54
4:A:604:GOL:O1	5:A:703:HOH:O	2.19	0.52
1:B:414:PRO:HA	1:B:419:PHE:CD1	2.44	0.52
1:B:61:LYS:HE2	1:B:124:SER:N	2.25	0.51
1:B:182:GLU:HA	1:B:188:LYS:HE2	1.90	0.51
1:A:61:LYS:HE2	1:A:124:SER:HA	1.94	0.50
1:B:7[A]:THR:HG22	1:B:214:ILE:HG22	1.93	0.50
1:B:82:GLU:HG2	1:B:115:LYS:CE	2.41	0.50
1:B:106:GLN:HE22	1:B:124:SER:HB2	1.77	0.50
1:B:82:GLU:HG2	1:B:115:LYS:HE2	1.94	0.50
1:B:57:SER:CB	1:B:58:GLU:HB3	2.37	0.49
1:B:38:VAL:HG22	1:B:46:VAL:HB	1.94	0.49
1:B:109:ASP:HB2	1:B:111:LYS:HE2	1.94	0.49
1:B:58:GLU:CD	1:B:60:GLY:H	2.10	0.48
1:B:202:PRO:HB2	1:B:225:PRO:HD3	1.94	0.48
1:B:361:GLN:HG3	1:B:371:VAL:HB	1.94	0.48
1:B:444:GLU:OE2	1:B:447:ARG:NH1	2.47	0.47
1:A:18:ASN:O	1:A:30:GLN:NE2	2.48	0.47
1:B:97:ARG:NH1	5:B:706:HOH:O	2.47	0.47
1:B:106:GLN:NE2	1:B:124:SER:HB2	2.30	0.47
1:B:74:SER:HB3	1:B:160:ILE:HG23	1.96	0.46
1:B:115:LYS:HA	1:B:115:LYS:HD2	1.57	0.46
1:B:58:GLU:CD	1:B:59:ASP:H	2.18	0.46
1:B:6:ILE:HG12	1:B:191:ILE:HG21	1.97	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:LYS:O	1:B:118:LEU:HD12	2.16	0.46
1:B:132:GLU:HA	1:B:135:LEU:HG	1.97	0.46
1:A:251:LEU:HB2	1:A:465:LEU:HB3	1.97	0.46
1:B:201:ASP:HA	1:B:202:PRO:HD2	1.72	0.46
1:B:171:LYS:HB3	1:B:174:GLN:OE1	2.15	0.46
1:B:217:LEU:HB3	1:B:223:LEU:HD21	1.98	0.46
1:B:422:GLN:HB3	1:B:428:LYS:HD3	1.97	0.45
1:B:206:ALA:O	1:B:210:GLU:HG3	2.15	0.45
1:B:40:TYR:CZ	1:B:401:PRO:HB3	2.52	0.45
1:B:8:THR:OG1	1:B:9:ALA:N	2.47	0.45
1:A:51:ALA:HA	1:A:67:LEU:HA	1.97	0.45
1:B:46:VAL:HG21	1:B:130:LEU:HB3	2.00	0.44
1:B:479:GLU:HG3	1:B:479:GLU:H	1.58	0.44
1:B:102:GLU:HB3	1:B:126:TYR:OH	2.18	0.43
1:B:203:THR:O	1:B:207:VAL:HG23	2.18	0.43
1:B:39:LYS:HG2	5:B:714:HOH:O	2.18	0.43
1:A:137:ARG:HH21	3:A:602:9YK:C	2.31	0.43
1:A:180:ARG:HD2	1:A:189:PRO:HD2	2.00	0.43
1:A:346:PHE:HE2	1:A:354:LYS:HD2	1.82	0.43
1:B:217:LEU:HB3	1:B:223:LEU:HD11	2.00	0.43
1:B:414:PRO:O	1:B:420:GLN:NE2	2.51	0.43
1:B:327:PRO:HB2	1:B:330:LYS:HD2	2.01	0.42
1:A:272:ALA:HB2	1:A:367:ILE:HD13	2.01	0.42
1:A:297:SER:HA	1:B:242:LEU:O	2.19	0.42
1:A:201:ASP:HA	1:A:202:PRO:HD3	1.92	0.42
1:B:300:TYR:CD1	1:B:448:GLN:HG2	2.55	0.42
1:B:124:SER:O	1:B:126:TYR:N	2.53	0.42
1:B:306:LYS:HA	1:B:307:PRO:HD3	1.89	0.42
1:A:126:TYR:OH	1:A:132:GLU:OE2	2.35	0.42
1:B:408:LEU:HA	1:B:411:MET:SD	2.59	0.42
1:A:135:LEU:HD22	1:A:402:TYR:CZ	2.55	0.41
1:A:19:PRO:HG3	1:A:142:ILE:HB	2.03	0.41
1:A:46:VAL:HG21	1:A:130:LEU:HB3	2.02	0.41
1:B:103:LEU:HD23	1:B:139:PHE:CE2	2.55	0.41
1:A:359:ILE:HG13	5:A:702:HOH:O	2.21	0.41
1:A:346:PHE:CE2	1:A:354:LYS:HD2	2.55	0.41
1:A:41:GLN:HG3	1:A:47:ILE:HG23	2.02	0.41
1:B:319:LEU:HD13	1:B:333:ARG:HD2	2.03	0.41
1:B:407:PHE:O	1:B:411:MET:HG3	2.22	0.41
1:A:306:LYS:HA	1:A:307:PRO:HD3	1.83	0.40
1:A:486:ALA:HB2	1:A:493:PRO:HD3	2.03	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:602:9YK:OBA	3:A:602:9YK:OAY	2.38	0.40
1:A:228:THR:HG23	1:A:231[B]:ARG:NH2	2.36	0.40
1:B:60:GLY:HA3	1:B:125:ALA:CB	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	504/502 (100%)	485 (96%)	19 (4%)	0	100	100
1	B	507/502 (101%)	480 (95%)	24 (5%)	3 (1%)	28	34
All	All	1011/1004 (101%)	965 (96%)	43 (4%)	3 (0%)	44	55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	59	ASP
1	B	309	GLN
1	B	125	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	430/425 (101%)	430 (100%)	0	100	100
1	B	433/425 (102%)	425 (98%)	8 (2%)	64	79
All	All	863/850 (102%)	855 (99%)	8 (1%)	85	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	57	SER
1	B	58	GLU
1	B	59	ASP
1	B	105	ASN
1	B	115	LYS
1	B	187	GLU
1	B	283[A]	LEU
1	B	283[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	ASN
1	B	106	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	9YK	A	602	2	24,30,31	0.70	0	33,39,41	1.50	5 (15%)
4	GOL	A	603	2	5,5,5	0.37	0	5,5,5	0.18	0
4	GOL	A	604	-	5,5,5	0.35	0	5,5,5	0.27	0
3	9YK	B	602	2	24,30,31	0.66	0	33,39,41	1.18	4 (12%)
4	GOL	B	603	-	5,5,5	0.36	0	5,5,5	0.27	0
4	GOL	B	604	-	5,5,5	0.36	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9YK	A	602	2	-	0/19/23/23	0/2/2/2
4	GOL	A	603	2	-	0/4/4/4	0/0/0/0
4	GOL	A	604	-	-	0/4/4/4	0/0/0/0
3	9YK	B	602	2	-	0/19/23/23	0/2/2/2
4	GOL	B	603	-	-	0/4/4/4	0/0/0/0
4	GOL	B	604	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	9YK	C-CA-N	-2.98	109.24	113.48
3	A	602	9YK	CAZ-CAT-NAL	-2.87	109.39	113.48
3	B	602	9YK	CAZ-CAT-NAL	-2.78	109.52	113.48
3	B	602	9YK	CAH-N-CAJ	-2.26	106.47	111.29
3	B	602	9YK	CAN-CAU-SBB	2.26	121.03	117.76
3	A	602	9YK	CAM-NAL-CAK	3.15	118.00	111.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	9YK	CBE-SBB-CAU	3.46	108.83	103.11
3	A	602	9YK	CAN-CAM-NAL	3.97	119.77	112.78
3	A	602	9YK	CBE-SBB-CAU	4.05	109.80	103.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	9YK	2	0
4	A	604	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	498/502 (99%)	0.27	9 (1%) 69 74	37, 55, 76, 98	0
1	B	498/502 (99%)	0.69	56 (11%) 6 8	36, 66, 94, 118	0
All	All	996/1004 (99%)	0.48	65 (6%) 20 26	36, 60, 89, 118	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	60	GLY	6.7
1	B	113	LEU	6.2
1	B	112	ALA	5.7
1	B	59	ASP	5.5
1	B	111	LYS	5.1
1	B	57	SER	4.9
1	B	121	THR	4.9
1	B	118	LEU	4.9
1	B	65	PHE	4.9
1	B	120	ILE	4.8
1	B	110	VAL	4.6
1	B	122	LEU	4.6
1	B	2	ALA	4.5
1	A	3	PRO	4.5
1	B	67	LEU	4.4
1	B	329	GLY	4.4
1	B	82	GLU	4.3
1	A	2	ALA	4.3
1	B	127	TYR	4.2
1	A	175	TYR	4.1
1	B	83	ALA	4.0
1	B	76	GLY	3.8
1	B	88	PHE	3.8
1	B	327	PRO	3.8

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	114	SER	3.7
1	A	192	LYS	3.6
1	B	75	ASN	3.4
1	B	119	GLN	3.3
1	B	63	TRP	3.3
1	B	115	LYS	3.3
1	B	125	ALA	3.3
1	B	3	PRO	3.2
1	B	54	TRP	3.2
1	B	331	ASP	3.2
1	B	107	ILE	3.2
1	B	382	TYR	3.2
1	A	127	TYR	3.2
1	B	58	GLU	3.1
1	B	62	THR	3.1
1	B	108	VAL	3.0
1	B	211	THR	2.8
1	B	50	LEU	2.8
1	B	323	GLY	2.7
1	B	338	GLN	2.7
1	B	47	ILE	2.7
1	B	84	ALA	2.7
1	A	172	LEU	2.6
1	B	55	THR	2.6
1	B	116	THR	2.6
1	B	96	GLN	2.4
1	B	199	ILE	2.4
1	A	499	PRO	2.3
1	B	340	LEU	2.3
1	B	160	ILE	2.3
1	B	37	LEU	2.3
1	B	442	HIS	2.3
1	B	87	ASN	2.2
1	B	61	LYS	2.1
1	A	58	GLU	2.1
1	B	191	ILE	2.0
1	B	192	LYS	2.0
1	B	375	GLY	2.0
1	A	150	HIS	2.0
1	B	56	HIS	2.0
1	B	195	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	9YK	A	602	29/30	0.88	0.28	3.12	50,74,111,116	0
3	9YK	B	602	29/30	0.94	0.16	-0.30	42,55,73,93	0
2	FE	B	601	1/1	0.98	0.14	-	51,51,51,51	0
4	GOL	A	603	6/6	0.90	0.45	-	65,76,81,84	0
4	GOL	B	604	6/6	0.77	0.21	-	57,78,85,90	0
4	GOL	A	604	6/6	0.90	0.16	-	60,79,86,98	0
4	GOL	B	603	6/6	0.68	0.21	-	65,79,85,89	0
2	FE	A	601	1/1	0.96	0.21	-	61,61,61,61	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.