



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 21, 2017 – 07:52 AM EST

PDB ID : 5ONB  
Title : Drosophila Bag-of-marbles CBM peptide bound to human CAF40  
Authors : Raisch, T.; Sgromo, A.; Backhaus, C.; Izaurralde, E.; Weichenrieder, O.  
Deposited on : 2017-08-03  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	C	273	
1	E	273	
1	G	273	
2	B	24	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	24	<div><div></div><div>4%</div><div>92%</div><div>8%</div></div>
2	F	24	<div><div></div><div>4%</div><div>88%</div><div>8%</div><div></div></div>
2	H	24	<div><div></div><div>4%</div><div>71%</div><div>17%</div><div>13%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCR4-NOT transcription complex subunit 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	0	0
			2161	1385	371	394	11			
1	C	268	Total	C	N	O	S	0	0	0
			2145	1374	369	392	10			
1	E	269	Total	C	N	O	S	0	0	0
			2153	1380	370	393	10			
1	G	270	Total	C	N	O	S	0	0	0
			2161	1385	371	394	11			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	GLY	-	expression tag	UNP Q92600
A	14	PRO	-	expression tag	UNP Q92600
A	15	HIS	-	expression tag	UNP Q92600
A	16	MET	-	expression tag	UNP Q92600
A	17	LEU	-	expression tag	UNP Q92600
A	18	GLU	-	expression tag	UNP Q92600
C	13	GLY	-	expression tag	UNP Q92600
C	14	PRO	-	expression tag	UNP Q92600
C	15	HIS	-	expression tag	UNP Q92600
C	16	MET	-	expression tag	UNP Q92600
C	17	LEU	-	expression tag	UNP Q92600
C	18	GLU	-	expression tag	UNP Q92600
E	13	GLY	-	expression tag	UNP Q92600
E	14	PRO	-	expression tag	UNP Q92600
E	15	HIS	-	expression tag	UNP Q92600
E	16	MET	-	expression tag	UNP Q92600
E	17	LEU	-	expression tag	UNP Q92600
E	18	GLU	-	expression tag	UNP Q92600
G	13	GLY	-	expression tag	UNP Q92600
G	14	PRO	-	expression tag	UNP Q92600
G	15	HIS	-	expression tag	UNP Q92600

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	16	MET	-	expression tag	UNP Q92600
G	17	LEU	-	expression tag	UNP Q92600
G	18	GLU	-	expression tag	UNP Q92600

- Molecule 2 is a protein called Protein bag-of-marbles.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	22	Total	C	N	O	S	0	0	0
			182	111	31	38	2			
2	D	22	Total	C	N	O	S	0	0	0
			182	111	31	38	2			
2	F	23	Total	C	N	O	S	0	0	0
			190	115	33	40	2			
2	H	21	Total	C	N	O	S	0	0	0
			178	109	30	37	2			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	G	1	Total	C	O	0	0
			6	3	3		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCR4-NOT transcription complex subunit 9




- Molecule 1: CCR4-NOT transcription complex subunit 9



- Molecule 1: CCR4-NOT transcription complex subunit 9





- Molecule 1: CCR4-NOT transcription complex subunit 9









- Molecule 2: Protein bag-of-marbles



- Molecule 2: Protein bag-of-marbles



- Molecule 2: Protein bag-of-marbles



- Molecule 2: Protein bag-of-marbles



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.57Å 200.93Å 59.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.73 – 3.00 46.73 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.73-3.00) 99.6 (46.73-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.216 , 0.268 0.230 , 0.277	Depositor DCC
$R_{free}$ test set	1263 reflections (5.09%)	DCC
Wilson B-factor (Å <sup>2</sup> )	79.1	Xtriage
Anisotropy	0.449	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 62.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9358	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/2203	0.65	0/2991
1	C	0.55	0/2187	0.69	0/2970
1	E	0.47	0/2195	0.62	0/2981
1	G	0.49	0/2203	0.63	0/2991
2	B	0.46	0/184	0.57	0/245
2	D	0.44	0/184	0.55	0/245
2	F	0.42	0/192	0.58	0/256
2	H	0.43	0/180	0.57	0/240
All	All	0.50	0/9528	0.64	0/12919

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2161	0	2236	25	0
1	C	2145	0	2216	32	0
1	E	2153	0	2227	13	0
1	G	2161	0	2236	27	0
2	B	182	0	163	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	182	0	163	0	0
2	F	190	0	169	3	0
2	H	178	0	160	3	0
3	G	6	0	8	0	0
All	All	9358	0	9578	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:63:ILE:CD1	1:G:63:ILE:CG1	1.76	1.61
1:G:74:TYR:HA	1:G:77:ILE:HD12	1.54	0.90
1:E:114:HIS:HE1	1:G:60:PHE:CD1	1.94	0.85
1:C:76:SER:HA	1:C:81:THR:HB	1.62	0.81
1:A:168:LEU:HB3	1:A:212:ILE:HD13	1.68	0.76
1:G:208:HIS:O	1:G:212:ILE:HG12	1.92	0.70
1:G:124:HIS:HA	1:G:169:ARG:HH22	1.62	0.65
1:A:208:HIS:O	1:A:212:ILE:HD12	1.97	0.65
1:A:177:LEU:CD2	2:B:25:GLU:HA	2.28	0.64
1:C:124:HIS:HA	1:C:169:ARG:HH22	1.62	0.64
1:E:124:HIS:HA	1:E:169:ARG:HH22	1.63	0.63
1:C:132:PHE:HB3	1:C:136:ARG:HH12	1.61	0.63
1:A:124:HIS:HA	1:A:169:ARG:HH22	1.64	0.62
1:A:174:GLY:O	1:A:179:LYS:HE2	2.00	0.61
1:E:19:ARG:HE	1:E:23:TYR:HE2	1.49	0.61
1:A:137:LEU:HD11	1:A:177:LEU:HD23	1.83	0.60
1:C:66:LEU:O	1:C:70:ILE:HG12	2.03	0.57
1:G:174:GLY:H	1:G:179:LYS:HE3	1.69	0.56
1:C:151:GLU:HB3	1:C:154:VAL:HG23	1.87	0.55
1:C:19:ARG:HA	1:C:22:ILE:HD12	1.89	0.55
1:C:180:THR:HG21	1:C:227:ARG:HG3	1.87	0.54
1:C:132:PHE:HB3	1:C:136:ARG:NH1	2.22	0.54
1:C:82:LEU:HD12	1:C:86:GLN:HB3	1.89	0.54
1:C:76:SER:CA	1:C:81:THR:HB	2.36	0.54
1:A:67:LEU:HD21	1:A:115:ILE:HG23	1.89	0.53
1:A:140:LEU:HD11	1:A:170:ILE:HG21	1.90	0.53
2:H:16:GLN:HE21	2:H:20:ASN:HD21	1.55	0.53
1:A:134:TYR:HE1	2:B:24:MET:HG3	1.72	0.53
1:G:137:LEU:HD11	1:G:177:LEU:HD23	1.92	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:168:LEU:HD13	1:C:208:HIS:CD2	2.45	0.52
1:E:114:HIS:CE1	1:G:60:PHE:CD1	2.87	0.51
1:A:203:TYR:CE2	1:G:55:MET:HG2	2.46	0.51
1:G:67:LEU:HD21	1:G:115:ILE:HG23	1.93	0.51
1:A:280:VAL:O	1:A:284:GLN:HG2	2.12	0.49
1:A:117:LEU:CD1	1:C:67:LEU:HD12	2.42	0.49
1:E:70:ILE:HG21	1:E:118:PHE:HB3	1.96	0.48
1:G:111:LEU:HD11	1:G:154:VAL:HG22	1.96	0.48
1:A:138:THR:HA	2:B:24:MET:HE2	1.95	0.48
1:A:68:GLN:HE22	1:C:120:TYR:HD2	1.61	0.48
1:A:203:TYR:CD2	1:G:55:MET:HG2	2.48	0.48
1:G:77:ILE:HD11	1:G:122:PHE:CE1	2.49	0.48
1:C:262:PHE:O	1:C:266:LEU:HB2	2.13	0.48
1:C:174:GLY:H	1:C:179:LYS:HE3	1.78	0.48
1:C:70:ILE:HG21	1:C:118:PHE:HB3	1.96	0.47
1:C:82:LEU:HD22	1:C:135:LEU:HD22	1.96	0.47
1:C:137:LEU:HD11	1:C:177:LEU:HD23	1.96	0.47
1:C:26:ILE:HG12	1:C:56:LEU:HD23	1.95	0.47
1:G:63:ILE:CD1	1:G:63:ILE:CB	2.83	0.47
1:G:110:PHE:HE2	1:G:146:LEU:HD11	1.79	0.46
1:C:213:LEU:HD22	1:C:232:VAL:HG13	1.97	0.46
1:G:101:ALA:HB1	1:G:146:LEU:HG	1.96	0.46
1:A:70:ILE:HG21	1:A:118:PHE:HB3	1.97	0.45
1:G:70:ILE:HG21	1:G:118:PHE:HB3	1.97	0.45
1:A:113:ALA:O	1:C:113:ALA:HA	2.17	0.45
1:C:114:HIS:HB3	1:C:117:LEU:HD12	1.97	0.45
1:A:177:LEU:HD22	2:B:25:GLU:HA	1.98	0.45
1:G:77:ILE:HD11	1:G:122:PHE:HE1	1.82	0.45
1:G:138:THR:HA	2:H:24:MET:CE	2.48	0.44
1:C:216:MET:O	1:C:220:LEU:HB2	2.17	0.44
1:E:134:TYR:HE1	2:F:24:MET:HG3	1.82	0.44
1:G:181:VAL:HG21	2:H:28:LEU:HD13	1.99	0.44
1:A:117:LEU:HD12	1:C:67:LEU:HD12	2.00	0.44
1:G:114:HIS:HB3	1:G:117:LEU:HD12	1.99	0.44
1:E:67:LEU:HD21	1:E:115:ILE:HG23	1.99	0.44
1:G:25:TRP:HA	1:G:28:GLU:HB2	2.00	0.44
1:E:138:THR:HA	2:F:24:MET:HE2	2.00	0.44
1:A:51:ASP:O	1:A:54:PRO:HD2	2.18	0.43
1:E:114:HIS:HB3	1:E:117:LEU:HD12	1.98	0.43
1:E:251:GLN:HG2	1:E:251:GLN:H	1.72	0.43
1:C:29:LEU:O	1:C:35:ARG:HD3	2.19	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:THR:HG22	1:E:194:THR:HA	2.01	0.42
1:A:114:HIS:CE1	1:C:60:PHE:CD1	3.07	0.42
1:E:29:LEU:O	1:E:35:ARG:HD3	2.19	0.42
1:C:19:ARG:HD3	1:C:19:ARG:HA	1.87	0.42
1:G:82:LEU:HB3	1:G:132:PHE:CE2	2.55	0.42
1:G:216:MET:HA	1:G:219:GLN:HG3	2.01	0.42
1:E:177:LEU:CD2	2:F:25:GLU:HA	2.49	0.42
1:G:51:ASP:O	1:G:54:PRO:HD2	2.20	0.41
1:A:254:PRO:HD2	1:A:257:LEU:HD12	2.02	0.41
1:G:29:LEU:O	1:G:35:ARG:HD3	2.21	0.41
1:A:29:LEU:O	1:A:35:ARG:HD3	2.20	0.41
1:C:37:ASN:O	1:C:41:GLU:HG2	2.20	0.41
1:C:216:MET:HA	1:C:219:GLN:HG3	2.03	0.41
1:A:70:ILE:HD11	1:A:97:LEU:CD1	2.51	0.41
1:C:77:ILE:HG23	1:C:125:THR:HG22	2.03	0.41
1:C:266:LEU:HD21	1:C:273:LYS:HA	2.03	0.41
1:C:67:LEU:O	1:C:70:ILE:HB	2.21	0.40
1:G:39:LEU:HD11	1:G:69:GLU:HG3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/273 (98%)	259 (97%)	9 (3%)	0	100	100
1	C	266/273 (97%)	256 (96%)	8 (3%)	2 (1%)	22	64
1	E	267/273 (98%)	258 (97%)	9 (3%)	0	100	100
1	G	268/273 (98%)	260 (97%)	8 (3%)	0	100	100
2	B	20/24 (83%)	20 (100%)	0	0	100	100
2	D	20/24 (83%)	20 (100%)	0	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	21/24 (88%)	21 (100%)	0	0	100	100
2	H	19/24 (79%)	19 (100%)	0	0	100	100
All	All	1149/1188 (97%)	1113 (97%)	34 (3%)	2 (0%)	51	86

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	82	LEU
1	C	80	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/246 (99%)	239 (98%)	5 (2%)	60	87
1	C	242/246 (98%)	235 (97%)	7 (3%)	48	82
1	E	243/246 (99%)	238 (98%)	5 (2%)	59	87
1	G	244/246 (99%)	240 (98%)	4 (2%)	68	90
2	B	20/22 (91%)	20 (100%)	0	100	100
2	D	20/22 (91%)	20 (100%)	0	100	100
2	F	21/22 (96%)	21 (100%)	0	100	100
2	H	20/22 (91%)	20 (100%)	0	100	100
All	All	1054/1072 (98%)	1033 (98%)	21 (2%)	60	87

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	73	ILE
1	A	99	CYS
1	A	146	LEU
1	A	219	GLN
1	A	284	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	19	ARG
1	C	73	ILE
1	C	99	CYS
1	C	149	THR
1	C	163	ILE
1	C	219	GLN
1	C	267	LYS
1	E	73	ILE
1	E	99	CYS
1	E	146	LEU
1	E	219	GLN
1	E	251	GLN
1	G	73	ILE
1	G	99	CYS
1	G	130	ARG
1	G	219	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	114	HIS
1	A	208	HIS
1	A	219	GLN
1	A	284	GLN
2	D	16	GLN
1	E	114	HIS
1	E	219	GLN
1	G	98	GLN
2	H	16	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	G	301	-	5,5,5	0.15	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	G	301	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	270/273 (98%)	-0.33	1 (0%) 92 77	42, 81, 130, 164	0
1	C	268/273 (98%)	0.41	23 (8%) 11 4	64, 110, 178, 203	0
1	E	269/273 (98%)	-0.05	3 (1%) 80 55	59, 100, 155, 196	0
1	G	270/273 (98%)	-0.16	5 (1%) 67 37	53, 86, 131, 186	0
2	B	22/24 (91%)	0.04	1 (4%) 34 13	75, 90, 144, 160	0
2	D	22/24 (91%)	0.36	1 (4%) 34 13	92, 109, 159, 166	0
2	F	23/24 (95%)	0.15	1 (4%) 36 15	94, 114, 168, 184	0
2	H	21/24 (87%)	-0.33	1 (4%) 31 12	90, 101, 131, 144	0
All	All	1165/1188 (98%)	-0.03	36 (3%) 49 22	42, 95, 160, 203	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	13	ASP	6.6
2	B	13	ASP	6.5
1	C	81	THR	4.9
1	C	51	ASP	4.3
1	G	16	MET	4.0
1	C	34	THR	3.9
1	C	237	LEU	3.7
1	C	257	LEU	3.6
1	C	221	SER	3.3
1	C	252	CYS	3.2
1	C	80	PRO	3.1
1	C	52	LEU	3.0
1	A	257	LEU	3.0
1	C	37	ASN	2.9
2	F	35	ASN	2.8
1	C	49	VAL	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	33	GLU	2.7
1	C	217	VAL	2.7
1	C	128	LYS	2.5
1	C	284	GLN	2.5
1	G	284	GLN	2.4
1	E	285	GLU	2.4
1	C	206	PHE	2.4
1	E	49	VAL	2.4
2	H	13	ASP	2.3
1	C	279	LEU	2.3
1	G	253	LEU	2.3
1	C	18	GLU	2.3
1	E	43	SER	2.3
1	G	17	LEU	2.2
1	C	43	SER	2.2
1	C	78	ASN	2.2
1	C	242	ASN	2.2
1	G	285	GLU	2.2
1	C	269	ASP	2.1
1	C	285	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	G	301	6/6	0.77	0.21	-	101,102,103,103	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.