



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:44 am GMT

PDB ID : 4OON  
Title : Crystal structure of PBP1a in complex with compound 17 ((4Z,8S,11E,14S)-5-(2-amino-1,3-thiazol-4-yl)-14-(5,6-dihydroxy-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)-8-formyl-2-methyl-6-oxo-3,10-dioxo-4,7,11-triazatetradeca-4,11-diene-2,12,14-tricarboxylic acid)  
Authors : Han, S.; Caspers, N.; Knafels, J.D.  
Deposited on : 2014-02-03  
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1015 (3.22-3.18)
Clashscore	112137	1009 (3.20-3.20)
Ramachandran outliers	110173	1118 (3.22-3.18)
Sidechain outliers	110143	1117 (3.22-3.18)
RSRZ outliers	101464	1020 (3.22-3.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	795	<div> <div> <div></div> <div>50%</div> <div>11%</div> <div>•</div> <div>37%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3959 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Penicillin-binding protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	502	Total	C	N	O	S	0	0	0
			3911	2470	688	737	16			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	INITIATING METHIONINE	UNP Q07806
A	29	SER	-	EXPRESSION TAG	UNP Q07806
A	30	HIS	-	EXPRESSION TAG	UNP Q07806
A	31	HIS	-	EXPRESSION TAG	UNP Q07806
A	32	HIS	-	EXPRESSION TAG	UNP Q07806
A	33	HIS	-	EXPRESSION TAG	UNP Q07806
A	34	HIS	-	EXPRESSION TAG	UNP Q07806
A	35	HIS	-	EXPRESSION TAG	UNP Q07806

- Molecule 2 is (4Z,8S,11E,14S)-5-(2-AMINO-1,3-THIAZOL-4-YL)-14-(5,6-DIHYDROXY-1,3-DIOXO-1,3-DIHYDRO-2H-ISOINDOL-2-YL)-8-FORMYL-2-METHYL-6-OXO-3,10-DIOXA-4,7,11-TRIAZATETRADECA-4,11-DIENE-2,12,14-TRICARBOXYLIC ACID (three-letter code: 2U4) (formula: C<sub>25</sub>H<sub>24</sub>N<sub>6</sub>O<sub>14</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			46	25	6	14	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total O 2 2	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.77Å 113.77Å 123.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.72 – 3.20 83.66 – 3.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (27.72-3.20) 100.0 (83.66-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.32 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, $R_{free}$	0.168 , 0.216 0.188 , 0.227	Depositor DCC
$R_{free}$ test set	695 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	86.5	Xtriage
Anisotropy	0.374	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 82.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3959	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2U4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/3997	0.76	2/5413 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	492	GLU	N-CA-C	6.12	127.52	111.00
1	A	508	LEU	C-N-CA	5.00	132.81	122.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3911	0	3868	40	0
2	A	46	0	18	0	0
3	A	2	0	0	0	0
All	All	3959	0	3886	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:ASN:HD21	1:A:730:ARG:H	1.32	0.77
1:A:430:PRO:HG3	1:A:715:VAL:HG23	1.71	0.72
1:A:564:VAL:HG23	1:A:569:ILE:HD11	1.71	0.71
1:A:296:GLN:HE22	1:A:438:LEU:H	1.39	0.69
1:A:483:ASN:HA	1:A:510:PRO:HA	1.75	0.68
1:A:339:ARG:HH22	1:A:445:GLU:HA	1.58	0.68
1:A:372:ASP:HA	1:A:375:LYS:HD2	1.77	0.66
1:A:537:ALA:O	1:A:541:ILE:HG12	1.97	0.65
1:A:433:GLY:O	1:A:585:PRO:HA	1.98	0.64
1:A:512:PRO:HG2	1:A:515:GLU:HB2	1.83	0.60
1:A:320:GLU:HB2	1:A:402:GLN:HG2	1.82	0.60
1:A:316:TYR:HA	1:A:417:GLN:HE22	1.67	0.60
1:A:48:LEU:O	1:A:60:GLU:HA	2.07	0.54
1:A:564:VAL:CG2	1:A:569:ILE:HD11	2.37	0.54
1:A:303:VAL:HG13	1:A:423:SER:HB3	1.91	0.53
1:A:713:ASP:OD1	1:A:756:HIS:HD2	1.92	0.53
1:A:323:LEU:HD22	1:A:326:GLN:HE22	1.75	0.52
1:A:591:ILE:HB	1:A:600:TYR:HB3	1.93	0.51
1:A:264:ILE:HG13	1:A:291:VAL:HG11	1.92	0.51
1:A:578:ASN:HB2	1:A:607:VAL:HG11	1.93	0.50
1:A:357:ILE:HG13	1:A:369:VAL:HB	1.95	0.49
1:A:674:ASP:HA	1:A:677:LYS:HB2	1.93	0.49
1:A:515:GLU:HA	1:A:518:TYR:CE2	2.49	0.47
1:A:768:ARG:HD2	1:A:784:ALA:HB1	1.97	0.47
1:A:301:GLN:HG2	1:A:305:ASP:OD2	2.15	0.47
1:A:465:PRO:HG3	1:A:672:LEU:HD21	1.97	0.46
1:A:518:TYR:HA	1:A:675:VAL:HG22	1.99	0.45
1:A:506:THR:HG21	1:A:523:MET:HG2	1.99	0.45
1:A:426:ILE:HD12	1:A:438:LEU:HD12	1.99	0.44
1:A:661:ASP:HB3	1:A:664:THR:OG1	2.17	0.44
1:A:768:ARG:HG2	1:A:786:PHE:CZ	2.52	0.44
1:A:262:PRO:HB2	1:A:444:PHE:CG	2.53	0.42
1:A:290:THR:HB	1:A:434:ALA:HB1	2.01	0.42
1:A:405:VAL:HB	1:A:413:LEU:HD22	2.01	0.42
1:A:427:SER:HA	1:A:715:VAL:O	2.19	0.41
1:A:322:ARG:C	1:A:324:PRO:HD3	2.40	0.41
1:A:351:GLN:HB2	1:A:358:MET:HB2	2.03	0.41
1:A:593:SER:HB3	1:A:599:LEU:HD12	2.02	0.40
1:A:708:SER:HG	1:A:717:SER:HG	1.69	0.40
1:A:768:ARG:HG2	1:A:786:PHE:CE1	2.56	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	494/795 (62%)	463 (94%)	26 (5%)	5 (1%)	18 61

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	508	LEU
1	A	283	GLU
1	A	377	ALA
1	A	507	PHE
1	A	791	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	411/657 (63%)	378 (92%)	33 (8%)	14 49

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	260	ASN
1	A	287	VAL
1	A	321	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	326	GLN
1	A	332	LEU
1	A	344	LEU
1	A	354	LYS
1	A	361	THR
1	A	372	ASP
1	A	382	SER
1	A	403	ILE
1	A	409	GLU
1	A	443	SER
1	A	445	GLU
1	A	492	GLU
1	A	494	LEU
1	A	507	PHE
1	A	532	LEU
1	A	595	ASP
1	A	599	LEU
1	A	609	VAL
1	A	680	THR
1	A	700	ASN
1	A	701	ASP
1	A	708	SER
1	A	730	ARG
1	A	731	ARG
1	A	757	THR
1	A	765	VAL
1	A	766	SER
1	A	767	LEU
1	A	768	ARG
1	A	776	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	GLN
1	A	297	ASN
1	A	312	GLN
1	A	326	GLN
1	A	384	ASN
1	A	554	ASN
1	A	673	GLN
1	A	756	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	2U4	A	901	1	36,48,48	1.20	3 (8%)	42,70,70	2.09	15 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2U4	A	901	1	-	0/25/64/64	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	2U4	C30-C1	-3.00	1.47	1.52
2	A	901	2U4	C35-N34	-2.01	1.36	1.40
2	A	901	2U4	C7-S6	3.85	1.76	1.70

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	2U4	C39-C40-C35	-3.06	105.67	108.27
2	A	901	2U4	C19-C16-C18	-3.02	105.44	110.20
2	A	901	2U4	C40-C39-C36	-2.40	106.23	108.27
2	A	901	2U4	O37-C35-C40	-2.33	124.09	128.68
2	A	901	2U4	O38-C36-C39	-2.30	124.16	128.68
2	A	901	2U4	C8-C7-S6	-2.18	109.12	111.79
2	A	901	2U4	C2-C3-N34	2.45	115.90	112.46
2	A	901	2U4	C39-C36-N34	2.62	108.39	105.89
2	A	901	2U4	C12-C13-N23	2.74	119.05	114.41
2	A	901	2U4	C41-C40-C35	2.76	133.81	129.29
2	A	901	2U4	C28-O27-N26	2.79	111.70	108.47
2	A	901	2U4	C42-C39-C36	2.90	134.04	129.29
2	A	901	2U4	C40-C35-N34	3.43	109.16	105.89
2	A	901	2U4	C2-C3-C4	3.73	116.41	110.81
2	A	901	2U4	O15-N14-C12	7.01	122.18	112.03

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	502/795 (63%)	0.32	8 (1%) 72 59	51, 79, 106, 138	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	506	THR	3.0
1	A	423	SER	2.6
1	A	424	ALA	2.6
1	A	720	VAL	2.3
1	A	705	GLY	2.2
1	A	708	SER	2.2
1	A	64	MET	2.2
1	A	487	ILE	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	2U4	A	901	46/46	0.91	0.26	-0.36	69,101,122,125	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.