



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2018 – 01:08 PM EST

PDB ID : 5OQU
Title : The crystal structure of CK2alpha in complex with compound 5
Authors : Brear, P.; De Fusco, C.; Iegre, J.; Yoshida, M.; Mitchell, S.; Rossmann, M.; Carro, L.; Sore, H.; Hyvonen, M.; Spring, D.
Deposited on : 2017-08-14
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4787 (2.34-2.30)
Clashscore	112137	5439 (2.34-2.30)
Ramachandran outliers	110173	5386 (2.34-2.30)
Sidechain outliers	110143	5385 (2.34-2.30)
RSRZ outliers	101464	4814 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	
1	B	352	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	405	-	-	-	X
3	ACT	A	406	-	-	-	X
3	ACT	B	402	-	-	-	X
3	ACT	B	403	-	-	-	X
6	I6P	A	408	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	324	Total	C	N	O	S	0	3	0
			2763	1770	485	497	11			
1	A	324	Total	C	N	O	S	0	3	0
			2763	1770	485	497	11			

There are 50 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	GLY	-	expression tag	UNP P68400
B	-21	SER	-	expression tag	UNP P68400
B	-20	MET	-	expression tag	UNP P68400
B	-19	ASP	-	expression tag	UNP P68400
B	-18	ILE	-	expression tag	UNP P68400
B	-17	GLU	-	expression tag	UNP P68400
B	-16	PHE	-	expression tag	UNP P68400
B	-15	ASP	-	expression tag	UNP P68400
B	-14	ASP	-	expression tag	UNP P68400
B	-13	ASP	-	expression tag	UNP P68400
B	-12	ALA	-	expression tag	UNP P68400
B	-11	ASP	-	expression tag	UNP P68400
B	-10	ASP	-	expression tag	UNP P68400
B	-9	ASP	-	expression tag	UNP P68400
B	-8	GLY	-	expression tag	UNP P68400
B	-7	SER	-	expression tag	UNP P68400
B	-6	GLY	-	expression tag	UNP P68400
B	-5	SER	-	expression tag	UNP P68400
B	-4	GLY	-	expression tag	UNP P68400
B	-3	SER	-	expression tag	UNP P68400
B	-2	GLY	-	expression tag	UNP P68400
B	-1	SER	-	expression tag	UNP P68400
B	0	GLY	-	expression tag	UNP P68400
B	1	SER	-	expression tag	UNP P68400
B	21	SER	ARG	engineered mutation	UNP P68400

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	GLY	-	expression tag	UNP P68400
A	-21	SER	-	expression tag	UNP P68400
A	-20	MET	-	expression tag	UNP P68400
A	-19	ASP	-	expression tag	UNP P68400
A	-18	ILE	-	expression tag	UNP P68400
A	-17	GLU	-	expression tag	UNP P68400
A	-16	PHE	-	expression tag	UNP P68400
A	-15	ASP	-	expression tag	UNP P68400
A	-14	ASP	-	expression tag	UNP P68400
A	-13	ASP	-	expression tag	UNP P68400
A	-12	ALA	-	expression tag	UNP P68400
A	-11	ASP	-	expression tag	UNP P68400
A	-10	ASP	-	expression tag	UNP P68400
A	-9	ASP	-	expression tag	UNP P68400
A	-8	GLY	-	expression tag	UNP P68400
A	-7	SER	-	expression tag	UNP P68400
A	-6	GLY	-	expression tag	UNP P68400
A	-5	SER	-	expression tag	UNP P68400
A	-4	GLY	-	expression tag	UNP P68400
A	-3	SER	-	expression tag	UNP P68400
A	-2	GLY	-	expression tag	UNP P68400
A	-1	SER	-	expression tag	UNP P68400
A	0	GLY	-	expression tag	UNP P68400
A	1	SER	-	expression tag	UNP P68400
A	21	SER	ARG	engineered mutation	UNP P68400

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is [3-chloranyl-4-(2-methoxyphenyl)phenyl]methanamine (three-letter code: A4B) (formula: C₁₄H₁₄ClNO).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			17	14	1	1	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is INOSITOL 1,2,3,4,5,6-HEXAKISPHOSPHATE (three-letter code: I6P) (formula: C₆H₁₈O₂₄P₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	P	0	0
			36	6	24	6		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	B	2	Total O 2 2	0	0
8	A	13	Total O 13 13	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.67Å 67.92Å 332.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	166.40 – 2.32 166.40 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.4 (166.40-2.32) 99.8 (166.40-2.32)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.32Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R, R_{free}	0.220 , 0.244 0.228 , 0.252	Depositor DCC
R_{free} test set	1600 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	54.1	Xtriage
Anisotropy	0.663	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.135 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5671	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, A4B, ADP, CL, I6P, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/2838	0.65	0/3839
1	B	0.52	0/2838	0.65	0/3839
All	All	0.54	0/5676	0.65	0/7678

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2763	0	2702	13	3
1	B	2763	0	2702	17	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	12	0	9	0	0
3	B	8	0	6	0	0
4	A	17	0	0	4	0
5	A	2	0	0	0	0
6	A	36	0	6	0	3
7	A	1	0	0	1	0
8	A	13	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	2	0	0	0	0
All	All	5671	0	5449	28	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:ILE:HG23	1:B:263:ILE:HB	1.66	0.77
1:B:224:SER:HB2	1:B:230:GLU:HG2	1.73	0.69
1:A:224:SER:HB2	1:A:230:GLU:HG2	1.74	0.68
4:A:402:A4B:N	7:A:409:CL:CL	2.64	0.66
1:B:266:ASP:OD1	1:B:267:PRO:HD2	2.03	0.59
1:B:50:TYR:HB3	1:B:71:LYS:HB3	1.86	0.57
1:B:50:TYR:CD1	1:B:50:TYR:N	2.72	0.55
1:B:158:LYS:HE2	1:B:194[A]:SER:OG	2.07	0.54
1:A:225:MET:CG	4:A:402:A4B:C2	2.87	0.53
1:A:5:VAL:HB	1:A:261:TYR:HA	1.92	0.52
1:B:20:PRO:HD2	1:A:105:VAL:HG21	1.91	0.51
1:B:20:PRO:CG	1:A:105:VAL:HG11	2.41	0.51
1:B:303:LYS:HB3	1:B:313:LEU:HG	1.94	0.50
1:A:225:MET:HG2	4:A:402:A4B:C2	2.41	0.50
1:A:120:ASP:HB2	1:A:123:GLN:HB2	1.96	0.48
1:A:134:ARG:HG2	1:A:323:TYR:CZ	2.48	0.47
1:B:120:ASP:HB3	1:B:123:GLN:HB2	1.96	0.47
1:B:5:VAL:HB	1:B:261:TYR:HA	1.98	0.45
1:B:20:PRO:HG2	1:A:105:VAL:HG11	1.97	0.45
1:A:224:SER:CB	1:A:230:GLU:HG2	2.45	0.45
1:A:225:MET:HG3	4:A:402:A4B:C2	2.47	0.44
1:B:224:SER:CB	1:B:230:GLU:HG2	2.44	0.42
1:A:258:ILE:HD13	1:A:265:LEU:HG	2.02	0.42
1:B:95:ILE:HB	1:B:174:ILE:HG22	2.02	0.41
1:B:128:LEU:HD22	1:B:132:ASP:HB3	2.01	0.41
1:B:118:ASN:HD22	1:B:164:ILE:H	1.69	0.41
1:B:42:VAL:HG23	1:B:56:ALA:HA	2.02	0.41
1:A:128:LEU:HD22	1:A:132:ASP:HB3	2.02	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:LYS:NZ	6:A:408:I6P:O32[4_597]	1.52	0.68
1:A:229:LYS:CE	6:A:408:I6P:O32[4_597]	2.02	0.18
1:A:234:HIS:ND1	6:A:408:I6P:O31[4_597]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/352 (92%)	310 (95%)	13 (4%)	2 (1%)	28	34
1	B	325/352 (92%)	308 (95%)	17 (5%)	0	100	100
All	All	650/704 (92%)	618 (95%)	30 (5%)	2 (0%)	44	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	267	PRO
1	A	72	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/319 (95%)	301 (100%)	1 (0%)	94	97
1	B	302/319 (95%)	302 (100%)	0	100	100
All	All	604/638 (95%)	603 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	58	ASN
1	B	118	ASN
1	A	118	ASN
1	A	262	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	A	401	5	25,29,29	0.72	0	24,45,45	0.77	0
4	A4B	A	402	-	18,18,18	0.14	0	24,24,24	0.26	0
3	ACT	A	405	-	1,3,3	5.16	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	A	406	-	1,3,3	5.20	1 (100%)	0,3,3	0.00	-
3	ACT	A	407	-	1,3,3	6.51	1 (100%)	0,3,3	0.00	-
6	I6P	A	408	-	36,36,36	1.05	3 (8%)	54,60,60	0.60	1 (1%)
2	ADP	B	401	-	25,29,29	1.05	2 (8%)	24,45,45	1.78	3 (12%)
3	ACT	B	402	-	1,3,3	5.70	1 (100%)	0,3,3	0.00	-
3	ACT	B	403	-	1,3,3	4.83	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	401	5	-	0/12/32/32	0/3/3/3
4	A4B	A	402	-	-	0/8/8/8	0/2/2/2
3	ACT	A	405	-	-	0/0/0/0	0/0/0/0
3	ACT	A	406	-	-	0/0/0/0	0/0/0/0
3	ACT	A	407	-	-	0/0/0/0	0/0/0/0
6	I6P	A	408	-	-	0/30/54/54	0/1/1/1
2	ADP	B	401	-	-	0/12/32/32	0/3/3/3
3	ACT	B	402	-	-	0/0/0/0	0/0/0/0
3	ACT	B	403	-	-	0/0/0/0	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	ADP	C2'-C1'	-2.40	1.49	1.53
6	A	408	I6P	P4-O14	2.22	1.63	1.59
2	B	401	ADP	C5-C4	2.31	1.45	1.40
6	A	408	I6P	P2-O12	2.99	1.64	1.59
6	A	408	I6P	P1-O11	4.15	1.66	1.59
3	B	403	ACT	CH3-C	4.83	1.55	1.48
3	A	405	ACT	CH3-C	5.16	1.55	1.48
3	A	406	ACT	CH3-C	5.20	1.55	1.48
3	B	402	ACT	CH3-C	5.70	1.56	1.48
3	A	407	ACT	CH3-C	6.51	1.57	1.48

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	ADP	N3-C2-N1	-6.49	123.20	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	401	ADP	C4-C5-N7	-3.24	106.28	109.41
2	B	401	ADP	C2-N1-C6	2.30	122.79	118.77
6	A	408	I6P	C2-C1-C6	2.63	116.30	110.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	402	A4B	4	0
6	A	408	I6P	0	3

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/352 (92%)	0.79	21 (6%) 20 26	34, 58, 96, 114	0
1	B	324/352 (92%)	1.20	74 (22%) 1 1	46, 81, 115, 140	0
All	All	648/704 (92%)	0.99	95 (14%) 3 4	34, 70, 111, 140	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	307	TYR	5.9
1	B	50	TYR	5.5
1	B	292	LEU	5.2
1	A	33	TRP	5.0
1	B	257	TYR	4.7
1	B	273	LEU	4.7
1	A	325	TYR	4.6
1	B	77	LYS	4.1
1	B	325	TYR	4.0
1	B	41	LEU	3.9
1	B	269	PHE	3.9
1	B	242	LEU	3.9
1	B	33	TRP	3.9
1	A	72	PRO	3.9
1	B	54	PHE	3.8
1	B	204	VAL	3.8
1	B	223	ALA	3.8
1	B	267	PRO	3.7
1	B	225	MET	3.7
1	B	75	LYS	3.5
1	B	254	LEU	3.5
1	B	125	TYR	3.5
1	B	62	ASN	3.4
1	B	227	PHE	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	42	VAL	3.4
1	A	71	LYS	3.4
1	B	216	TRP	3.3
1	B	76	LYS	3.3
1	A	50	TYR	3.3
1	A	271	ASP	3.3
1	B	249	LEU	3.2
1	B	265	LEU	3.2
1	B	245	ILE	3.1
1	B	206	TYR	3.1
1	B	270	ASN	3.0
1	B	250	GLY	3.0
1	B	326	THR	3.0
1	B	243	VAL	2.9
1	A	128	LEU	2.9
1	B	261	TYR	2.9
1	B	30	VAL	2.9
1	B	28	SER	2.9
1	B	298	LEU	2.9
1	B	305	LEU	2.9
1	B	51	SER	2.8
1	A	273	LEU	2.8
1	B	72	PRO	2.7
1	A	51	SER	2.7
1	B	5	VAL	2.7
1	B	202	LEU	2.7
1	B	281	TRP	2.7
1	B	171	LEU	2.7
1	B	178	LEU	2.7
1	A	124	LEU	2.6
1	B	121	PHE	2.6
1	B	173	LEU	2.6
1	A	70	LEU	2.6
1	B	255	TYR	2.6
1	B	239	TYR	2.5
1	B	52	GLU	2.5
1	B	271	ASP	2.4
1	B	66	VAL	2.4
1	B	124	LEU	2.4
1	B	57	ILE	2.4
1	A	326	THR	2.4
1	B	59	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	269	PHE	2.3
1	B	46	GLY	2.3
1	B	74	LYS	2.3
1	A	284	PHE	2.3
1	B	263	ILE	2.3
1	A	268	ARG	2.2
1	A	272	ILE	2.2
1	B	64	LYS	2.2
1	B	26	TYR	2.2
1	A	76	LYS	2.2
1	A	327	VAL	2.2
1	B	21	SER	2.2
1	B	82	ILE	2.2
1	B	213	LEU	2.2
1	B	69	ILE	2.2
1	B	100	ILE	2.2
1	B	45	LEU	2.2
1	B	244	ARG	2.2
1	B	272	ILE	2.1
1	A	222	LEU	2.1
1	B	160	HIS	2.1
1	B	232	PHE	2.1
1	A	239	TYR	2.1
1	B	191	ARG	2.0
1	A	225	MET	2.0
1	B	67	VAL	2.0
1	B	126	GLN	2.0
1	B	31	VAL	2.0
1	B	70	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ACT	B	403	4/4	0.81	0.33	6.23	78,81,83,86	0
3	ACT	A	406	4/4	0.91	0.26	5.63	66,70,71,73	0
3	ACT	A	405	4/4	0.81	0.31	4.87	72,73,74,75	0
3	ACT	B	402	4/4	0.71	0.28	4.12	79,80,80,81	0
6	I6P	A	408	36/36	0.74	0.24	2.33	124,128,132,133	36
7	CL	A	409	1/1	0.91	0.24	1.64	93,93,93,93	0
4	A4B	A	402	17/17	0.79	0.25	0.94	62,69,79,90	0
2	ADP	B	401	27/27	0.87	0.24	0.51	96,108,126,128	0
2	ADP	A	401	27/27	0.93	0.16	-0.61	54,71,87,90	0
5	MG	A	404	1/1	0.97	0.09	-3.06	64,64,64,64	0
5	MG	A	403	1/1	0.90	0.12	-	61,61,61,61	0
3	ACT	A	407	4/4	0.94	0.29	-	61,63,63,64	0

6.5 Other polymers

There are no such residues in this entry.