



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 03:22 pm GMT

PDB ID : 1OR7
Title : Crystal Structure of Escherichia coli sigmaE with the Cytoplasmic Domain of its Anti-sigma RseA
Authors : Campbell, E.A.; Tupy, J.L.; Gruber, T.M.; Wang, S.; Sharp, M.M.; Gross, C.A.; Darst, S.A.
Deposited on : 2003-03-12
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	<div> <div>13%</div> <div>63%</div> <div>26%</div> <div>7%</div> </div>
1	B	194	<div> <div>4%</div> <div>65%</div> <div>15%</div> <div>15%</div> </div>
2	C	90	<div> <div>6%</div> <div>49%</div> <div>20%</div> <div>27%</div> </div>
2	F	90	<div> <div>4%</div> <div>51%</div> <div>14%</div> <div>6%</div> <div>29%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA polymerase sigma-E factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	27	3	0
			1457	915	261	276	5			
1	B	164	Total	C	N	O	S	0	4	0
			1338	846	243	245	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	CLONING ARTIFACT	UNP P0AGB6
A	-1	SER	-	CLONING ARTIFACT	UNP P0AGB6
A	0	HIS	-	CLONING ARTIFACT	UNP P0AGB6
B	-2	GLY	-	CLONING ARTIFACT	UNP P0AGB6
B	-1	SER	-	CLONING ARTIFACT	UNP P0AGB6
B	0	HIS	-	CLONING ARTIFACT	UNP P0AGB6

- Molecule 2 is a protein called Sigma-E factor negative regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	66	Total	C	N	O	S	11	1	0
			536	329	89	113	5			
2	F	64	Total	C	N	O	S	0	0	0
			514	317	87	105	5			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total	O	0	0
			33	33		
3	B	69	Total	O	0	0
			69	69		
3	C	31	Total	O	0	0
			31	31		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	42	Total	O	0	0
			42	42		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: RNA polymerase sigma-E factor




• Molecule 1: RNA polymerase sigma-E factor



• Molecule 2: Sigma-E factor negative regulatory protein



• Molecule 2: Sigma-E factor negative regulatory protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.05Å 56.90Å 104.98Å 90.00° 130.51° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 34.33 – 1.99	Depositor EDS
% Data completeness (in resolution range)	100.0 (30.00-2.00) 99.0 (34.33-1.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.197 , 0.231 0.215 , 0.237	Depositor DCC
R_{free} test set	2134 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.187	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 59.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4020	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	5/1498 (0.3%)	0.92	8/2024 (0.4%)
1	B	0.81	4/1378 (0.3%)	0.97	5/1860 (0.3%)
2	C	0.93	4/549 (0.7%)	0.83	2/739 (0.3%)
2	F	1.02	4/522 (0.8%)	0.90	1/703 (0.1%)
All	All	0.87	17/3947 (0.4%)	0.92	16/5326 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	1

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	136	GLU	CB-CG	13.33	1.77	1.52
1	A	1	MET	CA-CB	-9.72	1.32	1.53
2	F	29	MET	CG-SD	7.55	2.00	1.81
1	B	163	MET	CG-SD	7.49	2.00	1.81
2	F	60	MET	CG-SD	7.48	2.00	1.81

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	28	ARG	NE-CZ-NH2	-13.33	113.64	120.30
1	A	1	MET	N-CA-CB	12.28	132.71	110.60
1	A	186	GLN	C-N-CD	-11.69	94.88	120.60
1	B	28	ARG	NE-CZ-NH1	10.32	125.46	120.30
1	A	141	ASP	CB-CG-OD2	7.21	124.79	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	1	MET	CA

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	186	GLN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1457	0	1451	61	6
1	B	1338	0	1351	44	6
2	C	536	0	509	20	0
2	F	514	0	495	28	0
3	A	33	0	0	2	0
3	B	69	0	0	1	0
3	C	31	0	0	1	0
3	F	42	0	0	4	0
All	All	4020	0	3806	126	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:10:MET:SD	2:C:10:MET:CE	2.02	1.47
2:C:29:MET:SD	2:C:29:MET:CE	2.01	1.46
1:B:163:MET:CE	1:B:163:MET:SD	2.03	1.46
2:F:43:MET:SD	2:F:43:MET:CE	2.05	1.43
1:A:144:MET:CE	1:A:148:LEU:HG	1.81	1.11

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:CE	1:B:91:ARG:CG[4_556]	0.85	1.35
1:A:1:MET:CE	1:B:91:ARG:CB[4_556]	0.97	1.23
1:A:1:MET:SD	1:B:91:ARG:CD[4_556]	1.74	0.46
1:A:1:MET:SD	1:B:91:ARG:CB[4_556]	1.83	0.37
1:A:1:MET:CE	1:B:91:ARG:CD[4_556]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	180/194 (93%)	175 (97%)	3 (2%)	2 (1%)	17	9
1	B	164/194 (84%)	161 (98%)	2 (1%)	1 (1%)	28	21
2	C	65/90 (72%)	64 (98%)	1 (2%)	0	100	100
2	F	62/90 (69%)	62 (100%)	0	0	100	100
All	All	471/568 (83%)	462 (98%)	6 (1%)	3 (1%)	28	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	110	GLY
1	A	186	GLN
1	B	90	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/168 (95%)	155 (98%)	4 (2%)	53	54
1	B	145/168 (86%)	140 (97%)	5 (3%)	42	40
2	C	61/81 (75%)	55 (90%)	6 (10%)	9	5
2	F	58/81 (72%)	56 (97%)	2 (3%)	42	40
All	All	423/498 (85%)	406 (96%)	17 (4%)	38	32

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	22[B]	GLU
2	C	31	LYS
1	B	133	ARG
2	C	22[A]	GLU
1	B	184	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	GLN
2	C	25	HIS
2	F	21	ASN
1	A	183	ASN
1	B	23	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/194 (93%)	0.63	25 (13%) 3 3	8, 18, 46, 59	6 (3%)
1	B	164/194 (84%)	0.06	7 (4%) 36 36	11, 18, 37, 52	0
2	C	66/90 (73%)	0.32	5 (7%) 15 15	10, 16, 37, 47	3 (4%)
2	F	64/90 (71%)	0.20	4 (6%) 21 21	10, 16, 23, 32	0
All	All	475/568 (83%)	0.33	41 (8%) 11 11	8, 18, 41, 59	9 (1%)

The worst 5 of 41 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	111	ALA	10.3
1	B	90	GLY	9.1
1	A	108	SER	7.1
1	A	85[A]	TYR	6.3
1	A	109	GLY	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.