



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 14, 2017 – 12:44 AM EST

PDB ID : 4OZO
Title : Crystal structure of an α -L-fucosidase GH29 from *Bacteroides thetaiotaomicron* (BT2192) in complex with oNPTG
Authors : Lafite, P.; Daniellou, R.; Guillotin, L.
Deposited on : unknown
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	469	<div> <div>2%</div> <div>87%</div> <div>10%</div> <div>•</div> </div>
1	B	469	<div> <div>3%</div> <div>84%</div> <div>14%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	501	-	-	-	X
2	GOL	B	502	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative lipoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	2	0	0
			3650	2332	618	684	16			
1	B	459	Total	C	N	O	S	3	0	0
			3650	2332	618	684	16			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	23	MET	-	expression tag	UNP Q8A5P6
A	24	SER	-	expression tag	UNP Q8A5P6
A	25	LEU	-	expression tag	UNP Q8A5P6
A	484	GLU	-	expression tag	UNP Q8A5P6
A	485	GLY	-	expression tag	UNP Q8A5P6
A	486	HIS	-	expression tag	UNP Q8A5P6
A	487	HIS	-	expression tag	UNP Q8A5P6
A	488	HIS	-	expression tag	UNP Q8A5P6
A	489	HIS	-	expression tag	UNP Q8A5P6
A	490	HIS	-	expression tag	UNP Q8A5P6
A	491	HIS	-	expression tag	UNP Q8A5P6
B	23	MET	-	expression tag	UNP Q8A5P6
B	24	SER	-	expression tag	UNP Q8A5P6
B	25	LEU	-	expression tag	UNP Q8A5P6
B	484	GLU	-	expression tag	UNP Q8A5P6
B	485	GLY	-	expression tag	UNP Q8A5P6
B	486	HIS	-	expression tag	UNP Q8A5P6
B	487	HIS	-	expression tag	UNP Q8A5P6
B	488	HIS	-	expression tag	UNP Q8A5P6
B	489	HIS	-	expression tag	UNP Q8A5P6
B	490	HIS	-	expression tag	UNP Q8A5P6
B	491	HIS	-	expression tag	UNP Q8A5P6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is (2R,3R,4S,5R,6S)-2-(hydroxymethyl)-6-(2-nitrophenyl)sulfanyl-oxane-3,4,5-triol (three-letter code: OTN) (formula: C₁₂H₁₅NO₇S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			21	12	1	7	1		
3	B	1	Total	C	N	O	S	0	0
			21	12	1	7	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total	O	0	0
			106	106		
4	B	97	Total	O	0	0
			97	97		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative lipoprotein



• Molecule 1: Putative lipoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	125.30Å 122.17Å 159.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 2.60 49.28 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.00-2.60) 100.0 (49.28-2.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.52 (at 2.61Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.165 , 0.220 0.157 , 0.218	Depositor DCC
R_{free} test set	1899 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	37.8	Xtriage
Anisotropy	0.879	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.074 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7569	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OTN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3753	0.53	0/5106
1	B	0.39	0/3753	0.54	0/5106
All	All	0.39	0/7506	0.53	0/10212

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3650	0	3485	28	0
1	B	3650	0	3485	35	0
2	A	12	0	16	3	0
2	B	12	0	16	2	0
3	A	21	0	15	2	0
3	B	21	0	15	2	0
4	A	106	0	0	2	0
4	B	97	0	0	2	0
All	All	7569	0	7032	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:ARG:NH2	1:B:214:CYS:O	2.22	0.71
1:A:339:GLN:NE2	1:B:343:GLU:OE2	2.25	0.67
1:A:57:GLU:OE2	4:A:650:HOH:O	2.15	0.64
1:A:310:ARG:HH21	2:A:502:GOL:H12	1.61	0.64
1:A:284:TRP:CD1	2:A:501:GOL:H11	2.34	0.62
1:A:99:HIS:NE2	2:A:501:GOL:O2	2.26	0.60
1:B:351:ARG:NH2	1:B:371:LEU:O	2.35	0.59
1:B:369:LYS:HD2	1:B:377:THR:HB	1.84	0.57
1:B:24:SER:N	4:B:689:HOH:O	2.37	0.56
1:B:387:LYS:NZ	1:B:462:GLN:O	2.38	0.56
1:B:343:GLU:HB2	1:B:479:SER:HB2	1.88	0.55
1:A:425:ASP:OD1	1:A:426:GLY:N	2.39	0.55
1:B:96:THR:HA	1:B:145:TYR:HB3	1.89	0.55
1:A:406:GLU:OE2	1:A:444:LYS:NZ	2.38	0.53
1:B:421:GLU:HG2	1:B:431:THR:HA	1.90	0.53
1:A:393:THR:HA	1:A:456:GLN:HG2	1.91	0.53
1:B:389:ASP:OD1	1:B:458:ARG:HD2	2.10	0.52
1:B:129:ARG:NH1	4:B:672:HOH:O	2.39	0.51
1:A:328:SER:HB3	1:B:328:SER:HB2	1.92	0.51
1:B:312:SER:OG	1:B:313:VAL:N	2.42	0.51
1:A:350:LEU:HG	1:A:475:VAL:HG22	1.93	0.51
1:B:367:PRO:HA	1:B:379:PHE:HB2	1.92	0.50
1:A:323:ARG:HG2	1:A:325:LEU:HG	1.94	0.49
1:A:336:LEU:HD21	1:B:339:GLN:HG3	1.94	0.49
1:A:425:ASP:OD1	1:A:427:LYS:N	2.36	0.49
1:B:408:VAL:HG22	1:B:471:HIS:CD2	2.48	0.49
1:A:116:LYS:NZ	4:A:601:HOH:O	2.37	0.48
3:A:503:OTN:S01	3:A:503:OTN:O3'	2.72	0.48
1:A:105:LEU:HB3	1:A:127:VAL:HB	1.95	0.47
1:B:39:TRP:CE2	1:B:228:VAL:HG11	2.50	0.47
1:B:230:TRP:CE2	3:B:503:OTN:H6	2.50	0.47
1:B:423:THR:HG22	1:B:456:GLN:HB2	1.96	0.47
1:A:369:LYS:HD2	1:A:377:THR:HB	1.96	0.47
3:B:503:OTN:S01	3:B:503:OTN:O3'	2.73	0.47
1:B:56:GLU:O	1:B:322:ARG:NH2	2.48	0.47
1:B:393:THR:HA	1:B:456:GLN:HG2	1.97	0.46
1:B:425:ASP:OD1	1:B:427:LYS:N	2.43	0.46
1:A:298:ARG:NH1	1:A:482:PHE:O	2.49	0.46
1:A:224:PRO:HG2	1:A:225:PHE:CE2	2.50	0.46
1:A:39:TRP:CE2	1:A:228:VAL:HG11	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:LEU:HD12	1:B:450:ALA:O	2.16	0.45
1:B:49:HIS:NE2	2:B:502:GOL:O2	2.41	0.44
1:A:389:ASP:OD1	1:A:458:ARG:NH2	2.48	0.44
1:B:120:TRP:CZ3	1:B:125:GLY:HA3	2.53	0.44
1:A:323:ARG:O	1:A:323:ARG:HG3	2.17	0.43
1:B:200:TYR:CD1	1:B:222:SER:HB3	2.54	0.43
1:B:306:THR:HA	1:B:310:ARG:HG3	2.01	0.43
1:B:146:LEU:HD23	1:B:168:TYR:CZ	2.53	0.43
1:B:296:SER:N	1:B:299:GLU:OE2	2.40	0.43
1:B:204:TYR:CE1	1:B:208:ARG:HD2	2.54	0.43
1:A:223:TYR:CG	1:A:224:PRO:HD3	2.54	0.42
1:A:39:TRP:CZ2	1:A:313:VAL:HB	2.55	0.42
1:B:310:ARG:HH21	2:B:501:GOL:H12	1.85	0.42
1:A:120:TRP:CZ3	1:A:125:GLY:HA3	2.55	0.42
1:B:353:ALA:HB2	1:B:394:LEU:HD23	2.02	0.41
1:B:39:TRP:HB2	1:B:215:VAL:HG21	2.03	0.41
1:B:342:ASP:O	1:B:346:SER:HB3	2.21	0.41
1:B:418:TRP:CE2	1:B:438:LYS:HB2	2.56	0.41
1:A:106:TRP:CE2	1:A:108:SER:HB2	2.56	0.40
1:A:406:GLU:HB2	1:A:472:THR:HB	2.03	0.40
1:A:96:THR:HA	1:A:145:TYR:HB3	2.03	0.40
1:A:230:TRP:CE2	3:A:503:OTN:H6	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/469 (97%)	440 (96%)	16 (4%)	1 (0%)	51	76
1	B	457/469 (97%)	441 (96%)	16 (4%)	0	100	100
All	All	914/938 (97%)	881 (96%)	32 (4%)	1 (0%)	55	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	377/393 (96%)	367 (97%)	10 (3%)	50	77
1	B	377/393 (96%)	364 (97%)	13 (3%)	42	69
All	All	754/786 (96%)	731 (97%)	23 (3%)	45	73

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	TYR
1	A	47	PHE
1	A	58	TYR
1	A	122	ASN
1	A	145	TYR
1	A	186	TRP
1	A	231	MET
1	A	389	ASP
1	A	397	THR
1	A	449	LEU
1	B	40	TYR
1	B	41	ASN
1	B	47	PHE
1	B	58	TYR
1	B	145	TYR
1	B	186	TRP
1	B	202	HIS
1	B	231	MET
1	B	293	ARG
1	B	330	ASP
1	B	389	ASP
1	B	397	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	449	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	501	-	5,5,5	1.20	0	5,5,5	0.38	0
2	GOL	A	502	-	5,5,5	0.98	0	5,5,5	0.51	0
3	OTN	A	503	-	20,22,22	1.95	3 (15%)	25,31,31	1.29	3 (12%)
2	GOL	B	501	-	5,5,5	0.96	0	5,5,5	0.23	0
2	GOL	B	502	-	5,5,5	1.18	0	5,5,5	0.43	0
3	OTN	B	503	-	20,22,22	2.26	4 (20%)	25,31,31	1.20	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	0/4/4/4	0/0/0/0
2	GOL	A	502	-	-	0/4/4/4	0/0/0/0
3	OTN	A	503	-	-	0/8/30/30	0/2/2/2
2	GOL	B	501	-	-	0/4/4/4	0/0/0/0
2	GOL	B	502	-	-	0/4/4/4	0/0/0/0
3	OTN	B	503	-	-	0/8/30/30	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	503	OTN	C1'-S01	-5.66	1.70	1.77
3	A	503	OTN	C1-C2	-2.32	1.49	1.53
3	B	503	OTN	C1-S01	-2.01	1.78	1.82
3	B	503	OTN	O5-C1	2.40	1.46	1.42
3	A	503	OTN	O5-C1	3.11	1.47	1.42
3	A	503	OTN	O3'-N1'	7.25	1.35	1.22
3	B	503	OTN	O3'-N1'	7.38	1.36	1.22

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	503	OTN	C6'-C1'-S01	-2.63	116.74	121.58
3	A	503	OTN	C6'-C1'-S01	-2.52	116.93	121.58
3	A	503	OTN	O2-C2-C3	-2.21	105.55	110.36
3	B	503	OTN	O5-C5-C6	2.08	111.39	106.41
3	B	503	OTN	C2'-C1'-S01	2.26	124.25	122.14
3	A	503	OTN	C2'-C1'-S01	2.77	124.74	122.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	GOL	2	0
2	A	502	GOL	1	0
3	A	503	OTN	2	0
2	B	501	GOL	1	0
2	B	502	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	503	OTN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	459/469 (97%)	-0.20	11 (2%)	59 52	7, 17, 33, 59	3 (0%)
1	B	459/469 (97%)	-0.20	12 (2%)	56 49	7, 17, 34, 62	4 (0%)
All	All	918/938 (97%)	-0.20	23 (2%)	58 50	7, 17, 33, 62	7 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	24	SER	3.7
1	A	24	SER	3.7
1	A	352	GLY	3.2
1	B	355	VAL	3.0
1	A	393	THR	2.9
1	B	368	GLU	2.7
1	B	351	ARG	2.7
1	B	347	THR	2.5
1	B	424	VAL	2.5
1	A	351	ARG	2.5
1	A	368	GLU	2.5
1	A	424	VAL	2.5
1	B	457	VAL	2.5
1	A	355	VAL	2.4
1	A	376	ASN	2.4
1	B	423	THR	2.2
1	B	455	LYS	2.1
1	B	363	ALA	2.1
1	B	376	ASN	2.1
1	B	357	ALA	2.1
1	A	363	ALA	2.1
1	A	427	LYS	2.1
1	A	394	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	B	501	6/6	0.85	0.20	4.83	25,28,33,34	0
2	GOL	B	502	6/6	0.94	0.18	4.24	10,13,17,17	0
2	GOL	A	502	6/6	0.87	0.17	1.75	22,24,30,36	0
3	OTN	A	503	21/21	0.94	0.18	1.58	16,25,31,54	0
3	OTN	B	503	21/21	0.97	0.17	1.27	9,17,36,48	0
2	GOL	A	501	6/6	0.97	0.15	0.96	15,16,25,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.