



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:44 am GMT

PDB ID : 1P18  
Title : Hypoxanthine Phosphoribosyltransferase from Trypanosoma cruzi, K68R mutant, ternary substrates complex  
Authors : Canyuk, B.; Eakin, A.E.; Craig III, S.P.  
Deposited on : 2003-04-11  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3202 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypoxanthine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1464	943	248	266	7			
1	B	194	Total	C	N	O	S	0	0	0
			1561	998	271	285	7			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	LYS	MET	SEE REMARK 999	UNP Q27796
A	52	ARG	LYS	ENGINEERED	UNP Q27796
A	66	CYS	SER	SEE REMARK 999	UNP Q27796
A	86	LEU	VAL	SEE REMARK 999	UNP Q27796
B	23	LYS	MET	SEE REMARK 999	UNP Q27796
B	52	ARG	LYS	ENGINEERED	UNP Q27796
B	66	CYS	SER	SEE REMARK 999	UNP Q27796
B	86	LEU	VAL	SEE REMARK 999	UNP Q27796

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Mg	0	0
			2	2		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 7-HYDROXY-PYRAZOLO[4,3-D]PYRIMIDINE (three-letter code: 7HP) (formula: C<sub>5</sub>H<sub>4</sub>N<sub>4</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	5	4	1		
3	B	1	Total	C	N	O	0	0
			10	5	4	1		

- Molecule 4 is ALPHA-PHOSPHORIBOSYLPYROPHOSPHORIC ACID (three-letter code: PRP) (formula:  $C_5H_{13}O_{14}P_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			22	5	14	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	0
			22	5	14	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	60	Total	O	0	0
			60	60		
5	B	49	Total	O	0	0
			49	49		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: hypoxanthine phosphoribosyltransferase

Chain A: 



- Molecule 1: hypoxanthine phosphoribosyltransferase

Chain B: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	39.50Å 102.10Å 51.80Å 90.00° 94.30° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.00)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.210 , 0.293	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3202	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 7HP, PRP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/1494	0.77	0/2032
1	B	0.57	0/1592	0.76	0/2157
All	All	0.56	0/3086	0.76	0/4189

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	172	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1464	0	1438	36	0
1	B	1561	0	1554	36	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	10	0	3	1	0
3	B	10	0	3	0	0
4	A	22	0	6	6	0
4	B	22	0	6	0	0
5	A	60	0	0	1	0
5	B	49	0	0	2	0
All	All	3202	0	3010	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ILE:HG23	4:A:801:PRP:H51	1.47	0.96
1:B:116:THR:HB	1:B:148:ARG:HB2	1.68	0.76
1:B:79:VAL:HA	1:B:93:ARG:O	1.87	0.74
1:A:23:LYS:HE2	1:A:68:PHE:CZ	2.23	0.73
1:B:84:GLU:HB2	1:B:197:ARG:NH2	2.06	0.70
1:B:79:VAL:HG22	1:B:92:VAL:HG12	1.72	0.69
1:B:27:LYS:HD2	1:B:68:PHE:CE2	2.28	0.68
1:A:21:ARG:HB3	1:A:157:VAL:HG13	1.79	0.65
1:A:118:LEU:HD11	1:A:148:ARG:HG2	1.79	0.63
1:A:69:ASN:HD22	1:A:69:ASN:N	1.96	0.63
1:B:116:THR:OG1	1:B:118:LEU:HG	1.99	0.61
1:A:28:ARG:NH1	1:A:31:ASP:HB2	2.16	0.61
1:B:80:SER:HB3	1:B:93:ARG:HD2	1.81	0.60
1:B:79:VAL:CG2	1:B:92:VAL:HG12	2.32	0.59
1:B:197:ARG:HG3	1:B:197:ARG:HH11	1.67	0.56
1:A:164:PHE:HE2	1:A:172:TYR:HB2	1.72	0.55
1:A:23:LYS:HE2	1:A:68:PHE:CE1	2.42	0.54
1:A:8:ALA:C	1:A:186:ARG:HG3	2.29	0.54
1:B:125:HIS:O	1:B:129:THR:HG23	2.09	0.53
1:B:21:ARG:HB3	1:B:157:VAL:HG13	1.91	0.52
1:B:197:ARG:HG3	1:B:197:ARG:NH1	2.24	0.52
1:B:136:LYS:HE3	5:B:2002:HOH:O	2.09	0.52
1:A:113:ILE:HG22	4:A:801:PRP:H2	1.91	0.52
1:A:90:GLY:HA2	5:A:6000:HOH:O	2.09	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:TYR:HE1	5:B:1514:HOH:O	1.93	0.51
3:A:800:7HP:H9	4:A:801:PRP:O4	2.12	0.50
1:B:187:PRO:HA	1:B:190:TYR:CE2	2.46	0.50
1:A:64:ALA:O	1:A:67:ASP:HB2	2.13	0.49
1:B:193:ARG:O	1:B:197:ARG:HG2	2.13	0.49
1:A:33:TYR:HB3	1:A:38:LEU:HD11	1.93	0.49
1:B:193:ARG:CZ	1:B:197:ARG:HD3	2.43	0.48
1:A:115:ASP:OD1	1:A:146:GLY:HA3	2.12	0.48
1:B:102:ILE:HB	1:B:107:VAL:HG21	1.96	0.48
1:A:140:LEU:HD21	1:A:182:ILE:HD11	1.96	0.47
1:A:52:ARG:HG2	1:B:76:PHE:HD1	1.79	0.47
1:A:58:THR:HG22	1:A:74:MET:CE	2.45	0.46
1:B:51:LEU:HB3	1:B:52:ARG:HA	1.98	0.46
1:A:164:PHE:CE2	1:A:172:TYR:HB2	2.51	0.46
1:A:69:ASN:N	1:A:69:ASN:ND2	2.60	0.45
1:A:87:THR:HG23	1:A:88:SER:N	2.31	0.45
1:A:58:THR:HG22	1:A:74:MET:HE1	1.98	0.45
1:A:28:ARG:HH12	1:A:31:ASP:HB2	1.79	0.45
1:A:7:PHE:HA	1:A:189:VAL:HG21	1.97	0.45
1:A:113:ILE:O	4:A:801:PRP:H52	2.16	0.45
1:A:171:ASP:HB3	1:A:177:ARG:HG2	1.99	0.45
1:B:140:LEU:O	1:B:157:VAL:HB	2.17	0.45
1:A:87:THR:HG23	1:A:88:SER:H	1.82	0.44
1:A:75:GLU:HB3	1:A:98:THR:HG22	1.99	0.44
1:A:113:ILE:CG2	4:A:801:PRP:H2	2.48	0.43
1:B:69:ASN:O	1:B:71:PRO:HD3	2.18	0.43
1:A:118:LEU:HD21	1:A:148:ARG:HG3	1.99	0.43
1:A:44:PRO:HD2	1:A:105:HIS:NE2	2.32	0.43
1:A:165:VAL:HG23	1:A:170:LEU:HD22	1.99	0.43
1:B:50:VAL:O	1:B:54:SER:HB3	2.19	0.43
1:B:142:ASP:HB3	1:B:159:ASN:ND2	2.34	0.43
1:B:176:TYR:O	1:B:179:LEU:HG	2.18	0.43
1:A:115:ASP:HB3	4:A:801:PRP:O3P	2.19	0.42
1:B:84:GLU:O	1:B:197:ARG:NH2	2.52	0.42
1:A:52:ARG:HG2	1:B:76:PHE:CD1	2.54	0.42
1:B:145:GLU:O	1:B:162:ASN:ND2	2.52	0.42
1:B:39:ARG:HB2	1:B:42:VAL:CG2	2.50	0.42
1:A:102:ILE:HB	1:A:107:VAL:HG21	2.02	0.41
1:B:84:GLU:HB3	1:B:173:ASP:OD1	2.19	0.41
1:B:91:GLN:H	1:B:91:GLN:HG2	1.67	0.41
1:A:21:ARG:HD2	1:A:21:ARG:HA	1.81	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:TYR:HE2	1:B:130:ARG:NH1	2.17	0.41
1:B:52:ARG:NH1	1:B:174:ASP:OD2	2.53	0.41
1:B:51:LEU:CB	1:B:52:ARG:HA	2.50	0.41
1:B:141:LEU:HD21	1:B:182:ILE:HD13	2.01	0.41
1:A:95:LEU:HA	1:A:95:LEU:HD23	1.80	0.40
1:B:115:ASP:OD1	1:B:143:LYS:HD2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	185/221 (84%)	174 (94%)	11 (6%)	0	100	100
1	B	192/221 (87%)	184 (96%)	8 (4%)	0	100	100
All	All	377/442 (85%)	358 (95%)	19 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/194 (79%)	149 (97%)	5 (3%)	44	42
1	B	167/194 (86%)	164 (98%)	3 (2%)	64	68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	321/388 (83%)	313 (98%)	8 (2%)	53 54

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ARG
1	A	69	ASN
1	A	87	THR
1	A	93	ARG
1	A	174	ASP
1	B	16	GLU
1	B	84	GLU
1	B	99	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	ASN
1	A	121	ASN
1	B	105	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	7HP	A	800	-	10,11,11	1.72	3 (30%)	4,15,15	4.39	4 (100%)
4	PRP	A	801	2	20,22,22	2.51	2 (10%)	30,35,35	2.26	11 (36%)
3	7HP	B	810	-	10,11,11	2.08	5 (50%)	4,15,15	4.28	3 (75%)
4	PRP	B	811	2	20,22,22	1.57	4 (20%)	30,35,35	1.90	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	7HP	A	800	-	-	0/0/0/0	0/2/2/2
4	PRP	A	801	2	-	0/16/33/33	0/1/1/1
3	7HP	B	810	-	-	0/0/0/0	0/2/2/2
4	PRP	B	811	2	-	0/16/33/33	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	801	PRP	PB-O3A	-7.53	1.47	1.60
4	A	801	PRP	C1-C2	-6.67	1.44	1.52
4	B	811	PRP	PB-O3A	-3.18	1.54	1.60
3	B	810	7HP	N8-N7	-3.01	1.32	1.37
4	B	811	PRP	PB-O2B	-2.97	1.42	1.54
3	A	800	7HP	N8-N7	-2.74	1.32	1.37
4	B	811	PRP	O4-C4	2.13	1.49	1.45
3	A	800	7HP	C2-N3	2.48	1.36	1.32
3	B	810	7HP	C2-N1	2.69	1.38	1.33
3	B	810	7HP	C2-N3	2.69	1.36	1.32
3	A	800	7HP	C2-N1	2.73	1.39	1.33
4	B	811	PRP	C3-C4	2.75	1.60	1.53
3	B	810	7HP	C5-N7	2.85	1.43	1.37
3	B	810	7HP	C6-N1	3.22	1.44	1.36

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	810	7HP	N3-C2-N1	-7.00	122.76	128.86
3	A	800	7HP	C2-N3-C4	-5.87	109.11	115.27
4	B	811	PRP	O3P-P-O5	-5.17	92.98	106.73
4	A	801	PRP	O3-C3-C4	-5.08	96.24	111.09
3	A	800	7HP	N3-C2-N1	-4.96	124.54	128.86
4	A	801	PRP	O2-C2-C1	-4.54	99.39	111.85
4	B	811	PRP	O2-C2-C1	-3.75	101.57	111.85
4	B	811	PRP	C2-C3-C4	-3.30	96.19	102.62
4	A	801	PRP	C1-C2-C3	-3.25	98.12	102.29
4	B	811	PRP	O3-C3-C4	-2.91	102.58	111.09
4	A	801	PRP	O3A-PA-O1	-2.43	97.71	102.05
4	A	801	PRP	O3P-P-O5	-2.08	101.20	106.73
4	A	801	PRP	O4-C4-C5	2.08	116.42	109.40
3	B	810	7HP	C2-N1-C6	2.16	125.09	117.26
3	A	800	7HP	C2-N1-C6	2.48	126.23	117.26
4	B	811	PRP	O3P-P-O2P	2.48	117.62	107.61
4	A	801	PRP	O5-C5-C4	2.51	117.89	109.00
4	B	811	PRP	O4-C4-C3	2.66	110.45	105.17
4	B	811	PRP	O4-C1-C2	2.71	108.54	104.97
4	A	801	PRP	P-O5-C5	2.78	125.96	118.30
3	A	800	7HP	O6-C6-C5	3.43	126.29	119.71
4	A	801	PRP	O4-C1-O1	4.01	116.99	108.97
4	B	811	PRP	O1-C1-C2	4.02	113.66	106.70
3	B	810	7HP	O6-C6-C5	4.18	127.75	119.71
4	A	801	PRP	O2-C2-C3	4.30	125.61	111.83
4	A	801	PRP	O4-C1-C2	4.68	111.12	104.97

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	800	7HP	1	0
4	A	801	PRP	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.