



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 07:03 am GMT

PDB ID : 1P19  
Title : Hypoxanthine Phosphoribosyltransferase from Trypanosoma cruzi, in complex with the product IMP  
Authors : Medrano, F.J.; Eakin, A.E.; Craig III, S.P.  
Deposited on : 2003-04-11  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	221	
1	B	221	
1	C	221	
1	D	221	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6928 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called hypoxanthine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	187	Total	C	N	O	S	0	0	0
			1519	978	259	276	6			
1	B	184	Total	C	N	O	S	0	0	0
			1507	967	262	272	6			
1	C	185	Total	C	N	O	S	0	0	0
			1507	970	257	274	6			
1	D	186	Total	C	N	O	S	0	0	0
			1523	977	265	275	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	23	LYS	MET	SEE REMARK 999	UNP Q27796
A	66	CYS	SER	SEE REMARK 999	UNP Q27796
A	86	LEU	VAL	SEE REMARK 999	UNP Q27796
B	23	LYS	MET	SEE REMARK 999	UNP Q27796
B	66	CYS	SER	SEE REMARK 999	UNP Q27796
B	86	LEU	VAL	SEE REMARK 999	UNP Q27796
C	23	LYS	MET	SEE REMARK 999	UNP Q27796
C	66	CYS	SER	SEE REMARK 999	UNP Q27796
C	86	LEU	VAL	SEE REMARK 999	UNP Q27796
D	23	LYS	MET	SEE REMARK 999	UNP Q27796
D	66	CYS	SER	SEE REMARK 999	UNP Q27796
D	86	LEU	VAL	SEE REMARK 999	UNP Q27796

- Molecule 2 is INOSINIC ACID (three-letter code: IMP) (formula: C<sub>10</sub>H<sub>13</sub>N<sub>4</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
2	B	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
2	C	1	Total	C	N	O	P	0	0
			23	10	4	8	1		
2	D	1	Total	C	N	O	P	0	0
			23	10	4	8	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	202	Total	O	0	0
			202	202		
3	B	193	Total	O	0	0
			193	193		
3	C	183	Total	O	0	0
			183	183		
3	D	202	Total	O	0	0
			202	202		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: hypoxanthine phosphoribosyltransferase




- Molecule 1: hypoxanthine phosphoribosyltransferase




- Molecule 1: hypoxanthine phosphoribosyltransferase





● Molecule 1: hypoxanthine phosphoribosyltransferase



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	156.94Å 121.02Å 52.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.30)	Depositor
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.193 , 0.261	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6928	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/1550	0.99	8/2099 (0.4%)
1	B	0.45	3/1536 (0.2%)	0.93	1/2078 (0.0%)
1	C	0.33	0/1538	0.91	4/2082 (0.2%)
1	D	0.30	0/1552	0.86	2/2100 (0.1%)
All	All	0.36	3/6176 (0.0%)	0.92	15/8359 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4	GLU	CB-CG	8.18	1.67	1.52
1	B	4	GLU	CD-OE1	5.56	1.31	1.25
1	B	4	GLU	CD-OE2	5.47	1.31	1.25

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	28	ARG	CD-NE-CZ	7.54	134.15	123.60
1	A	131	ARG	NE-CZ-NH2	6.45	123.52	120.30
1	A	28	ARG	CD-NE-CZ	6.22	132.31	123.60
1	A	28	ARG	NE-CZ-NH1	6.05	123.32	120.30
1	A	131	ARG	CD-NE-CZ	5.98	131.98	123.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1519	0	1526	116	0
1	B	1507	0	1514	117	0
1	C	1507	0	1509	123	0
1	D	1523	0	1531	124	0
2	A	23	0	11	6	0
2	B	23	0	11	1	0
2	C	23	0	11	4	0
2	D	23	0	11	1	0
3	A	202	0	0	14	0
3	B	193	0	0	16	0
3	C	183	0	0	13	0
3	D	202	0	0	12	0
All	All	6928	0	6124	463	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 463 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:102:ILE:HD12	1:B:107:VAL:HG21	1.43	0.97
1:A:15:GLU:HA	1:A:18:ILE:HD12	1.49	0.94
1:D:193:ARG:HH21	1:D:197:ARG:HD3	1.32	0.92
1:A:185:LEU:HD11	1:A:189:VAL:HG11	1.54	0.89
1:C:102:ILE:HD11	1:C:130:ARG:HB2	1.58	0.86

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/221 (83%)	167 (91%)	15 (8%)	1 (0%)	32	39
1	B	180/221 (81%)	163 (91%)	13 (7%)	4 (2%)	8	6
1	C	181/221 (82%)	162 (90%)	16 (9%)	3 (2%)	11	9
1	D	182/221 (82%)	163 (90%)	17 (9%)	2 (1%)	17	18
All	All	726/884 (82%)	655 (90%)	61 (8%)	10 (1%)	13	13

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	173	ASP
1	D	162	ASN
1	A	146	GLY
1	B	197	ARG
1	B	10	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/194 (86%)	134 (81%)	32 (19%)	1	1
1	B	162/194 (84%)	128 (79%)	34 (21%)	1	1
1	C	164/194 (84%)	127 (77%)	37 (23%)	1	1
1	D	164/194 (84%)	132 (80%)	32 (20%)	1	1
All	All	656/776 (84%)	521 (79%)	135 (21%)	1	1

5 of 135 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	174	ASP
1	C	24	GLU
1	D	147	ARG
1	B	188	GLU
1	C	6	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	162	ASN
1	D	105	HIS
1	B	198	GLN
1	B	159	ASN
1	D	43	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	IMP	A	1300	-	21,25,25	1.60	3 (14%)	22,38,38	2.06	7 (31%)
2	IMP	B	2300	-	21,25,25	1.64	3 (14%)	22,38,38	2.01	6 (27%)
2	IMP	C	3300	-	21,25,25	1.59	3 (14%)	22,38,38	2.01	5 (22%)
2	IMP	D	4300	-	21,25,25	1.62	3 (14%)	22,38,38	2.14	5 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMP	A	1300	-	-	0/6/26/26	0/3/3/3
2	IMP	B	2300	-	-	0/6/26/26	0/3/3/3
2	IMP	C	3300	-	-	0/6/26/26	0/3/3/3
2	IMP	D	4300	-	-	0/6/26/26	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2300	IMP	C8-N7	-2.71	1.29	1.34
2	A	1300	IMP	C8-N7	-2.60	1.29	1.34
2	C	3300	IMP	C8-N7	-2.59	1.29	1.34
2	D	4300	IMP	C8-N7	-2.56	1.29	1.34
2	C	3300	IMP	C2-N1	2.59	1.38	1.33

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4300	IMP	N3-C2-N1	-6.40	123.28	128.86
2	C	3300	IMP	N3-C2-N1	-6.27	123.40	128.86
2	A	1300	IMP	N3-C2-N1	-5.94	123.68	128.86
2	B	2300	IMP	N3-C2-N1	-5.86	123.75	128.86
2	B	2300	IMP	C4'-O4'-C1'	-3.43	106.12	109.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1300	IMP	6	0
2	B	2300	IMP	1	0
2	C	3300	IMP	4	0
2	D	4300	IMP	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.