



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:53 am GMT

PDB ID : 1P1V
Title : Crystal Structure of FALS-associated human Copper-Zinc Superoxide Dismutase (CuZnSOD) Mutant D125H to 1.4Å
Authors : Elam, J.S.; Malek, K.; Rodriguez, J.A.; Doucette, P.A.; Taylor, A.B.; Hayward, L.J.; Cabelli, D.E.; Valentine, J.S.; Hart, P.J.
Deposited on : 2003-04-14
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1307 (1.40-1.40)
Clashscore	112137	1411 (1.40-1.40)
Ramachandran outliers	110173	1373 (1.40-1.40)
Sidechain outliers	110143	1372 (1.40-1.40)
RSRZ outliers	101464	1315 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>5% • 5%</div> </div> </div>
1	B	153	<div> <div>14%</div> <div> <div></div> <div>76%</div> <div>14% • 6%</div> </div> </div>
1	C	153	<div> <div>%</div> <div> <div></div> <div>86%</div> <div>8% • 5%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Cu-Zn].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1066	656	197	209	4			
1	B	144	Total	C	N	O	S	0	0	0
			1048	644	193	207	4			
1	C	146	Total	C	N	O	S	0	0	0
			1066	656	197	209	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	111	CSX	CYS	MODIFIED RESIDUE	UNP P00441
A	125	HIS	ASP	ENGINEERED	UNP P00441
B	111	CSX	CYS	MODIFIED RESIDUE	UNP P00441
B	125	HIS	ASP	ENGINEERED	UNP P00441
C	111	CSX	CYS	MODIFIED RESIDUE	UNP P00441
C	125	HIS	ASP	ENGINEERED	UNP P00441

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	296	Total	O	0	0
			296	296		
4	B	148	Total	O	0	0
			148	148		
4	C	292	Total	O	0	0
			292	292		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Superoxide dismutase [Cu-Zn]



• Molecule 1: Superoxide dismutase [Cu-Zn]



• Molecule 1: Superoxide dismutase [Cu-Zn]



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.45Å 101.14Å 143.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.40 36.79 – 1.40	Depositor EDS
% Data completeness (in resolution range)	93.9 (10.00-1.40) 94.6 (36.79-1.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 1.40Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.146 , 0.212 0.188 , 0.203	Depositor DCC
R_{free} test set	2908 reflections (3.06%)	DCC
Wilson B-factor (Å ²)	12.8	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 63.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3937	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, CSX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/1076	1.03	0/1447
1	B	0.48	0/1058	1.20	5/1425 (0.4%)
1	C	0.60	0/1076	1.07	1/1447 (0.1%)
All	All	0.56	0/3210	1.10	6/4319 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	83	ASP	CB-CG-OD1	9.46	126.82	118.30
1	B	115	ARG	NE-CZ-NH2	7.39	124.00	120.30
1	C	69	ARG	CD-NE-CZ	6.75	133.05	123.60
1	B	96	ASP	CB-CG-OD1	6.47	124.12	118.30
1	B	79	ARG	NE-CZ-NH1	-6.16	117.22	120.30
1	B	79	ARG	NE-CZ-NH2	5.22	122.91	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1066	0	1043	6	0
1	B	1048	0	1017	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1066	0	1043	11	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	296	0	0	1	0
4	B	148	0	0	7	0
4	C	292	0	0	7	0
All	All	3937	0	3103	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:LYS:HG2	4:B:832:HOH:O	1.84	0.77
1:B:70:LYS:HE2	1:B:78:GLU:OE2	1.89	0.71
1:B:1:ALA:HA	1:B:22:GLN:O	1.91	0.70
1:B:23:LYS:HD3	4:B:836:HOH:O	1.94	0.67
1:A:136:LYS:HG3	1:A:136:LYS:O	1.96	0.65
1:B:58:THR:HG23	4:B:916:HOH:O	1.99	0.62
1:B:66:PRO:HG2	1:B:67:LEU:HD13	1.83	0.60
1:B:23:LYS:CE	1:B:24:GLU:HB3	2.38	0.53
1:A:125:HIS:CD2	1:A:139:ASN:HB3	2.44	0.52
1:A:2:THR:HG21	4:A:1041:HOH:O	2.09	0.52
1:B:28:PRO:HD2	4:B:874:HOH:O	2.10	0.51
1:B:23:LYS:HE3	1:B:24:GLU:HB3	1.92	0.51
1:B:24:GLU:HG3	1:B:25:SER:N	2.25	0.51
1:A:125:HIS:HD2	1:A:139:ASN:HB3	1.77	0.50
1:C:126:LEU:HB2	1:C:128:LYS:HE3	1.93	0.50
1:B:27:GLY:HA3	4:B:836:HOH:O	2.11	0.49
1:C:125:HIS:CE1	1:C:138:GLY:HA2	2.48	0.49
1:B:118:VAL:HG11	1:B:143:ARG:HG2	1.93	0.49
1:C:23:LYS:HB2	4:C:896:HOH:O	2.12	0.47
1:C:128:LYS:HD2	4:C:991:HOH:O	2.16	0.46
1:C:1:ALA:N	4:C:966:HOH:O	2.49	0.45
1:B:69:ARG:NH1	4:B:882:HOH:O	2.50	0.44
1:C:3:LYS:NZ	4:C:978:HOH:O	2.50	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2:THR:HG22	4:C:1072:HOH:O	2.17	0.44
1:C:91:LYS:NZ	4:C:1060:HOH:O	2.50	0.43
1:B:66:PRO:HG2	1:B:67:LEU:CD1	2.48	0.43
1:C:136:LYS:HZ3	1:C:136:LYS:HB2	1.84	0.42
1:A:121:GLU:HA	1:A:144:LEU:HD11	2.02	0.42
1:C:8:LEU:HD11	1:C:117:LEU:HD23	2.01	0.41
1:B:58:THR:HG23	4:B:923:HOH:O	2.21	0.41
1:C:36:LYS:HD2	4:C:875:HOH:O	2.21	0.41
1:A:125:HIS:HD2	1:A:138:GLY:O	2.04	0.40
1:B:23:LYS:HG2	1:B:24:GLU:N	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/153 (92%)	139 (99%)	2 (1%)	0	100	100
1	B	139/153 (91%)	136 (98%)	3 (2%)	0	100	100
1	C	141/153 (92%)	140 (99%)	1 (1%)	0	100	100
All	All	421/459 (92%)	415 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/117 (96%)	110 (98%)	2 (2%)	64	30
1	B	110/117 (94%)	98 (89%)	12 (11%)	7	0
1	C	112/117 (96%)	109 (97%)	3 (3%)	50	14
All	All	334/351 (95%)	317 (95%)	17 (5%)	28	4

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LYS
1	A	136	LYS
1	B	2	THR
1	B	23	LYS
1	B	24	GLU
1	B	25	SER
1	B	75	LYS
1	B	78	GLU
1	B	91	LYS
1	B	99	ILE
1	B	102	SER
1	B	107	SER
1	B	137	THR
1	B	139	ASN
1	C	15	GLN
1	C	125	HIS
1	C	136	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	C	15	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSX	A	111	1	4,6,7	1.57	1 (25%)	2,6,8	1.13	0
1	CSX	B	111	1	4,6,7	0.85	0	2,6,8	1.78	1 (50%)
1	CSX	C	111	1	4,6,7	1.16	1 (25%)	2,6,8	1.53	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	A	111	1	-	0/1/5/7	0/0/0/0
1	CSX	B	111	1	-	0/1/5/7	0/0/0/0
1	CSX	C	111	1	-	0/1/5/7	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	111	CSX	CA-C	2.18	1.53	1.50
1	A	111	CSX	CA-C	2.82	1.54	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	111	CSX	CA-CB-SG	-2.33	108.23	113.47
1	C	111	CSX	O-C-CA	-2.04	119.37	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	801	3	4,4,4	0.94	0	6,6,6	0.45	0
2	SO4	B	802	3	4,4,4	1.47	1 (25%)	6,6,6	1.07	0
2	SO4	C	803	3	4,4,4	0.97	0	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	801	3	-	0/0/0/0	0/0/0/0
2	SO4	B	802	3	-	0/0/0/0	0/0/0/0
2	SO4	C	803	3	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	802	SO4	O1-S	2.27	1.57	1.45

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	145/153 (94%)	-0.18	2 (1%) 75 75	11, 15, 30, 41	0
1	B	143/153 (93%)	1.05	22 (15%) 2 2	14, 29, 58, 71	0
1	C	145/153 (94%)	-0.27	1 (0%) 87 86	9, 13, 26, 46	0
All	All	433/459 (94%)	0.19	25 (5%) 24 22	9, 17, 46, 71	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	9.3
1	B	137	THR	6.5
1	B	139	ASN	5.3
1	B	23	LYS	4.8
1	B	125	HIS	4.1
1	B	103	VAL	4.1
1	B	68	SER	3.7
1	B	127	GLY	3.6
1	B	104	ILE	3.5
1	B	71	HIS	3.4
1	B	70	LYS	3.3
1	B	138	GLY	3.3
1	B	26	ASN	3.0
1	A	136	LYS	3.0
1	B	28	PRO	2.8
1	C	1	ALA	2.7
1	B	109	ASP	2.7
1	B	2	THR	2.6
1	B	25	SER	2.5
1	B	69	ARG	2.5
1	B	76	ASP	2.2
1	B	81	VAL	2.2
1	B	82	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	23	LYS	2.1
1	B	140	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
1	CSX	C	111	7/8	0.98	0.06	-	8,10,14,37	0
1	CSX	A	111	7/8	0.96	0.10	-	12,13,17,42	0
1	CSX	B	111	7/8	0.94	0.10	-	29,29,36,47	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	SO4	B	802	5/5	0.92	0.17	0.25	30,30,43,50	0
2	SO4	A	801	5/5	0.98	0.06	-0.16	12,13,17,19	0
2	SO4	C	803	5/5	0.98	0.06	-0.65	10,11,17,18	0
3	ZN	A	202	1/1	1.00	0.02	-0.84	11,11,11,11	0
3	ZN	C	206	1/1	1.00	0.02	-1.06	10,10,10,10	0
3	ZN	B	204	1/1	0.81	0.10	-1.19	33,33,33,33	0
3	ZN	B	203	1/1	0.91	0.09	-1.27	25,25,25,25	0
3	ZN	C	205	1/1	1.00	0.02	-3.38	9,9,9,9	0
3	ZN	A	201	1/1	1.00	0.02	-4.60	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.